

References

3.1 INTRODUCTION

Intergovernmental Panel on Climate Change (IPCC). (1997). Houghton J.T., Meira Filho L.G., Lim B., Treanton K., Mamaty I., Bonduki Y., Griggs D.J. and Callander B.A. (Eds). *Revised 1996 IPCC Guidelines for National Greenhouse Inventories*. IPCC/OECD/IEA, Paris, France.

Intergovernmental Panel on Climate Change (IPCC). (2000). Penman J., Kruger D., Galbally I., Hiraishi T., Nyenzi B., Emmanuel S., Buendia L., Hoppaus R., Martinsen T., Meijer J., Miwa K., and Tanabe K. (Eds). *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*. IPCC/OECD/IEA/IGES, Hayama, Japan.

3.2 FOREST LAND

Andreae M.O., and Merlet P. (2002). Emission of trace gases and aerosols from biomass burning. *Global Biogeochemical Cycles*, 15 (4): pp. 955-966.

Bernoux M., Carvalho M. da CS, Volkoff B., Cerri C.C., Carvalho M. da CS. (2002). Brazil's soil carbon stocks. *Soil Science Society of America Journal*, 66(3), pp. 888-896.

Bhatti J.S., Apps M.J., and Jiang H. (2001). Examining the carbon stocks of boreal forest ecosystems at stand and regional scales. In: Lal R. et al. (eds.) *Assessment Methods for Soil Carbon*, Lewis Publishers, Boca Raton FL. pp. 513-532.

Brown S. (1997). Estimating biomass and biomass change of tropical forests: a primer. FAO-Forestry-Paper 134. Forest Resources Assessment Publication, pp. 55.

Brumme R., Borken W., and Finke S. (1999). Hierarchical control on nitrous oxide emission in forest ecosystem. *Global Biogeochemical Cycles*, 13: pp. 1137-1148.

Butterbach Bahl K., Breuer L., Gasche R., Willibald G., and Papen H. (2002). Exchange of trace gases between soils and the atmosphere in Scots pine forest ecosystems of the northeastern German lowlands. 1. Fluxes of N₂O, NO/NO₂ and CH₄ at forest sites with different N-deposition. *Forest Ecology and Management*, 167: pp. 123-134.

Chojnacky D.C. and Heath L.S. (2002). Estimating down deadwood from FIA forest inventory variables in Maine. *Environmental-Pollution*, 116: Suppl. 1 : S25-S30.

Conen F., Dobbie K.E., and Smith K.A. (2000). Predicting N₂O emissions from agricultural land through related parameters. *Global Change Biology*, 5: pp. 471-426.

Crutzen P.J., and Andreae M.O. (1990). Biomass burning in the tropics: impact on atmospheric chemistry and biogeochemical cycles. *Science*, 4988: pp. 1669-1678.

Obale-Ebanga F., Sevink J., de Groot W., and Nolte C. (2003). Myths of slash and burn on physical degradation of savannah soils: Impacts on Vertisols in North Cameroon. *Soil-Use and Management*, 19 (1) : pp. 83-86.

Dixon R.K., Brown S., Houghton R. A., Solomon A. M., Trexler M. C., and Wisniewski J. (1994). Carbon pools and flux of global forest ecosystems. *Science* 263(1544): pp. 185-190.

Hay G.J., Marceau D.J., Dube P., and Bouchard A. (2001). A multi-scale framework for landscape analysis: Object-specific analysis and upscaling. *Landscape-Ecology*, 16: pp. 471-490.

Duvall M.D., and Grigal D.F. (1999). Effects of timber harvesting on coarse woody debris in red pine forests across the Great Lakes states, USA. *Canadian Journal of Forest Research* 29 (12): pp. 1926-1934.

Food and Agriculture Organisation (FAO) (2001). *Global Forest Resources Assessment 2000. Main Report*. Forestry Paper 140, FAO, Rome: pp. 479.

FAO (1995). *Forest Resources Assessment 1990: Global Synthesis*. Forestry Papers 124, FAO, Rome: pp. 44.

Filipchuk A.N., Strakhov V.V., Borisov B.A. et al. (2000). A Brief National Overview on Forestry Sector and Wood Products: Russian Federation. UN ECE, FAO. New York, Geneva. ECE/TIM/SP/18 (in Russian): pp. 94.

Fisher R.F. and Binkley D. (2000). *Ecology and management of forest soils*. John Wiley & Sons. New York: pp. 489.

- Harmon M.E. and Marks B. (2002). Effects of silvicultural practices on carbon stores in Douglas-fir-western hemlock forests in the Pacific Northwest, USA: results from a simulation model. *Canadian Journal of Forest Research*, 32 (5): pp. 863-877.
- Harmon M. E., Krankina O.N., Yatskov M., and Matthews E. (2001). Predicting broad-scale carbon stores of woody detritus from plot-level data. In: Lai, R., J. Kimble, B. A. Stewart (eds.). *Assessment Methods for Soil Carbon*, CRC Press, New York, pp. 533-552.
- Hoover C.M., Birdsey R.A., and Heath L.S. (2000). How to estimate carbon sequestration on small forest tracts. *Journal Forestry*, 98 (9): pp. 13-19.
- Houghton R.A. (1999). The annual net flux of carbon to the atmosphere from changes in land-use 1850-1990. *Tellus*, 51B: pp.298-313.
- Ilic J., Boland D., McDonald M., Downes G., and Blakemore P. (2000). Woody density Phase 1 – state of Knowledge. National Carbon Accounting System, Technical Report No 18. Australian greenhouse Office.
- Intergovernmental Panel on Climate Change (IPCC). (1997). Houghton J.T., Meira Filho L.G., Lim B., Treanton K., Mamaty I., Bonduki Y., Griggs D.J. and Callander B.A. (Eds). *Revised 1996 IPCC Guidelines for National Greenhouse Inventories*. IPCC/OECD/IEA, Paris, France.
- Intergovernmental Panel on Climate Change (IPCC). (2000). Penman J., Kruger D., Galbally I., Hiraishi T., Nyenzi B., Emmanuel S., Buendia L., Hoppaus R., Martinsen T., Meijer J., Miwa K., and Tanabe K. (Eds). *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*. IPCC/OECD/IEA/IGES, Hayama, Japan.
- Jackson R.B., Banner J.L., Jobbagy E.B., Pockman W.T., and Wall D.H. (2002). Ecosystem carbon loss with woody plan invasion of grasslands. *Nature*, 418: pp. 623-626.
- Jobbagy E.G., and Jackson R.B. (2000). The vertical distribution of soil organic carbon and its relation to climate and vegetation. *Ecological Applications*, 19(2): pp. 423-436.
- Johnson D.W., and Curtis P.S. (2001). Effects of forest management on soil C and N storage: meta analysis. *Forest Ecology and Management*, 140: pp. 227-238.
- Johnson D.W., Knoepp J.D., Swank W.T. (2002). Effects of forest management on soil carbon: results of some long-term resampling studies. *Environment Pollution*, 116: pp. 201-208.
- Johnson M.G., Levine E.R., and Kern J.S. (1995). Soil organic matter: distribution, genesis, and management to reduce greenhouse gas emissions. *Water, Air and Soil Pollution* 82: pp. 593-615.
- Jacinthe P.A., Lal R., Kimble J.M. (2002). Carbon dioxide evolution in runoff from simulated rainfall on long-term no-till and plowed soils in Southwestern Ohio. *Soil Tillage Research* 66 (1): pp. 23-33.
- Kirschbaum-MUF (2000). How should forest fires be treated in the National Greenhouse Gas Inventory? *Australian-Forestry*, 63(2): pp. 136-141.
- Klemetsson L., Klemetsson A.K., Moldan F., and Weslien P. (1997). Nitrous oxide emission from Swedish forest soils in relation to liming and simulated increased N-deposition. *Biology & Fertility of Soils* 25: pp. 290-295.
- Koehl M. (2000). Reliability and comparability of TBFRA 2000 results. In: TBFRA 2000, Geneva, UN-ECE/FAO: pp. 27-61.
- Kramer H. (1982). *Nutzungsplanung in der Forsteinrichtung*. Sauerländer Verlag, Frankfurt: pp.128.
- Kurz W. A. and Apps M. J. (1992). Atmospheric carbon and Pacific Northwest Forests. In: Wall, G. (ed.). *Implication of climate change for Pacific Northwest Forest management*. Un. of Waterloo. Dept. of Geography. *Occasional Paper* No. 15: pp. 69-80.
- Lafleur P.M., Roulet N.T., Bubier J.L., Frohling S., and Moore T.R. (2003). Inter-annual variability in the peatland-atmosphere carbon dioxide exchange at an ombrotrophic bog - art. no. 1036. *Global-Biogeochemical-Cycles*, 17 (2): pp. 00-00.
- Laitat É., Karjalainen T., Loustau D., and Lindner M. (2000). Introduction: Towards an integrated scientific approach for carbon accounting in forestry. *Biotechnologie, Agronomie, Société et Environnement* 2000 4(4): pp. 315-319.
- Izaurre R.C., Rosenberg N.J., and Lal R. (2001). Mitigation of climatic change by soil carbon sequestration: issues of science, monitoring, and degraded lands. *Advances-in-Agronomy*, 70: pp. 1-75.

- Lehtonen X, Mäkipää R., Heikkinen J., Sievänen R., and Liski J. (2003). Biomass expansion factors (BEFs) for Scots pine, Norway spruce and birch according to stand age for boreal forests, *Forest Ecology and Management, In Press, Corrected Proof, Available online 24 September 2003*.
- Levine J.S., Wesley III R.C., Winstead E.L., Thinehart R.P., Cahoon Jr. D.R., Sebacher D.K., Sebacher S., and Stocks B.J. (1991). Biomass burning: combustion emissions, satellite imagery, and biogenic emissions. In: J.S. Levine (Ed.) *Global Biomass Burning* MIT Press, Cambridge: pp. 264-271.
- Li Z., Kurz W.A., Apps M.J., and Beukema S.J. (2003) Belowground biomass dynamics in the Carbon Budget Model of the Canadian Forest Sector: recent improvements and implications for the estimation of NPP and NEP. *Canadian Journal of Forest Research* 33 (1): pp. 126-136.
- Li C.S., Aber J, Stange F, Butterbach Bahl K., and Papen H. (2000). A process-oriented model of N₂O and NO emissions from forest soils: 1. Model development. *Journal of Geophysical Research Atmospheres*. 105 (D4): pp. 4369-4384.
- Liski J., Pussinen A., Pingoud K., Makipaa R., Karjalainen T. (2001). Which rotation length is favourable to carbon sequestration? *Canadian Journal of Forest Research* 31: pp. 2004-2013.
- Löewe H, Seufert G., and Raes F. (2000). Comparison of methods used within Member States for estimating CO₂ emissions and sinks according to UNFCCC and EU Monitoring Mechanism: forest and other wooded land. *Biotechnologie, Agronomie, Société et Environnement* 2000 4(4): pp. 315-319.
- Martikainen P.J., Nykanen H., Alm J., and Silvola J. (1995). Change in fluxes of carbon dioxide, methane and nitrous oxide due to forest drainage of mire sites of different trophic. *Plant & Soil* 169: pp. 571-577.
- McKenzie N.J., Cresswell H.P., Ryan P.J., and Grundy M. (2000). Opportunities for the 21st century: Expanding the horizons for soil, plant, and water analysis. *Communications in Soil Science and Plant Analysis* 31: pp. 1553-1569.
- Mosier A. and Kroeze C. (1999). Contribution of agroecosystems to the global atmospheric N₂O budget. Proceedings of International workshop on reducing N₂O emission from agroecosystems, Banff, Canada, March 1999.
- Mosier A. R., Delgado J.A., and Keller M. (1998). Methane and nitrous oxide fluxes in an acid Oxisol in Western Puerto Rico: Effects of tillage, liming and fertilization. *Soil Biology & Biochemistry* 30: pp. 2087-2098.
- Page-Dumroese D, Jurgensen M, Elliot W, *et al.* (2000). Soil quality standards and guidelines for forest sustainability in northwestern North America. *Forest Ecology Management* 138 (1-3): pp. 445-462.
- Papen H., and Butterbach-Bahl K. (1999). A 3-year continuous record of nitrogen trace gas fluxes from untreated and limed soil of a N-saturated spruce and beech forest in Germany - 1. N₂O emissions. *Journal of Geophysical Research* 104: pp. 18487-18503.
- Paul K.I., Polglase P.J., Nyakuengama J.G., and Khanna P.K. (2002). Change in soil carbon following afforestation. *Forest Ecology and Management* 168: pp. 241-257.
- Polglase P.J., Paul K.I., Khanna P.K., Nyakuengama J.G., O'Connell A.M., Grove T.S., and Battaglia M. (2000). Change in soil Carbon Following Afforestation or Reforestation. National Carbon Accounting system technical report no. 20 Australian Greenhouse Gas Office, Canberra.
- Post W.M. and Kwon K.C. (2000). Soil carbon sequestration and land-use changes: processes and potential. *Global Change Biology* 6: pp. 317-327.
- Pregitzer K.S. (2003). Woody plants, carbon allocation and fine roots. *New Phytologist* 158 (3): pp. 421-424.
- Renault P. (1999). Les modèles opérationnels d'émission de N₂O par les sols aux échelles régionales. *C.R. Acad. Agri. Fr.* 85, 6 : pp. 163-176.
- Richter D.D., Markewitz D., Trumbore S.E. and Wells C.B. (1999). Rapid accumulation and turnover of soil carbon in a re-establishing forest. *Nature* 400: pp. 56-58.
- Schelhaas M.J., Varis S., and Schuck A. (2001). Database on Forest Disturbances in Europe (DFDE), European Forest Institute, Joensuu, Finland, <http://www.efi.fi/projects/dfde/>.
- Scott N.A., Tate K.R., Giltrap D.J., *et al.* (2002). Monitoring land-use change effects on soil carbon in New Zealand: quantifying baseline soil carbon stocks. *Environmental Pollution* 116: pp. 167-186.
- Siltanen *et al.* (1997). A soil profile and organic carbon data base for Canadian forest and tundra mineral soils. Natural Resources Canada, Canadian Forest Service, Northern Forestry Centre, Edmonton, Alberta.

- Smith J.E., and Heath L.S. (2002). A model of forest floor carbon mass for United States forest types. General Technical Report, USDA Forest Service, Northeastern Research Station, Newtown Square, PA. In press.
- Smith K. A., Dobbie K.E., Ball B.C., Bakken L.R., Sitaula B.K., Hansen S., Brumme R., Borken W., Christensen S., Priemé A., Fowler D., MacDonald J.A., Skiba U., Klemmedtsson L., Kasimir-Klemmedtsson A., Degórska A., and Orlanski P. (2000). Oxidation of atmospheric methane in Northern European soils, comparison with other ecosystems, and uncertainties in the global terrestrial sink. *Global Change Biology* 6: pp. 791-803.
- Smith K.A., Bouwman L., and Braatz B. (1999). Nitrous oxide: direct emissions from agricultural soils. Background paper for IPCC Workshop on Good Practice in Inventory Preparation : Agricultural sources of methane and nitrous oxide, 24-26 February 1999, Wageningen, The Netherlands.
- Spies T.A., Franklin J.F., and Thomas T.B. (1988). Coarse woody debris in Douglas-fir forests of Western Oregon and Washington. *Ecology* 6: pp. 1689-1702.
- Stange F., Butterbach-Bahl K., Papen H., *et al.* (2000). A process-oriented model of N₂O and NO emissions from forest soils 2. Sensitivity analysis and validation. *Journal of Geophysical Research Atmosphere* 105 (D4): pp. 4385-4398.
- Thormann M.N., Szumigalski A.R., Bayley S.E. (1999). Aboveground peat and carbon accumulation potentials along a bog-fen-marsh wetland gradient in southern boreal Alberta, Canada. *Wetlands* 19 (2): pp. 305-317.
- Tremblay S., Ouimet R. and Houle D. (2002). Prediction of organic carbon content in upland forest soils of Quebec, Canada. *Can. J. For. Res.* 32: pp. 903-914.
- UN-ECE/FAO (20000)
- Vogt K.A., Vogt D.J., Pamiotto P.A., Boon P., O'Hara J., and Asbjornsen H. (1996). Review of root dynamics in forest ecosystems grouped by climate, climatic forest type, and species. *Plant and Soil* 187: pp. 159-219.
- Yavitt J. B., Fahey T.J., and Simmons J.A. (1997). Methane and carbon dioxide dynamics in a northern hardwood ecosystem. *Soil Science Society of America Journal* 59: pp. 796-804.
- Zagreev V.V., Sukhikh B.I., Shvidenko A.Z., Gusev N.N., and Moshkalev A.G. (1993). The All-Union Standards for Forest Inventory. *Kolos*, Moscow, p. 495. (In Russian).
- Zoltai S.C. and Vitt D.H. (1995). Canadian wetlands – environmental gradients and classification. *Vegetation* 118 (1-2): pp. 131-137.

3.3 CROPLAND

- Armentano T.V. and Menges E.S. (1986). Patterns of change in the carbon balance of organic soil-wetlands of the temperate zone. *Journal of Ecology* 74: pp. 755-774.
- Barbosa R.I. and Fearnside P.M. (1996). Pasture burning in Amazonia: Dynamics of residual biomass and the storage and release of aboveground carbon. *Journal of Geophysical Research (Atmospheres)* 101(D20): pp. 25847-25857.
- Bernoux M., Carvalho M.D.S., Volkoff B., and Cerri C.C. (2002). Brazil's soil carbon stocks. *Soil Science Society of America Journal* 66: pp. 888-896.
- Conant R.T. and Paustian K. (2002). Spatial variability of soil organic carbon in grasslands: implications for detecting change at different scales. *Environmental Pollution* 116: pp. 127-135.
- Davidson E. A. and Ackerman I.L. (1993). Changes in soil carbon inventories following cultivation of previously untilled soils. *Biogeochemistry* 20: pp. 161-164.
- Dixon R.K., Winjum J.K., and Schroeder P.E. (1993). Conservation and sequestration of carbon: the potential of forests and agroforest management practices. *Global Environmental Change* 3: pp. 159-173.
- Dobbie K.E., McTaggart I.P., and Smith K.A. (1999). Nitrous oxide emissions from intensive agricultural systems: Variations between crops and seasons, key driving variables, and mean emission factors. *Journal of Geophysical Research-Atmospheres* 104: pp. 26891-26899.
- Fearnside P.M. (1990). Fire in the tropical rainforest of the Amazon basin. In: Fire in the tropical biota Goldammer, J.G.(ed). Ecological Studies 84, Springer-Verlag, N.Y. : pp. 106-116.

- Fearnside P.M. (2000). Global warming and tropical land-use change: greenhouse gas emissions from biomass burning, decomposition and soils in forest conversion, shifting cultivation and secondary vegetation. *Climatic Change* 46: pp. 115-158.
- Firestone M.K. and Davidson E.A. (1989) Microbiological basis of NO and N₂O production and consumption in soil. In: Andreae M.O. and D. S. Schimel (eds) . Exchange of Trace Gases Between Terrestrial Ecosystems and the Atmosphere, Wiley, N.Y: pp. 7-21.
- Intergovernmental Panel on Climate Change (IPCC). (1997). Houghton J.T., Meira Filho L.G., Lim B., Treanton K., Mamaty I., Bonduki Y., Griggs D.J. and Callander B.A. (Eds). *Revised 1996 IPCC Guidelines for National Greenhouse Inventories*. IPCC/OECD/IEA, Paris, France.
- Intergovernmental Panel on Climate Change (IPCC). (2000). Penman J., Kruger D., Galbally I., Hiraishi T., Nyenzi B., Emmanuel S., Buendia L., Hoppaus R., Martinsen T., Meijer J., Miwa K., and Tanabe K. (Eds). *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*. IPCC/OECD/IEA/IGES, Hayama, Japan.
- Intergovernmental Panel on Climate Change (IPCC). (2000b). Watson R., Noble I.R., Bolin B., Ravindranath, N.H., Verardo D.J. and Dokken D.J. (Eds) *Land use, Land-use Change, and Forestry: A Special Report*. Cambridge University Press. Cambridge, UK.
- Jobbagy E.G. and Jackson R.B. (2000). The vertical distribution of soil organic carbon and its relation to climate and vegetation. *Ecological Applications* 19(2): pp. 423-436.
- Klemedtsson L., Klemedtsson A.K., Moldan F., and Weslien P. (1997). Nitrous oxide emission from Swedish forest soils in relation to liming and simulated increased N-deposition. *Biology & Fertility of Soils* 25: pp. 290-295.
- Masera O. R., Garza-Caligaris J.F., Kanninen M., Karjalainen T., Liski J., Nabuurs G.J., Pussinen A., de Jong B.H.J., and Mohren G.M.J. (2003). Modeling carbon sequestration in afforestation, agroforestry and forest management projects. *Ecological Modelling* 164: pp. 177-199.
- McGill W. B. (1996). Review and classification of ten soil organic matter models. In: Powlson D.S., Smith P., and Smith J.U. (eds.). *Evaluation of Soil Organic Matter Models Using Existing Long-Term Datasets*. Springer-Verlag, Heidelberg: pp. 111-132.
- Neill C., Melillo J.M., Steudler P.A., Cerri C.C., de Moraes J.F.L., Piccolo M.C., and Brito M. (1997). Soil carbon and nitrogen stocks following forest clearing for pasture in the southwestern Brazilian Amazon. *Ecological Applications* 7 : pp. 1216-1225.
- Nusser S.M. and Goebel J.J. (1997). The National Resources Inventory: a long-term multi-resource monitoring programme. *Environmental and Ecological Statistics* 4: pp. 181-204.
- Ogle S. M., Breidt F.J., Eve M.D., and Paustian K. (2003). Uncertainty in estimating land use and management impacts on soil organic carbon storage for U.S. agricultural lands between 1982 and 1997. *Global Change Biology* (in press).
- Paul E.A., Paustian K., Elliott E.T. and Cole C.V. (eds) (1997) Soil organic matter in temperate agroecosystems: Long-term Experiments in North America. CRC Press, Boca Raton, 414 p.
- Schroeder P. (1994). Carbon storage benefits of agroforestry systems. *Agroforestry Systems* 27: pp. 89-97.
- Schroth G., D'Angelo S.A., Teixeira W.G., Haag D., and Lieberei R. (2002). Conversion of secondary forest into agroforestry monoculture plantations in Amazonia: consequences for biomass, litter and soil carbon stocks after 7 years. *Forest Ecology and Management* 163: pp. 131-150.
- Smith P., Powlson D.S., Smith J.U., and Elliott E.T. (eds) (1997). Evaluation and comparison of soil organic matter models. Special Issue, *Geoderma* 81: pp. 1-225.
- Smith P., Powlson D., Glendinning, M. (1996). Establishing a European GCTE soil organic matter network (SOMNET). In: Powlson D.S., Smith P., and Smith J.U. (eds.), *Evaluation of Soil Organic Matter Models Using Existing Long-Term Datasets*. NATO ASI Series, Vol 38, Springer-Verlag, Berlin: pp. 81-97.
- Soil Organic Matter Network (SOMNET) (1996). Model and Experimental Metadata. GCTE Task 3.3.1. Smith P., Smith J.U., and Powlson D.S. (eds). *Global Change and Terrestrial Ecosystems Report No 7*. GCTE Focus 3 Office, Wallingford, UK. 255 p.

3.4 GRASSLAND

- Anderson D.J., Perry R.A., and Leigh J.H. (1972). Some perspectives on shrub/environment interactions. In: McKell C.M., Blaisdell J.P., Goodon J.R. (eds), *Wildland Shrubs – Their Biology and Utilization*.). USDA Forest Service, General Tech. Report INT-1.
- Armentano T.V. and Menges E.S. (1986). Patterns of change in the carbon balance of organic soil-wetlands of the temperate zone. *Journal-of-Ecology*. 74, 3: pp. 755-774.
- Baldocchi D., Kelliher F.M., Black T.A., Jarvis P. (2000). Climate and vegetation controls on boreal zone energy exchange. *Global Change Biology*,-Supplement. 6: Supplement 1,pp. 69-83.
- Baldocchi D., Falge E., Gu L.H., Olson R., Hollinger D., Running S., Anthoni P., Bernhofer C., Davis K., Evans R., Fuentes J., Goldstein A., Katul G., Law B., Lee X.H., Malhi Y., Meyers T., Munger W., Oechel W., Pilegaard K., Schmid H.P., Valentini R., Verma S., Vesala T., Wilson K., and Wofsy S. (2001). FLUXNET: A new tool to study the temporal and spatial variability of ecosystem-scale carbon dioxide, water vapor, and energy flux densities. *Bulletin of the American Meteorological Society* 82: pp. 2415-2434.
- Barbosa R.I. and Fearnside P.M. (1996). Pasture burning in Amazonia: Dynamics of residual biomass and the storage and release of aboveground carbon. *Journal of Geophysical Research (Atmospheres)* 101(D20): pp. 25,847-25,857.
- Cairns M.A., Brown S., Helmer E.H., and Baumgardner G.A. (1997). Root biomass allocation in the world's upland forests. *Oecologia* 111: pp. 1-11.
- Conant R.T. and Paustian K. (2002a). Spatial variability of soil organic carbon in grasslands: implications for detecting change at different scales. *Environmental Pollution* 116: pp. 127-135.
- Conant R.T. and Paustian K. (2002b). Potential soil carbon sequestration in overgrazed grassland ecosystems. *Global Biogeochemical Cycles* 16: pp. 90_1-90_9.
- Conant R.T., Paustian K., and Elliott E.T. (2001). Grassland management and conversion into grassland: Effects on soil carbon. *Ecological Application* 11: pp. 343-355.
- Davidson E. A. and Ackerman I.L. (1993). Changes in soil carbon inventories following cultivation of previously untilled soils. *Biogeochemistry* 20: pp. 161–164.
- Delmas R.A., Loudjana P., Podaire A., and Menaut J.C. (1991). Biomass burning in Africa: An assessment of annually burnt biomass. In; Levine J.S. (ed), *Global Biomass Burning: Atmosphere, Climatic and Biosphere Implications*, MIT Press, Cambridge, Mass.: pp. 147-154.
- Ellert B.H., Janzen H.H., and McConkey B.G. (2001). Measuring and comparing soil carbon storage. In: R. Lal, J.M. Kimble, R.F. Follett and B.A. Stewart (eds.). *Soil Management for Enhancing Carbon Sequestration*. CRC Press, Boca Raton, FL.: pp. 593-610.
- Fearnside P.M. (1990). Fire in the tropical rainforest of the Amazon basin. pp 106-116 *In: Goldammer J.G.(ed.) Fire in the tropical biota*. Ecological Studies 84, Springer-Verlag, N.Y. 497 p.
- Fearnside P.M. (2000). Global warming and tropical land-use change: greenhouse gas emissions from biomass burning, decomposition and soils in forest conversion, shifting cultivation and secondary vegetation. *Climatic Change* 46: pp. 115-158.
- Guo L.B. and Gifford R.M. (2002). Soil carbon stocks and land use change: a meta analysis. *Global Change Biology* 8: pp. 345-360.
- Hao W.M., Darold E.W., Olbu G., and Baker S.P. (1996). Emissions of CO₂, CO and hydrocarbons from fires in diverse African savanna ecosystems. *Journal of Geophysical Research*, Vol 101, No. D19: pp. 23577-23584.
- Intergovernmental Panel on Climate Change (IPCC). (1997). Houghton J.T., Meira Filho L.G., Lim B., Treanton K., Mamaty I., Bonduki Y., Griggs D.J. and Callander B.A. (Eds). *Revised 1996 IPCC Guidelines for National Greenhouse Inventories*. IPCC/OECD/IEA, Paris, France.
- Intergovernmental Panel on Climate Change (IPCC). (2000). Penman J., Kruger D., Galbally I., Hiraishi T., Nyenzi B., Emmanuel S., Buendia L., Hoppaus R., Martinsen T., Meijer J., Miwa K., and Tanabe K. (Eds). *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*. IPCC/OECD/IEA/IGES, Hayama, Japan.

- Intergovernmental Panel on Climate Change (IPCC). (2000b). Watson R., Noble I.R., Bolin B., Ravindranath, N.H., Verardo D.J. and Dokken D.J. (Eds) *Land use, Land-use Change, and Forestry: A Special Report*. Cambridge University Press. Cambridge, UK.
- Jackson R.B., Banner J.L., Jobbagy E.B., Pockman W.T., and Wall D.H. (2002). Ecosystem carbon loss with woody plant invasion of grasslands. *Nature* 418: pp. 623-626.
- Jackson R.B., Canadell J., Ehleringer J.R., Mooney H.A., Sala O.E., and Schulze E.D. (1996). A global analysis of root distributions for terrestrial biomes. *Oecologia* 108: pp. 389-411.
- Kuhlbusch T.A.J., Andreae M.O., Cachier H., Goldammer J.G., Lacaux J.P., Shea R., and Crutzen P.J. (1996). Black carbon formation by savanna fires: Measurements and implications for the global carbon cycle. *Journal of Geophysical Research*, Vol 101, No D19: pp. 23651-23665.
- Lacaux J.P., Delmas R., and Jambert C. (1996). NO_x emissions from African savanna fires. *Journal of Geophysical Research*, Vol 101, No D19: pp. 23585-23595.
- Lober J. M., Scharffé D.H., Hao W.M., Kuhlbusch T.A., Seuwen R., Warneck P., and Crutzen P.J. (1993). Experimental evaluation of biomass burning emissions: Nitrogen carbon containing compounds. In: Levine J.S. (ed) *Global Biomass Burning*, MIT Press.
- Lober J.M. and Warnatz J. (1993). Emissions from combustion process in vegetation. In: Crutzen P.J. and Goldammer J.G., *Fire in the environment*, John Wiley, New York: pp. 15-37.
- McGill W. B. (1996). Review and classification of ten soil organic matter models. In: Powlson D.S., Smith P., and Smith J.U. (eds.). *Evaluation of Soil Organic Matter Models Using Existing Long-Term Datasets*. Springer-Verlag, Heidelberg: pp. 111-132.
- Milchunas D.G. and Lauenroth W.K. (1993). Quantitative effects of grazing on vegetation and soils over a global range of environments. *Ecological Monographs* 63: pp. 327-366.
- Neill C., Melillo J.M., Steudler P.A., Cerri C.C., de Moraes J.F.L., Piccolo M.C. and Brito M. (1997). Soil carbon and nitrogen stocks following forest clearing for pasture in the southwestern Brazilian Amazon. *Ecological Applications* 7 : pp. 1216-1225.
- Nihlgård B. (1972). Plant biomass, primary production and distribution of chemical elements in abeech and replanted spruce forest in South Sweden. *Oikos* 23: pp. 69-81.
- Nusser S.M. and Goebel J.J. (1997). The National Resources Inventory: a long-term multi-resource monitoring programme. *Environmental and Ecological Statistics* 4: pp. 181-204.
- Ogle S. M., Breidt F. J., Eve M. D., and Paustian K. (2003). Uncertainty in estimating land use and management impacts on soil organic carbon storage for U.S. agricultural lands between 1982 and 1997. *Global Change Biology* (in press).
- Ojima D.S., Parton W.J., Schimel D.S., Scurlock J.M.O., and Kittel T.G.F. (1993). Modeling the effects of climatic and CO₂ changes on grassland storage of soil C. *Water, Air, and Soil Pollution* 70: pp. 643-657.
- Olson R. J., Scurlock, J. M. O., Prince S. D., Zheng D. L., and Johnson K. R. (eds.) (2001). NPP Multi-Biome: NPP and Driver Data for Ecosystem Model-Data Intercomparison. Data set. Available on-line [<http://www.daac.ornl.gov>] from the Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, U.S.A.
- Pacala S.W., Hurtt G.C., Baker D., Peylin P., Houghton R.A., Birdsey R.A., Heath L., Sundquist E.T., Stallard R.F., Ciais P., Moorcroft P., Caspersen J.P., Shevliakova E., Moore B., Kohlmaier G., Holland E., Gloor M., Harmon M.E., Fan S.M., Sarmiento J.L., Goodale C.L., Schimel D., and Field C.B. (2001). Consistent land- and atmosphere-based US carbon sink estimates. *Science* 292: pp. 2316-2320.
- Scholes R. J., Kendall J., and Justice C.O. (1996). The quantity of biomass burned in southern Africa, *Journal of Geophysical Research*, Vol 101. NO D19: pp. 23677-23682.
- Smith P., Powlson D. S., Glendining M. J., and Smith J. O. U. (1997). Potential for carbon sequestration in European soils: preliminary estimates for five scenarios using results from long-term experiments. *Global Change Biology* 3: pp. 67-79.
- Veldkamp E. (2001). Changes in soil carbon stocks following conversion of forest to pasture in the tropics. In: Holland E.A. (ed.): *Notes from Underground: Soil Processes and Global Change*. NATO ASI Series Berlin: Springer: in press.

Ward D.E., Hao W.M., Susott R.A., Babbitt R.E., Shea R.W., Kauffman J.B. and Justice C.O. (1996). Effect of fuel composition on combustion efficiency and emission factors for African savanna ecosystems. *Journal of Geophysical Research*, Vol 101, No D19: pp. 23569-23574.

3.5 WETLANDS

Alm J., Saarnio S., Nykanen H., Silvola J., and Martikainen P. J. (1999). Winter CO₂, CH₄ and N₂O fluxes on some natural and drained boreal peatlands. *Biogeochemistry* 44: pp. 163-186.

Intergovernmental Panel on Climate Change (IPCC). (1997). Houghton J.T., Meira Filho L.G., Lim B., Treanton K., Mamaty I., Bonduki Y., Griggs D.J. and Callander B.A. (Eds). *Revised 1996 IPCC Guidelines for National Greenhouse Inventories*. IPCC/OECD/IEA, Paris, France.

Intergovernmental Panel on Climate Change (IPCC). (2000). Penman J., Kruger D., Galbally I., Hiraishi T., Nyenzi B., Emmanuel S., Buendia L., Hoppaus R., Martinsen T., Meijer J., Miwa K., and Tanabe K. (Eds). *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*. IPCC/OECD/IEA/IGES, Hayama, Japan.

Laine J. and Minkinen K. (1996). Effect of forest drainage on the carbon balance of a mire - a case study. *Scandinavian Journal of Forest Research* 11: pp. 307-312.

Laine J., Silvola J., Tolonen K., Alm J., Nykanen H., Vasander H., Sallantausta T., Savolainen I., Sinisalo J., and Martikainen P. J. (1996). Effect of water-level drawdown on global climatic warming - northern peatlands. *Ambio* 25: pp. 179-184.

LUSTRA (2002). Land Use Strategies for Reducing Net Greenhouse Gas Emissions. Annual Report 2002, Uppsala, Sweden. <http://www.sml.slu.se/lustra/index.phtml>

Minkinen K., Korhonen R., Savolainen I., and Laine J. (2002). Carbon balance and radiative forcing of Finnish peatlands 1900-2100 the impact of forestry drainage. *Global Change Biology* 8: pp. 785-799.

Sundh I., Nilsson M., Mikkela C., Granberg G., and Svensson B. H. (2000). Fluxes of methane and carbon dioxide on peat-mining areas in Sweden. *Ambio* 29: pp. 499-503.

3.6 SETTLEMENTS

Nowak D.J. and Crane D.E. (2002). Carbon storage and sequestration by urban trees in the United States. *Environmental Pollution* 116(3): pp. 381-389.

Nowak D.J., Rowntree R.A., McPherson E.G., Sisinni S.M., Kerkmann E.R., and Stevens J.C. (1996). Measuring and analyzing urban tree cover. *Landscape and Urban Planning* 36: pp. 49-57.

APPENDIX 3A.1 HARVESTED WOOD PRODUCTS: BASIS FOR FUTURE METHODOLOGICAL DEVELOPMENT

Burden R.L. and Faires J.D. (2001). *Numerical Analysis*, 7th ed. Brooks/ Cole Publishing. 810 p.

Food and Agriculture Organization 1999. *FAO Yearbook – Forest Products 1997*. FAO Forestry Series No. 42. Rome. 245 pp. <ftp://ftp.fao.org/fo/fon/fons/FOYB1997.pdf>

Food and Agriculture Organization 2002a. FAOSTAT Forestry data. Web site: <http://apps.fao.org/page/collections?subset=forestry> accessed July 1, 2002.

Flugsrud K., Hoem B., Kvingedal E. and Rypdal R. (2001). Estimating the net emissions of CO₂ from harvested wood products. SFT report 1831/200. Norwegian Pollution Control Authority, Oslo. 47 p. <http://www.sft.no/publikasjoner/luft/1831/ta1831.pdf>

Gjesdal S.F.T., Flugsrud K., Mykkelbost T.C., and Rypdal K. (1996). A balance of use of wood products in Norway, Norwegian Pollution Control Authority SFT, Report 96:04, 54 p.

Haynes *et al.* (1990). *An Analysis of the timber situation in the United States: 1989-2040*. USDA Forest Service. Gen. Tech. Rpt. RM-199. 268 p.

- Heath L. S., Birdsey R.A., Row C., and Plantinga A.J. (1996). Carbon pools and fluxes in U.S. forest products. In: Apps M.J. and Price D.T. (eds.), *Forest Ecosystems, Forest Management and the Global Carbon Cycle*. NATO ASI Series, Springer-Verlag, Berlin: pp. 271-278.
- Intergovernmental Panel on Climate Change (IPCC). (1997). Houghton J.T., Meira Filho L.G., Lim B., Treanton K., Mamaty I., Bonduki Y., Griggs D.J. and Callander B.A. (Eds). *Revised 1996 IPCC Guidelines for National Greenhouse Inventories*. IPCC/OECD/IEA, Paris, France.
- Intergovernmental Panel on Climate Change (IPCC) (1998). Evaluating Approaches for Estimating Net Emissions of Carbon Dioxide from Forest Harvesting and Wood Products. Meeting Report, Dakar, Senegal, 5-7 May 1998. Brown S., Lim B. and Schlamadinger B. IPCC/OECD/IEA, Paris, France. See <http://www.ipcc-nggip.iges.or.jp/public/mtdocs/dakar.htm>
- Intergovernmental Panel on Climate Change (IPCC). (2000). Penman J., Kruger D., Galbally I., Hiraishi T., Nyenzi B., Emmanuel S., Buendia L., Hoppaus R., Martinsen T., Meijer J., Miwa K., and Tanabe K. (Eds). *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*. IPCC/OECD/IEA/IGES, Hayama, Japan.
- Karjalainen T., Kellomaki S., and Pussinen A. (1994). Role of Wood-based Products in Absorbing Atmospheric Carbon. *Silva Fennica* 28 (2): pp. 67-80.
- Micales J.A. and Skog K.E. (1997). The decomposition of forest products in landfills. *International Biodeterioration and Biodegradation* 39 (2-3): pp. 145-158.
- Nabuurs G.J. and Sikkema R. (1998). Application and Evaluation of the Alternative IPCC Methods for Harvested Wood Products in the National Communications; Proceedings for the IPCC Expert Meeting on Evaluating approaches for estimating net emissions from harvested wood products, Wageningen, The Netherlands
- Pingoud K., Savolainen I., and Seppälä H. (1996). Greenhouse impact of the Finnish forest sector including forest products and waste management. *Ambio* 25: pp. 318-326.
- Pingoud K., Perälä A.L., and Pussinen A. (2001). Carbon dynamics in wood products. *Mitigation and Adaptation Strategies for Global Change* 6: pp. 91-111.
- Skog K., and Nicholson G. (1998). Carbon Cycling through Wood Products: The Role of Wood and Paper Products in Carbon Sequestration. *Forest Products Journal* 48 (7/8): pp. 75-83.

APPENDIX 3A.2 NON-CO₂ FROM DRAINAGE AND REWETTING OF FOREST SOILS: BASIS FOR FUTURE METHODOLOGY DEVELOPMENT

- Alm J., Saarnio S., Nykanen H., Silvola J. and Martikainen P.J. (1999). Winter CO₂, CH₄ and N₂O fluxes on some natural and drained boreal peatlands. *Biogeochemistry* 44: pp. 163 – 186.
- Bartlett K.B. and Harriss R.C. (1993). Review and assessment of methane emissions from wetlands. *Chemosphere* 26: pp. 261 -320.
- Intergovernmental Panel on Climate Change (IPCC). (1997). Houghton J.T., Meira Filho L.G., Lim B., Treanton K., Mamaty I., Bonduki Y., Griggs D.J. and Callander B.A. (Eds). *Revised 1996 IPCC Guidelines for National Greenhouse Inventories*. IPCC/OECD/IEA, Paris, France.
- Intergovernmental Panel on Climate Change (IPCC). (2000). Penman J., Kruger D., Galbally I., Hiraishi T., Nyenzi B., Emmanuel S., Buendia L., Hoppaus R., Martinsen T., Meijer J., Miwa K., and Tanabe K. (Eds). *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*. IPCC/OECD/IEA/IGES, Hayama, Japan.
- Klemedtsson L., Weslien P., Arnold K., Agren G., Nilsson M., and Hanell B. (2002). Greenhouse gas emissions from drained forests in Sweden. In: Olsson M. (ed.) *Land use strategies for reckoning net greenhouse gas emissions*. Mistra Programme: Progress report 1999 – 2002. Swedish University of Agricultural Sciences. Uppsala: pp. 44 – 67.
- Komulainen V.M., Nykanen H., Martikainen P.J. and Laine J. (1998). Short-term effect of restoration on vegetation change and methane emissions from peatlands drained for forestry in southern Finland. *Can. J. For. Res.* 28: pp. 402 – 411.

- Laine J., Silvola J., Tolonen K., Alm J., Nykanen H., Vasander H., Sallantausta T., Savolainen I., Sinisalo J., and Martikainen P. J. (1996). Effect of water-level drawdown on global climatic warming - northern peatlands. *Ambio* 25: pp. 179-184.
- Martikainen P. J., Nykanen H., Alm J., and Silvola J. (1995). Change in fluxes of carbon dioxide, methane and nitrous oxide due to forest drainage of mire sites of different trophic. *Plant & Soil* 169: pp. 571-577.
- Regina K., Nykänen H., Silvola J., and Martikainen P. J. (1996). Nitrous oxide production in boreal peatlands of different hydrology and nutrient status. In: Northern peatlands in global climatic change. Proceedings of the international workshop. Academy of Finland, Hyytiälä: pp. 158-166.
- Roulet N.T. and Moore T.R. (1995). Methane Emissions from Canadian Peatlands. In: Lal R., Kimble J., Levine E., and Stewart B.A., *Soils and Global Change*, CRC Lewis Publishers, Boca Raton: pp. 153-164.
- Tuittila, E-S., Komulainen, V-M., Vasander, H., Nykänen, H., Martikainen, P.J. & Laine, J. (2000). Methane dynamics of a restored cut-away peatland. *Global Change Biology* 6: 569-581.

APPENDIX 3A.3 WETLANDS REMAINING WETLANDS: BASIS FOR FUTURE METHODOLOGICAL DEVELOPMENT

ORGANIC SOILS MANAGED FOR PEAT EXTRACTION

- Alm, J., S. Saarnio, H. Nykanen, J. Silvola, and P. J. Martikainen. (1999). Winter CO₂, CH₄ and N₂O fluxes on some natural and drained boreal peatlands. *Biogeochemistry* 44:163-186.
- Andriessse, J. P. (1988) Nature and Management of Tropical Peat Soils, FAO SOILS BULLETIN 59, <http://www.fao.org/docrep/x5872e/x5872e04.htm>
- Feehan, J. & O'Donovan, G. (1996) *The Bogs of Ireland*. The Environmental Institute, University College Dublin, Ireland.
- Fey, A., G. Benckiser and J.C.G. Ottow (1999). Emissions of nitrous oxide from a constructed wetland using a groundfilter and macrophytes in waste-water purification of a dairy farm. *Biol Fertil Soils* 29, 354-359.
- Huttunen, J.T., T.S. Vaisanen, S. K. Hellsten, M. Heikkinen, H. Nykanen, H. Jungner, A. Niskanen, M. O. Virtanen, O.V. Lindqvist, O. S. Nenonen, and P.J. Martikainen, (2002), Fluxes of CH₄, CO₂, and N₂O in hydroelectric reservoir Lokka and Porttipahata in the northern boreal zone in Finland, *Global Biogeochemical Cycles*, 16,1.
- Laine, J., and K. Minkinen. (1996). Effect of forest drainage on the carbon balance of a mire - a case study. *Scandinavian Journal of Forest Research* 11:307-312
- Laine, J., J. Silvola, K. Tolonen, J. Alm, H. Nykanen, H. Vasander, T. Sallantausta, I. Savolainen, J. Sinisalo, and P. J. Martikainen. (1996). Effect of water-level drawdown on global climatic warming - northern peatlands. *Ambio* 25:179-184.
- Lappalainen, E. (1996) Global Peat Resources. Saarijärvi, Finland, Saarijärven Offset Oy.
- Lappalainen, E. and Zurek, S. (1996a). Peatlands in other African countries. In: Lappalainen (Ed.), *Global Peat Resources*, International Peat Society, Finland, pp 239-242.
- Lappalainen, E. and Zurek, S. (1996b). Peatlands in other Asian countries. In: Lappalainen (Ed.), *Global Peat Resources*, International Peat Society, Finland, pp 209-212.
- Lappalainen, E. and Zurek, S. (1996c). Peatlands in central and south America. In: Lappalainen (Ed.), *Global Peat Resources*, International Peat Society, Finland, pp 279-282.
- LUSTRA (2002): Land Use Strategies for Reducing Net Greenhouse Gas Emissions. Annual Report 2002, Uppsala, Sweden. <http://www.sml.slu.se/lustra/index.phtml>
- Martikainen, P. J., H. Nykanen, J. Alm, and J. Silvola. (1995). Change in fluxes of carbon dioxide, methane and nitrous oxide due to forest drainage of mire sites of different trophic. *Plant & Soil* 169:571-577.
- Minkinen, K., R. Korhonen, I. Savolainen, and J. Laine. (2002). Carbon balance and radiative forcing of Finnish peatlands 1900-2100 the impact of forestry drainage. *Global Change Biology* 8:785-799.
- Mosier A. and C. Kroeze, (1999). Contribution of agroecosystems to the global atmospheric N₂O budget. Proceedings of International workshop on reducing N₂O emission from agroecosystems, Banff, Canada, March 1999.

- OECD/IUCN. (1996). Guidelines for aid agencies for improved conservation and sustainable use of tropical and sub-tropical wetlands. OECD, Paris.
- Regina, K., H. Nykänen, J. Silvola, and P. J. Martikainen. (1996). Nitrous oxide production in boreal peatlands of different hydrology and nutrient status. Pages 158-166 in Northern peatlands in global climatic change. Proceedings of the international workshop. Academy of Finland, Hyytiälä
- Rubec, C. (1996). The status of peatland resources in Canada. In: Lappalainen (Ed.), Global Peat Resources, International Peat Society, Finland, pp 243-252.
- Smith K.A., L. Bouwman, B. Braatz, (1999). Nitrous oxide : direct emissions from agricultural soils. Background paper for IPCC Workshop on Good Practice in Inventory Preparation : Agricultural sources of methane and nitrous oxide, 24-26 february 1999, Wageningen, The Netherlands.
- Sundh, I., Nilsson, M., Mikkela, C., Granberg, G., Svensson, B.H.,(2000). Fluxes of methane and carbon dioxide on peat-mining areas in Sweden. *Ambio* 29(8), 499-503.
- Tarnocai, C., Kettles, I.M., Lacelle, B. (2000). Peatlands of Canada. Geological Survey of Canada, Ottawa, Ont. Open File 3152 (map).
- Umeda, Y. And Inoue, T. (1996). Peatlands of Japan. In: Lappalainen (Ed.), Global Peat Resources, International Peat Society, Finland, pp 179-182.
- Xuehui, M and Yan, H. (1996). Peat and peatlands in China. . In: Lappalainen (Ed.), Global Peat Resources, International Peat Society, Finland, pp 163-168
- FLOODED LAND REMAINING FLOODED LAND*
- Dos Santos, M.A., (2000), Inventário emissões de gases de efeito estufa derivadas de Hidrelétricas, PhD. Dissertation, University of Rio de Janeiro, Rio de Janeiro, Brazil, 154p.
- Duchemin, É., (2000), Hydroelectricity and greenhouse gases: Emission evaluation and identification of biogeochemical processes responsible for their production, PhD. Dissertaton, Université du Québec à Montréal, Montréal (Québec), Canada, 321 p (available on CD-ROM)
- Duchemin, É. (2002a), Greenhouse gases emissions from US reservoirs: Spot sampling in the Columbia River Basin and in the Sierra Nevada region, Report for Environmental Fund Defense, DREXenvironment, 47p. (available on CD-ROM)
- Duchemin, É., (2002b), Canadian Reservoir Database, Environment Canada/DREXenvironment, CD-ROM.
- Duchemin, É., M. Lucotte, R. Canuel and A. Chamberland, (1995), Production of the greenhouse gases CH₄ and CO₂ by hydroelectric reservoirs of the boreal region, *Global Biogeochemical Cycles*, 9, 4, 529-540.
- Duchemin, É., R. Canuel, P. Ferland, and M. Lucotte, (1999), Étude sur la production et l'émission de gaz à effet de serre par les réservoirs hydroélectriques d'Hydro-Québec et des lacs naturels (Volet 2), Scientific report, Direction principal Planification Stratégique - Hydro-Québec, 21046-99027c, 48p.
- Duchemin, É., M. Lucotte, R. Canuel, D. Almeida Cruz, H. C. Pereira, J. Dezincourt and A. G. Queiroz, (2000), Comparison of Greenhouse Gas Emissions from an Old Tropical Reservoir and from other Reservoirs Worldwide, *Verh. International Verein. Limnol.*, 27, 3, 1391-1395.
- Duchemin, É., M. Lucotte, R. Canuel, (2002a), CH₄ and CO₂ emissions from boreal reservoirs upon ice break-up, submitted to *Global Biogeochemical Cycles*.
- Duchemin, É, M. Lucotte, V. St-Louis, and R. Canuel, (2002b), Hydroelectric reservoirs as anthropogenic source of greenhouse gases, *World Resources Review*, 27, 3,
- Fearnside, P.M. (2002). Greenhouse gas emissions from a hydroelectric reservoir (Brazil's Tucuruí Dam) and the energy policy implications. *Water, Air and Soil Pollution* 133(1-4): 69-96.
- Galy-Lacaux, C., R. Delmas, C. Jambert, J.-F. Dumestre, L. Labroue, S. Richard and P. Gosse, (1997), Gaseous emissions and oxygen consumption in hydroelectric dams: a case study in French Guyana, *Global Biogeochemical Cycle*, 11,4, 471-483.
- Hélie, (2003), Approche isotopique des flux et de la géochimie du carbone dans les milieux aquatiques de l'est du Canada : exemple du Saint-Laurent et du réservoir Robert-Bourassa, PhD. Dissertation, Université du Québec à Montréal, Montréal (Québec), Canada.
- Houel, (2002), Dynamique de la matière organique terrigène dans les réservoirs boréaux, PhD. Dissertation, Université du Québec à Montréal, Montréal (Québec), Canada, 111p.

- Huttunen, J.T., T.S. Vaisanen, S. K. Hellsten, M. Heikkinen, H. Nykanen, H. Jungner, A. Niskanen, M. O. Virtanen, O.V. Lindqvist, O. S. Nenonen, and P.J. Martikainen, (2002), Fluxes of CH₄, CO₂, and N₂O in hydroelectric reservoir Lokka and Porttipahata in the northern boreal zone in Finland, *Global Biogeochemical Cycles*, 16,1.
- International Commission on Large Dams (ICOLD). 1998. World register of Dams (1998). Paris. International Committee on large Dams (Ed.). Metadatabase..
- Junk, W.J., J.A.S.N. Mello, (1990), Impactos ecologico das represas hidreletricas na bacia amazonica brasileira, *Estudo Avançado*, 4 (8), 126-143
- Keller, M. and R.F. Stallard. (1994). Methane emission by bubbling from Gatun lake, Panama, *J. Geophys. Res.*, 99, D4, 8307-8319.
- Malik, L.K., Koronkevich, N.I., Zaitseva, I.S., Barabanova, E.A. (2000). *Development of Dams in the Russian Federation and NIS Countries*, A WCD briefing paper prepared as an input to the World Commission on Dams, Cape Town, <http://www.dams.org>
- Pinguelli Rosa, L., B. Matvienko Sikar, M.A. dos Santos, E. Matvienko Sikar, (2002), Emissoes de dióxido de carbono e de metano pelos reservatorios hidroelectricos brasileiros, Relatorio de referencia – Inventorio brasileiro de emissoes antropicas de gase de efeito de estufa, Ministerio da Ciencia e tecnologia, Brazil, 199p.
- Schlellhase, H.U. (1994). B.C. Hydro Strategic R&D; Carbon project - Reservoir case study, Powertech Labs inc., Final Report, 1-57.
- Scott, K.J., C.A. Kelly, J.W.M. Rudd, (1999), The importance of floating peat to methane fluxes from flooded peatlands, *Biogeochemistry*, 47, 187-202.
- Smith, L.K., and W.M. Lewis, (1992), Seasonality of methane emissions from five lakes and associated wetlands of the Colorado Rockies, *Global Biogeochemical Cycles*, 6, 4, 323-338
- St-Louis, V., C. A. Kelly, É. Duchemin, J. W. M. Rudd and D.M. Rosenberg. (2000). Reservoir surfaces as sources of greenhouse gases: a global estimate, *Bioscience*, 50,9, 766-775.
- Tavares de Lima I. (2002). Emissao de metano em reservatorio hidreletricos amazonicos atraves de leis de potencia (Methane emission from Amazonian hydroelectric reservoirs through power laws), PhD Dissertation, Universidade de Sao Paulo, Sao Paulo, Brazil, 119 p.
- US Army Corps. (1996). United States Army Corps of Engineers' national Inventory of Dams. Metadatabase.US Army Corps (Ed.). USA.
- WCD, (2000), Dams and Development a new framework for Decision-Making, The report of the World Commission on Dams, Earthscan Publications Ltd, London and Sterling, VA, 356 p.
- WCD, (2001), Dams and Development a new framework for Decision-Making, The report of the World Commission on Dams, Earthscan Publications Ltd, London and Sterling, VA, 356 p.
- Xue, Y., D.A. Kovacic, M.B. David, L.E. Gentry, R.L. Mulvaney and C.W. Lindau (1999). In situ measurements of denitrification in constructed wetlands. *J. Environ. Qual.* 28, 263-269.
- Xuehui, M, and Yan, H. 1996. Peat and peatlands in China. . In: Lappalainen (Ed.), *Global Peat Resources*, International Peat Society, Finland, pp 163-168.

APPENDIX 3A.4 SETTLEMENTS: BASIS FOR FUTURE METHODOLOGICAL DEVELOPMENT

- Brack, C.L. (2002). Pollution mitigation and carbon sequestration by a urban forest. *Environmental Pollution* 116(Suppl. 1): S195-S200.
- Nowak, D.J.; Crane, D.E. (2002). Carbon storage and sequestration by urban trees in the United States. *Environmental Pollution* 116(3): 381-389.
- Nowak, D.J.; Rowntree, R.A.; McPherson, E.G.; Sisinni, S.M.; Kerkmann, E.R. and Stevens, J.C. (1996). Measuring and analyzing urban tree cover. *Landscape and Urban Planning* 36: 49-57.