

References

METHODS FOR ESTIMATION, MEASUREMENT, MONITORING AND REPORTING OF LULUCF ACTIVITIES UNDER ARTICLES 3.3 AND 3.4

- Coleman K. and Jenkinson D.S. (1996). RothC-26.3- A Model for the turnover of carbon in soil. In: Powlson D.S., Smith P., and Smith J.U. (eds.) Evaluation of Soil Organic Matter Models Using Existing, Long-Term Datasets, NATO ASI Series I, Vol.38, Springer-Verlag, Heidelberg, pp. 237-246. 34
- Flanagan L.B., Wever L.A., and Carlson P.J. (2002). Seasonal and interannual variation in carbon dioxide exchange and carbon balance in a northern temperate grassland. *Global Change Biology*, 8: pp. 599-615.
- Follett R.F., Kimble R.F., and Lal R. (2000). The potential of U.S. grazing lands to sequester carbon. In: Follett R.F., Kimble J.M. and Lal R. (eds.) The potential of U.S. grazing lands to sequester carbon and mitigate the greenhouse effect. Lewis Publishers, Boca Raton: pp. 401-430.
- Griffis T.J., Rouse W.R., and Waddington J.M. (2000). Interannual variability of net ecosystem CO₂ exchange at a subarctic fen. *Global Biogeochemical Cycles*, 14: pp. 1109-1121.
- Guo, L.B. and Gifford R.M. (2002). Soil carbon stocks and land use change: a meta analysis. *Global Change Biology*, 8: pp. 345-360.
- Intergovernmental Panel on Climate Change (IPCC). (1997). Houghton J.T., Meira Filho L.G., Lim B., Treanton K., Mamaty I., Bonduki Y., Griggs D.J. and Callander B.A. (Eds). *Revised 1996 IPCC Guidelines for National Greenhouse Inventories*. IPCC/OECD/IEA, Paris, France.
- Intergovernmental Panel on Climate Change (IPCC). (2000). Penman J., Kruger D., Galbally I., Hiraishi T., Nyenzi B., Emmanuel S., Buendia L., Hoppaus R., Martinsen T., Meijer J., Miwa K., and Tanabe K. (Eds). *Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories*. IPCC/OECD/IEA/IGES, Hayama, Japan.
- Intergovernmental Panel on Climate Change (IPCC). (2000b). Watson R., Noble I.R., Bolin B., Ravindranath, N.H., Verardo D.J. and Dokken D.J. (Eds) *Land use, Land-use Change, and Forestry: A Special Report*. Cambridge University Press. Cambridge, UK.
- Kurz W.A. and Apps M.J. (1999). A 70-year retrospective analysis of carbon fluxes in the Canadian forest sector. *Ecological Applications*, 9(2): pp. 526-547.
- Kurz W.A., Apps M.J., Webb T.M., and McNamee P.J. (1992). The carbon budget of the Canadian forest sector: phase I. Forestry Canada, Northwest Region. Information Report NOR-X-326, 93 pp.
- Lal R., Kimble J.M., Follet R.F., Cole C.V. (1998). The potential of U.S. cropland to sequester carbon and mitigate the greenhouse effect. Ann Arbor Press, Chelsea, MI. 128pp.
- Linn D.M., Doran J.W. (1984). Effect of water-filled pore space on carbon dioxide and nitrous oxide production in tilled and non-tilled soils. *Soil Science Society of America Journal*, 48: pp. 1267-1272.
- MacKenzie A.F., Fan M.X. and Cadrin F. (1998). Nitrous oxide emission in three years as affected by tillage, corn-soybean-alfalfa rotations, and nitrogen fertilization. *Journal of Environmental Quality* 27: pp. 698-703.
- Parton W.J., Schimel D.S., Cole C.V. and Ojima D.S. (1987). Analysis of factors controlling soil organic matter levels in Great Plains grasslands. *Soil Science Society of America Journal* 51, 1173-1179.
- Paustian K., Andr n O., Janzen H.H., Lal R., Smith P., Tian G., Tiessen H., van Noordwijk M., and Woomer P.L. (1997). Agricultural soils as a sink to mitigate CO₂ emissions. *Soil Use and Management*, 13: pp. 229-244.
- Robertson G.P., Paul E.A., Harwood R.R. (2000). Greenhouse gases in intensive agriculture: Contributions of individual gases to the radiative forcing of the atmosphere. *Science*, 289 (5486): pp. 1922-1925.
- Smith P., Goulding K.W., Smith K.A., Powlson D.S., Smith J.U., Falloon P.D., Coleman K. (2001). Enhancing the carbon sink in European agricultural soils: Including trace gas fluxes in estimates of carbon mitigation potential. *Nutrient Cycling in Agroecosystems*, 60: pp. 237-252.
- Smith P., Powlson D.S., Glendining M.J., and Smith J.U. (1997). Potential for carbon sequestration in European soils: preliminary estimates for five scenarios using results from long-term experiments. *Global Change Biology*, 3: pp. 67-79.
- Smith P., Powlson D.S., Glendining M.J., Smith J.U. (1998). Preliminary estimates of the potential for carbon mitigation in European soils through no-till farming. *Global Change Biology*, 4: pp. 679-685.

- Smith P., Powlson D.S., Smith J.U., Falloon P.D. and Coleman K. (2000). Meeting Europe's Climate Change Commitments: Quantitative Estimates of the Potential for Carbon Mitigation by agriculture. *Global Change Biology*, 6: pp. 525-539.
- Tian H., Melillo J.M., Kicklighter D.W., McGuire A.D., Helfrich J.V.K. III, Moore B.I., and Vorosmarty C.J. (1998). Effect of interannual climate variability on carbon storage in Amazonian ecosystems. *Nature*, 396: pp. 664-667.
- Tate K.R., Scott N.A., Saggart S., Giltrap D.J., Baisden W.T., Newsome P.F., Trotter C.M., Wilde R.H. (2003). Land-use change alters New Zealand's terrestrial carbon budget: uncertainties associated with estimates of soil carbon change between 1990-2000. *Tellus*, 55B: pp. 364-377.
- Vinten A.J.A., Ball B.C., O'Sullivan M.F., and Henshall J.K. (2002). The effects of cultivation method, fertilizer input and previous sward type on organic C and N storage and gaseous losses under spring and winter barley following long-term leys. *J. Agric. Sci. Camb.*, 139 (3), pp. 231-243.
- Weier K.L., McEwan C.W., Vallis I., Catchpoole V.R., and Myers R.J. (1996). Potential for biological denitrification of fertilizer nitrogen in sugarcane soils. *Aust. J. Agric. Res.*, 47: pp. 67-79.

LULUCF PROJECTS

- Araújo T.M., Higuchi N., and de Carvalho Junior J.A. (1999). Comparison of formulae for biomass content determination in a tropical rain forest site in the state of Pará, Brazil. *For. Ecol. Manage.*, 117: pp. 43-52.
- Avery T.E. and Burkhart H.E. (eds.). (1983). *Forest Measurements*. 3rd edition. McGraw-Hill, New York.
- Beets P.N., Robertson K.A., Ford-Robertson J.B., Gordon J., and Maclaren J.P. (1999). Description and validation of C change: a model for simulating carbon content in managed *Pinus radiata* stands. *New Zealand Journal of Forestry Science* 29(3): pp. 409-427.
- Bennie A.T.P., Taylor H.M., and Georgen P.G. (1987). An assessment of the core-break method for estimating root density of different crops in the field. *Soil Till. Res.* 9: pp. 347-353.
- Blake, G.R. and Hartage K.H. (1986). Bulk density. In Klute A. (ed.) *Methods of Soil Analysis*. Part 1. Physical and Mineralogical Methods. *Agronomy No. 9*. ASA, Madison, WI. pp. 363-375.
- Bland W.L. (1989). Estimating root length density by the core-break method. *Soil Sci. Soc. Am. J.* 53: pp. 1595-1597.
- Böhm W. (1979). *Methods of Studying Root Systems*. Springer-Verlag, New York.
- Boscolo M., Powell M., Delaney M., Brown S., and Faris R. (2000). The cost of inventorying and monitoring carbon. Lessons from the Noel Kempff Climate Action Project. *Journal of Forestry*, September, pp. 24-27 and 29-31.
- Brown J.K. and Roussopoulos J.K. (1974). Eliminating biases in the planar intercept method for estimating volumes of small fuels. *Forest Science* 20: pp. 350-356.
- Brown S. (1997). Estimating biomass and biomass change of tropical forests. A primer. FAO Forestry Paper No.134. Rome, Italy. 55 p.
- Brown S. (2002). Measuring, monitoring, and verification of carbon benefits for forest-based projects. *Phil. Trans. R. Soc. Lond. A* 360: pp. 1669-1684.
- Brown S., Burnham M., Delaney M., Vaca R., Powell M., and Moreno A. (2000a). Issues and challenges for forest-based carbon-offset projects: a case study of the Noel Kempff Climate Action Project in Bolivia. *Mitigation and Adaptation Strategies for Climate Change* 5: pp. 99-121.
- Brown S., Delaney M., and Shoch D. (2001). Carbon monitoring, analysis, and status report for the Rio Bravo Carbon Sequestration Pilot Project. Report to the Programme for Belize, Winrock International, Arlington, VA, USA.
- Brown S., Masera O., and Sathaye J. (2000b). Project-based activities. In: Intergovernmental Panel on Climate Change (IPCC). (2000b). Watson R., Noble I.R., Bolin B., Ravindranath, N.H., Verardo D.J. and Dokken D.J. (Eds) *Land use, Land-use Change, and Forestry: A Special Report*. Cambridge University Press. Cambridge, UK. Chapter 5, pp.283-338.
- Brown S. and Schroeder P. (1999). Spatial patterns of aboveground production and mortality of woody biomass for eastern US forests. *Ecological Applications*, 9: pp. 968-980.

- Cahoon G. A. and Morton E.S. (1961). An apparatus for the quantitative separation of plant roots from soil. *Am. Soc. Hort. Sci.* 78: pp. 593-596.
- Cairns M.A., Brown S., Helmer E.H., and Baumgardner G.A. (1997). Root biomass allocation in the world's upland forests. *Oecologia*, 111: pp. 1-11.
- Clark D. A., Brown S., Kicklighter D.W., Chambers J.Q., Thomlinson J.R., and Jian Ni, (2000). Measuring net primary production in forests: concepts and field methods. *Ecological Applications*, 11: pp. 356-370.
- Dawkins H.C. (1957). Some results of stratified random sampling of tropical high forest. *Seventh British Commonwealth Forestry Conference*, 7 (iii): pp. 1-12.
- Delaney M., Brown S., and Powell M. (1999). 1999 Carbon-Offset Report for the Noel Kempff Climate Action Project, Bolivia. Report to The Nature Conservancy, Winrock International, Arlington, VA, USA.
- Delaney M., Brown S., Lugo A.E., Torres-Lezama A., and Bello Quintero N. (1998). The quantity and turnover of dead wood in permanent forest plots in six life zones of Venezuela. *Biotropica*, 30: pp. 2-11.
- Drew M.C. and Saker L.R. (1980). Assessment of a rapid method, using soil cores, for estimating the amount and distribution of crop roots in the field. *Plant Soil*, 55: pp. 297-305.
- Fang J., Chen A., Peng C., Zhao S., and Ci L. (2001). Changes in forest biomass carbon storage in China between 1949 and 1998. *Science*, 292: pp. 2320-2322.
- Fearnside P.M.(1997). Wood density for estimating forest biomass in Brazilian Amazonia. *Forest Ecology and Management*, 90(1): pp. 59-89.
- Freese F.(1962). Elementary forest sampling. USDA Forest Service Handbook 232, US Government Printing Office, Washington, DC.
- Harmon M.E., Brown S., Gower S.T. (1993). Consequences of tree mortality to the global carbon cycle. In Vinson T.S. and Kolchugina T.P. (eds.). Carbon cycling in boreal and subarctic ecosystems, biospheric response and feedbacks to global climate change. Symposium Proceedings, USEPA, Corvallis, OR, pp. 167-176.
- Harmon M. E. and Sexton J. (1996). Guidelines for Measurements of Woody Detritus in Forest Ecosystems. US LTER Publication No. 20. US LTER Network Office, University of Washington, Seattle, WA, USA. Available at <http://www.lternet.edu/documents/Publications/woodydetritus/>
- Intergovernmental Panel on Climate Change (IPCC). (2000b). Watson R., Noble I.R., Bolin B., Ravindranath, N.H., Verardo D.J. and Dokken D.J. (Eds) *Land use, Land-use Change, and Forestry: A Special Report*. Cambridge University Press. Cambridge, UK.
- Körner C. (1994). Biomass fractionation in plants: a reconsideration of definitions based on plant functions. In: Roy J and Garnier E. (eds.). A Whole Plant Perspective on Carbon-Nitrogen Interactions. SPB Academic Publishing, The Hague, pp. 173-185.
- Kraenzel M., Castillo A., Moore T., and Potvin C. (2003). Carbon storage of harvest-age teak (*Tectona grandis*) plantations, Panama. *Forest Ecology and Management*, 173: pp. 213-225.
- Kurz W. A., Beukema S.J. , and Apps M.J. (1996). Estimation of root biomass and dynamics for the carbon budget model of the Canadian forest sector. *Canadian Journal of Forest Research*, 26: pp. 1973-1979.
- Li Z., Kurz W.A., Apps M.J., and Beukema S. (2003). Belowground biomass dynamics in eh carbon budget model of the Canadian forest sector: recent improvements and implications for the estimation of NPP and NEP. *Canadian Journal of Forest Research*, 33: pp. 126-136.
- Lund G.H. (ed.). (1998). IUFRO Guidelines for designing multipurpose resource inventories. IUFRO World Service Volume 8, Vienna, Austria.
- MacDicken K.G. (1997). *A Guide to Monitoring Carbon Storage in Forestry and Agroforestry Projects*. Winrock International, Arlington, VA, USA, 87 pp, available at: http://www.winrock.org/REEP/PDF_Pubs/carbon.pdf; also in Spanish from Fundacion Solar, Guatemala, http://www.winrock.org/REEP/PDF_Pubs/fundacionsolar.pdf
- Masera O.R., Garza-Caligaris J.F., Kanninen M., Karjalainen T., Nabuurs G.J., Pussinen A., de Jong B.J., and Mohren F. (2003). Modeling Carbon Sequestration in Afforestation and Forest Management Projects: The CO2fix V.2 Approach. *Ecological Modelling* 3237, pp. 1-23
- Page-Dumroese D.S., Jurgensen M.F., Brown R.E., and Mroz G.D. (1999). Comparison of methods for determining bulk densities of rocky forest soils. *Soil Science Society of America Journal*, 63: pp. 379-383.

- Paivinen R., Lund G.H., Poso S., and Zawila-Niedzwiecki T. (eds.). (1994). IUFRO international guidelines for forest monitoring. IUFRO World Series Report 5. Vienna, Austria. 102 p.
- Parresol B.R. (1999). Assessing tree and stand biomass: a review with examples and critical comparisons. *Forest Science*, 45(4): pp. 573-593.
- Parton W.J., Schimel D.S., Cole C.V., and Ojima D.S. (1987). Analysis of factors controlling soil organic matter levels in Great Plains grasslands. *Soil Science Society of America Journal* 51: pp. 1173-1179.
- Pérez L.D. and Kanninen M. (2002). Wood specific gravity and aboveground biomass of *Bombacopsis quinata* plantations in Costa Rica. *Forest Ecology and Management* 165 (1-3): pp. 1-9.
- Pérez L.D. and Kanninen M. (2003). Aboveground biomass of *Tectona grandis* plantations in Costa Rica. *Journal of Tropical Forest Science* 15(1): pp. 199-213.
- Pinard M. and Putz F. (1997): Monitoring carbon sequestration benefits associated with a reduced impact logging project in Malaysia. *Mitigation and Adaptation Strategies for Global Change* 2: pp. 203–215.
- Phillips D.L., Brown S.L., Schroeder P.E., and Birdsey R.A. (2000). Toward error analysis of large-scale forest carbon budgets. *Global Ecology and Biogeography*, 9(4): pp. 305-313.
- Post W.M., Izaurre R.C., Mann L.K., and Bliss N. (1999): Monitoring and verifying soil carbon sequestration. In: Rosenberg N., Izaurre R.C., and Malone E.L. (eds.). Carbon Sequestration in Soils. Batelle Press, pp. 41–82.
- Ritson P. and Sochacki S. (2003). Measurement and prediction of biomass and carbon content of *Pinus pinaster* trees in farm forestry plantations, south-western Australia. *Forest Ecology and Management* 175: pp. 103-117.
- Sampson, R.N. and Scholes R.J. (2000). Additional human-induced activities—Article 3.4. In: Intergovernmental Panel on Climate Change (IPCC). (2000b). Watson R., Noble I.R., Bolin B., Ravindranath, N.H., Verardo D.J. and Dokken D.J. (Eds) *Land use, Land-use Change, and Forestry: A Special Report*. Cambridge University Press. Cambridge, UK. Chapter 4, pp. 181-281.
- Schlegel B., Gayoso J., and Guerra J. (2001). Manual de procedimiento para inventarios de carbono en ecosistemas forestales. Medición de la capacidad de captura de carbono en bosques de Chile y promoción en el mercado mundial. Universidad Austral de Chile. 14 pp.
- Schroeder P., Brown S., Mo J., Birdsey R., and Cieszewski C. (1997). Biomass estimation for temperate broadleaf forests of the United States using inventory data. *Forest Science* 43 (3):pp. 424-434.
- Schroth G., D'Angelo S.A., Teixeira W.G., Haag D., and Lieberei R. (2002). Conversion of secondary forest to agroforestry and monoculture plantations in Amazonia: consequences for biomass, litter, and soil carbon stock after 7 years. *Forest Ecology and Management*, 163: pp. 131-150.
- Segura M. and Kanninen M. (2002). Inventario para estimar carbono en ecosistemas forestales tropicales. In: Orozco L. and Brumér C. (eds). Inventarios forestales para bosques latifoliados en América Central. CATIE - Centro Agronómico Tropical de Investigación y Enseñanza, pp. 202-216.
- Senelwa, K and Sims R.E.H. (1998). Tree biomass equations for short rotation eucalypts grown in New Zealand. *Biomass and Energy* 13(3): pp. 133-140.
- Shepherd D. and Montagnini F. (2001). Carbon Sequestration Potential in Mixed and Pure Tree Plantations in the Humid Tropics. *Journal of Tropical Forest Science* 13(3): pp. 450-459.
- Smucker A.J.M., McBurney S.L., and Srivastava A.K. (1982). Quantitative separation of roots from compacted soil profiles by the hydropneumatic elutriation system Root and soil separation, root response to adverse soil environment. *Agron. J.*, 74: pp. 499-503.
- Sokal R.R. and Rohlf F.J. (1995). Biometry: the principles and practice of statistics in biological research. 3rd Edition. W. H. Freeman and Co., New York.
- Zar J.H. (1996). Biostatistical analysis. Prentice Hall, Englewood Cliffs, New Jersey.