

Use of Inverse Modeling to Evaluate Emissions Inventories

Stephen M. Ogle

Natural Resources Ecology Laboratory
Colorado State University




Acknowledgements: Andrew Schuh, Dan Cooley, F. Jay Breidt and Scott Denning (Colorado State University), Ken Davis, Thomas Lauvaux, Andrew Richardson and Natasha Miles (Pennsylvania State University), Tristram West (DOE), Arlyn Andrews and Pieter Tans (NOAA), Linda Heath (US-Forest Service), Kevin Gurney (Purdue University)




US Mid-Continent Intensive Study


- Compare and reconcile CO₂ fluxes from inventories and atmospheric inversions




MCI Study Region



MCI Pre-Campaign Observations



MCI Campaign Region



Inventory

- **Emissions = Activity Data * Emission Factor**
 - Emission factor may be based measurements and/or modeling
- **Strength**
 - Ground-based activity data used to estimate sources and sinks of CO₂
 - Can be used to determine anthropogenic impact on greenhouse gas emissions
- **Weakness**
 - Does not provide the total CO₂ flux
 - As implemented in national GHG inventories, i.e., only anthropogenic sources
 - Estimates may not be consistent with total CO₂ flux in a region

CO₂ Sources

Included in National GHG Inventory


- Fossil emission of CO₂
- Biomass, dead biomass and soil C stock change
 - Forestland, Cropland, Settlements and Grassland
- Harvested woody product C
- Inorganic CO₂ (including liming)

CO₂ Sources


Typically Not Included in National GHG Inventory

- Harvested Crop Grain C
 - Lateral transport of carbon fixed during photosynthesis which is harvested and a portion exported from the region
- Livestock and Human Respiration of CO₂
 - Mass balance of CO₂ flux with uptake in crops, including lateral transport into region
- Biofuel CO₂ Emission from Combustion
 - Mass balance of CO₂ flux with uptake during photosynthesis in biofuel feedstock crop production
- Wetland CO₂ Emissions
 - Generally considered non-anthropogenic
- Landfill Emissions of CO₂
 - Biogenic and therefore not estimated

Comparing Sources (Gigagrams CO₂)



Inventory Results (Gigagrams CO₂)




West et al., JGR, in review

Inversion


- Can be used to estimate CO₂ flux based on prior information about flux, atmospheric CO₂ concentration data, and transport modeling
- Strength
 - Atmospheric CO₂ measurement constraint
 - Consistent with total flux for region (mass balance)
- Weakness
 - Does not typically disaggregate to source
 - No direct inference on the anthropogenic effect of emissions
 - Cannot capture point sources at current measurement density
 - e.g., power plant emissions are excluded from current inversion analyses

Basis for Inverse Modeling of CO₂




Transport of Air Parcels

June 15, 2004



July 1, 2004




Black = Air Parcels within Domain

Red = Air Parcels outside of Domain (Boundary Conditions)

Transport Model: Conceptualizing

"Observed" at 2PM on 7/7/2004: 368 ppm

"Calculated" for 2PM on 7/7/2004: $370 \text{ ppm} - 3 \text{ ppm} + 1 \text{ ppm} - 2 \text{ ppm} = 366 \text{ ppm}$




Carbon drawdown in upwind areas must be too strong since the observed CO₂ at the tower is higher than what we predict

WLEF Tower


Atmospheric CO₂ Inversion Results

June 15 - July 15, without Fossil Fuel Emissions (g C m⁻²)

Inversion Prior




Inversion Result



Schuh et al., *Biogeosciences Disc.* 6:10195-10241

Global Inversion vs. Inventory (Gigagrams CO₂) *Without CO₂ Tower Observations in the MCI*


2007 CarbonTracker Inversion



Carbon Sink: 1166 Tg CO₂

Excludes Fossil Fuel CO₂ Emissions

2007 Inventory




Carbon Sink: 477 Tg CO₂

Difference between inversion and inventory = 689 Tg CO₂

CarbonTracker: Peters et al.,
PNAS 104:18925-18930

Meso-Scale Inversion vs. Inventory (Gg CO₂) With Additional CO₂ Tower Observations in the MCI


2007 PSU Meso-Scale Inversion



Carbon Sink: 325 Tg CO₂

Excludes Fossil Fuel CO₂ Emissions

2007 Inventory





Carbon Sink: 406 Tg CO₂

Difference between inversion and inventory = 81 Tg CO₂


PSU Meso-Scale: Thomas et al., Preliminary Results

Mathematically Reconciling



Kalman Smoother

Example with simulated data



Combining inventory and inversion to obtain “reconciled” estimates and uncertainty (sources)

Research & Development

■ Inventory methods



- Requires estimates for sources not typically reported in national GHG inventory
 - Including non-anthropogenic impacts on sources
- Challenge to obtain all of the necessary activity data and emission “factor” estimates

■ Inversion methods

- Further development of sub-continental inversions frameworks
 - Observational data density, boundary conditions and transport methods
 - New satellite missions gathering CO₂ concentration data could be used to inform inversions and increase observation data density
- Uncertainties are not well quantified
 - Prior, boundary conditions and transport

What if uncertainties are not rigorous?

Combining data from inversion and inventory assumes that the underlying uncertainty is representative and contains the true estimate of CO₂ flux at a specified confidence level.



Concluding Remarks

- Atmospheric CO₂ inversions are a promising technology for verification of inventories
 - Constrained by mass balance of CO₂ concentration changes in the atmosphere across space and time
 - Requires more research and development, and an expanded network of atmospheric CO₂ observations before an inversion system could be operational
- Inventories will need to be further developed in order to compare with inversions
 - Requires additional sources not typically included a national inventory for reporting under IPCC GL

Thanks for your attention!



National Aeronautics
and Space Administration

