

CHAPTER 5

CROPLAND

Contents

5 CROPLAND

5.1 INTRODUCTION 4

5.2	CROPLAND REMAINING CROPLAND	4
5.2.1	Biomass	4
5.2.1.1	Choice of methods.....	4
5.2.1.2	Choice of emission factors	5
5.2.1.3	Choice of activity data.....	14
5.2.1.4	Calculation steps for tier 1 and tier 2.....	15
5.2.1.5	uncertainty assessment	15
5.2.2	Dead organic matter	15
5.2.3	Soil carbon	15
5.2.3.1	Choice of method	16
5.2.3.2	Choice of stock change and emission factors.....	17
5.2.3.3	Choice of activity data.....	21
5.2.3.4	Calculation steps for Tier 1	26
5.2.3.5	Uncertainty assessment	27
5.2.4	Non-CO ₂ greenhouse gas emissions from biomass burning.....	27
5.3	LAND CONVERTED TO CROPLAND	28
5.3.1	Biomass	28
5.3.1.1	Choice of method	28
5.3.1.2	Choice of emission factors	30
5.3.1.3	Choice of activity data.....	32
5.3.1.4	Calculation steps for tier 1 and tier 2.....	33
5.3.1.5	uncertainty assessment	33
5.3.2	Dead Organic Matter.....	33
5.3.3	Soil carbon	33
5.3.3.1	Choice of method	33
5.3.3.2	Choice of stock change and emission factors.....	34
5.3.3.3	Choice of activity data.....	36
5.3.3.4	Calculation steps for Tier 1	38
5.3.3.5	Uncertainty assessment	39
5.3.4	Non-CO ₂ greenhouse gas emissions from biomass burning.....	39
5.4	COMPLETENESS, TIME SERIES, QA/QC, AND REPORTING	39
5.5	METHANE EMISSIONS FROM RICE CULTIVATION	40
5.5.1	Choice of method	40
5.5.2	Choice of emission and scaling factors	44
5.5.3	Choice of activity data.....	49
5.5.4	Uncertainty assessment	51
ANNEX 5A.1ESTIMATION OF DEFAULT STOCK CHANGE FACTORS FOR MINERAL SOIL C EMISSIONS/REMOVALS FOR CROPLAND	60
ANNEX 5A.2ESTIMATION OF DEFAULT EMISSION FACTORS AND SCALING FACTORS FOR CH ₄ EMISSION FROM RICE CULTIVATION.....	88
ANNEX 5A.3ESTIMATION OF COEFFICIENTS AND FACTORS FOR BIOMASS C EMISSIONS AND REMOVALS IN CROPLAND.....	89

EQUATIONS

Equation 5. 1 Cropland litter carbon input for three-pool steady-state C model.....	24
Equation 5. 2 CH ₄ emissions from rice cultivation	40
Equation 5. 3 Adjusted daily emission factor.....	44
Equation 5. 4 Adjusted CH ₄ emission scaling factors for organic amendments	48

FIGURES

Figure 5. 1 Classification scheme for cropping systems. In order to classify cropland management systems, the inventory compiler should start at the top and proceed through the diagram answering questions (move across branches if answer is yes) until reaching a terminal point on the diagram. The classification diagram is consistent with default stock change factors in Table 5.5. C input classes (i.e., low, medium, high and high with organic amendment) are further subdivided by tillage practice.	23
Figure 5. 2 Decision tree for CH ₄ emissions from rice production	43

TABLES

UPDATED-Table 5. 1 Default coefficients for above-ground woody biomass and harvest cycles in cropping systems containing perennial species.....	6
Updated - Table 5. 2 Potential C storage for agroforestry systems in different ecoregions of the world.....	7
Table 5. 3 Examples of classification of agroforestry system.....	10
Updated- Table 5. 4 Default maximum and time-averaged mean above-ground biomass for various types of perennial croplands (tonnes ha ⁻¹)	11
Updated Table 5. 5 Examples of perennial cropland subcategories which a country may have.....	15
Updated Table 5. 6 Relative stock change factors (FLU, FMG, AND FI) (over 20 years) for different management activities on cropland.....	19
NEW GUIDANCE Table 5. 7 Default values for nitrogen and lignin contents in crops for three-pool steady-state C model.....	21
Table 5. 8 Text to be provided in SOD	29
Table 5. 9 Example of a simple disturbance matrix (Tier 2) for the impacts of land conversion activities on carbon pools	30
Updated Table 5. 10 Default biomass carbon stocks removed due to land conversion to cropland.....	31
UPDATED Table 5. 11 Default biomass carbon stocks present on land converted to cropland in the year following conversion.....	31
Table 5. 12 Soil stock change factors (FLU, FMG, FI) for land-use conversions to cropland	35
UPDATED- Table 5. 13 Default CH ₄ baseline emission factor assuming no flooding for less than 180 days prior to rice cultivation, and continuously flooded during rice cultivation without organic amendments.....	45
UPDATED Table 5. 14 Default CH ₄ emission scaling factors for water regimes during the cultivation period relative to continuously flooded fields.....	47
UPDATED - Table 5. 15 Default CH ₄ emission scaling factors for water regimes before the cultivation period.....	48
Table 5. 16 Default conversion factors for different types of organic amendments	49

BOXES

Box 5. 1 Text to be provided in SOD.....	41
Box 5. 2 Conditions influencing CH ₄ emissions from rice cultivation	42
Box 5. 3 Good practice guidance for developing baseline emission factors (EF) for methane emission from rice cultivation	46
New Guidance Box 5. 4 Example of how to estimate methane emission from rice cultivation using Tier 1 method	49

First Order Draft

5.1 INTRODUCTION

No Refinement

5.2 CROPLAND REMAINING CROPLAND

No Refinement

5.2.1 Biomass

5.2.1.1 CHOICE OF METHODS

This section provides elaboration on methods, clarifying how to use updated factors.

Carbon can be stored in the biomass of croplands that contain perennial woody vegetation including, but not limited to, monocultures such as coffee, oil palm, coconut, rubber plantations, fruit and nut orchards, and polycultures such as agroforestry systems. The default methodology for estimating carbon stock changes in woody biomass is provided in Chapter 2, Section 2.2.1. This section elaborates this methodology with respect to estimating changes in carbon stocks in biomass in *Cropland Remaining Cropland*.

The change in biomass is only estimated for perennial woody crops. For annual crops, increase in biomass stocks in a single year is assumed equal to biomass losses from harvest and mortality in that same year - thus there is no net accumulation of biomass carbon stocks.

Changes in carbon in cropland biomass (ΔC_{CCB}) may be estimated from either: (a) annual rates of biomass gain and loss (Chapter 2, Equation 2.7) or (b) carbon stocks at two points in time (Chapter 2, Equation 2.8). The first approach (gain-loss method) provides the default Tier 1 method and can also be used at Tier 2 or 3 with refinements described below. The second approach (the stock-difference method) applies either at Tier 2 or Tier 3, but not Tier 1. It is *good practice* to improve inventories by using the highest feasible tier given national circumstances. It is *good practice* for countries to use a Tier 2 or Tier 3 method if carbon emissions and removals in *Cropland Remaining Cropland* is a key category and if the sub-category of biomass is considered significant. It is *good practice* for countries to use the decision tree in Figure 2.2 in Chapter 2 to identify the appropriate tier to estimate changes in carbon stocks in biomass.

Tier 1

The default method is to multiply the area of perennial woody cropland by a net estimate of biomass accumulation from growth and subtract losses associated with harvest or gathering or disturbance (according to Equation 2.7 in Chapter 2). Losses are estimated by multiplying a carbon stock value by the area of cropland on which perennial woody crops are harvested.

Default Tier 1 assumptions are: all carbon in perennial woody biomass removed (e.g., biomass cleared and replanted with a different crop) is emitted in the year of removal; and perennial woody crops accumulate carbon for an amount of time equal to a nominal harvest/maturity cycle. The latter assumption implies that perennial woody crops accumulate biomass for a finite period until they are removed through harvest or reach a steady state where there is no net accumulation of carbon in biomass because growth rates have slowed and incremental gains from growth are offset by losses from natural mortality, pruning or other losses.

Under Tier 1, updated default factors shown in updated Table 5.1 to Table 5.4, are applied to nationally derived estimates of land areas. For perennial cropland C uptake, multiply unharvested area that is still younger than the age of maturity by the above-ground growth rate. If harvest and immature areas are unknown, it is assumed C uptake in growth is balanced by emissions due to crop turnover in cropland remaining cropland. For perennial cropland C losses, it should be noted that updated tables provide two types of carbon stocks of perennial woody biomass per area. One is maximum carbon stock at harvest/maturity state and the other is mean carbon stock for whole lifetime of perennial woody biomass. These values should be used appropriately to calculate carbon losses following the guidance in 5.2.1.2.

Tier 2

Two methods can be used for Tier 2 estimation of changes in biomass. Method 1 (also called the **Gain-Loss Method**) requires the biomass carbon loss to be subtracted from the biomass carbon increment for the reporting year (Chapter 2, Equation 2.7). Method 2 (also called the **Stock-Difference Method**) requires biomass carbon stock inventories for a given land-use area at two points in time (Chapter 2, Equation 2.8).

A Tier 2 estimate, in contrast, will generally develop estimates for the major woody crop types by climate zones, using country-specific carbon accumulation rates and stock losses where possible or country-specific estimates of carbon stocks at two points in time. Under Tier 2, carbon stock changes are estimated for above-ground and below-ground biomass in perennial woody vegetation. Tier 2 methods involve country-specific or region-specific estimates of biomass stocks by major cropland types and management system, and estimates of stock change as a function of major management system (e.g., dominant crop, productivity management). To the extent possible, it is *good practice* for countries to incorporate changes in perennial crop or tree biomass using country-specific or region-specific data. Where data are missing, default data may be used.

Tier 3

A Tier 3 estimate will use a highly disaggregated Tier 2 approach or a country-specific method involving process modelling and/or detailed measurement. Tier 3 involves inventory systems using statistically-based sampling of carbon stocks over time and/or process models, stratified by climate, cropland type and management regime. For example, validated species-specific growth models that incorporate management effects such as harvesting and fertilization, with corresponding data on management activities, can be used to estimate net changes in cropland biomass carbon stocks over time. Models, perhaps accompanied by measurements like those in forest inventories, can be used to estimate stock changes and extrapolate to entire cropland areas, as in Tier 2.

Key criteria in selecting appropriate models are that they are capable of representing all of the management practices that are represented in the activity data. It is critical that the model be validated with independent observations from country-specific or region-specific field locations that are representative of climate, soil and cropland management systems in the country.

5.2.1.2 CHOICE OF EMISSION FACTORS

This section has updated factors and an elaboration on the methods.

Emission and removal factors required to estimate the changes in carbon stocks include (a) annual biomass accumulation or growth rate, and (b) biomass loss factors which are influenced by such activities as removal (harvesting), fuelwood gathering and disturbance.

Above-ground woody biomass growth rate

Tier 1

Updated Tables 5.1 to 5.4 provide estimates of biomass stocks and biomass growth rates and losses for major climatic regions and agricultural systems. Updated Table 5.1 provides default values of biomass growth and losses applicable to general perennial cropping system for each climate region. Updated Table 5.2 provides default potential carbon storage for agro-forestry system in tropical and sub-tropical region. Updated Table 5.3 provides default values of biomass growth and losses for various perennial cropping systems for each climate region. Countries should use appropriate default values of above-ground biomass growth rate relative to each climate region and cropping system from updated Table 5.1 or Table 5.4. However, given the large variation in cropping systems, incorporating trees or tree crops, it is *good practice* to seek national data on above-ground woody biomass growth rate.

Tier 2

Annual woody biomass growth rate data can be, at a finer or disaggregated scale, based on national data sources for different cropping and agroforestry systems. Rates of change in annual woody biomass growth rate should be estimated in response to changes in specific management/land-use activities (e.g., fertilization, harvesting, thinning). Results from field research should be compared to estimates of biomass growth from other sources to verify that they are within documented ranges. It is important, in deriving estimates of biomass accumulation rates, to recognize that biomass growth rates will occur primarily during the first 20 years following changes in management, after which time the rates will tend towards a new steady-state level with little or no change occurring unless further changes in management conditions occur.

Tier 3

For Tier 3, highly disaggregated factors for biomass accumulation are needed. These may include categorisation of species, specific for growth models that incorporate management effects such as harvesting and fertilization. Measurement of above-ground biomass, similar to forest inventory with periodic measurement of above-ground biomass accumulation, is necessary.

First Order Draft

UPDATED-TABLE 5.1 DEFAULT COEFFICIENTS FOR ABOVE-GROUND WOODY BIOMASS AND HARVEST CYCLES IN CROPPING SYSTEMS CONTAINING PERENNIAL SPECIES

Domain	Ecological zone	Continent	Cropping-system	Maximum above-ground biomass carbon stock at harvest (L_{max}) (tonnes C ha ⁻¹)	Harvest/Maturity cycle (yr)	Biomass accumulation rate (G) (tonnes C ha ⁻¹ yr ⁻¹)	Mean biomass carbon loss (L_{mean}) (tonnes C ha ⁻¹ yr ⁻¹)	Error range ¹
Tropical ³	Highland	All	General perennial agroforestry	117 ²	30	3.9 ²	63	± 75%
Tropical	Dry	All	General perennial agroforestry	13 ²	5	2.6 ²	9	± 75%
Tropical	Moist	All	General perennial agroforestry	13 ²	8	6.1 ²	21	± 75%
Tropical	Wet	All	General perennial agroforestry	13 ²	5	10.0	25 ²	± 75%
Temperate and subtropical ⁴		All	General perennial cropland	TBD	TBD	TBD	TBD	TBD
<p>TBD – To be determined, Temperate (and subtropical) values will be prepared for the second order draft.</p> <p>Note: Values for tropical zones are derived from the literature survey and synthesis published by Schroeder (1994).</p> <p>¹ Represents a nominal estimate of error, equivalent to two times standard deviation, as a percentage of the mean.</p> <p>² The default factors are modified from the original Table 5.1 in the 2006 IPCC guidelines in order to be consistent with the information provided in Schroeder (1994).</p> <p>³ The climate region are modified from the original Table 5.1 in the 2006 IPCC guidelines in order to be consistent with the information provided in Schroeder (1994).</p> <p>⁴ Values for temperate domain are derived from cropping systems in Table 5.4.</p>								

UPDATED -
TABLE 5.2 POTENTIAL C STORAGE FOR AGROFORESTRY SYSTEMS IN DIFFERENT ECOREGIONS OF THE WORLD

Domain	Ecological zone	Continent	Agroforestry System	Above-ground biomass (tonnes ha ⁻¹)	Range (tonnes ha ⁻¹)	Reference	Note
Tropical	Tropical rainforest	Africa	Multi-strata (Togo: Coffee-Albizia)	140		Dossa et al. (2008)	The climate is a Sudan-Guinean type characterized by a bimodal rainfall regime with a mean annual rainfall of 1,400 mm
		North and South America					
		Asia	Multi-strata (Philippines: Gmelina arborea + coffee)	116	sd= 31	Lasco et al., 2001	
			Improved fallow (Philippines: Leucaena leucocephala)	32	sd= 11.6	Lasco et al. (2010)	Average annual biomass for 6 year cycle
			Multi-strata (SE Asia: Jungle rubber)	304	Error= 17	Tomich et al., 1998	From GPG 2006
			Multi-strata (SE Asia: Jungle rubber)	116	Error= 53	Lasco et al., 2001	From GPG 2006
	Tropical moist deciduous forest	Africa					
		North and South America	Mexico: various AF systems	56.5	35.3 to 74.0	Soto-Pinto et al., (2010)	Low-tropical agro-climatic zone of Chiapas
			Improved fallow (Columbia)	20.9	6.7 to 37	Barrios and Cabo, (2004)	Mean annual rainfall of 1900 mm
			Multi-strata (Guatemala: Coffee agroforests)	255.2 ± 13.2	148 to 518	Schmitt-Harshet et al. (2012)	Annual rainfall and temperature averages 2504 mm and 18–24 °C,
			Multi-strata (Cocoa agroforestry [six countries])	49 ± 35		Somarriba et al. (2013)	
			Agrosilvicultural (S America)	70.5	39 - 102	Albrecht and Kandji, 2003	From GPG 2006
			Silvopastoral (N America)	151.0	104 - 198	Albrecht and Kandji, 2003	From GPG 2006
		Australia	Silvopastoral	39.5	28 - 51	Albrecht and Kandji, 2003	From GPG 2006
		Asia	Agrosilvicultural (SE Asia)	120.0	12 - 228	Albrecht and Kandji, 2003	From GPG 2006
			Silvopastoral (N Asia)	16.5	15 - 18	Albrecht and Kandji, 2003	From GPG 2006

First Order Draft

UPDATED - TABLE 5.2 (CONTINUED) Potential C storage for agroforestry systems in different ecoregions of the world							
Domain	Ecological zone	Continent	Agroforestry System	Above-ground biomass (tonnes ha ⁻¹)	Range (tonnes ha ⁻¹)	Reference	Note
Tropical	Tropical dry forest	Africa					
		North and South America	Agrosilvicultural (S America)	117.0	39 - 195	Albrecht and Kandji, 2003	From GPG 2006
			Silvopastoral (N America)	132.5	90 - 175	Albrecht and Kandji, 2003	From GPG 2006
		Asia	Agrosilvicultural (SE Asia)	75.0	68 - 81	Albrecht and Kandji, 2003	From GPG 2006
	Tropical shrub land	Africa	Multi-strata (Kenya: Homegarden)	15.6	13.8-17.3	Henry et al. (2009)	East African highlands with average altitudes of 1600 and 1200 m.a.s.l., and annual rainfall of 1400-1800 mm (Kenya)
			Plantation (Kenya: Woodlots)	81	13.8-17.3	Henry et al. (2009)	East African highlands with average altitudes of 1600 and 1200 m.a.s.l., and annual rainfall of 1400-1800 mm (Kenya)
			Improved fallow (E Africa: 1 year fallow)	35.0	27 - 44	Albrecht and Kandji, 2003	From GPG 2006
			Improved fallow (E Africa: 2 year fallow)	12.0	7 - 21	Albrecht and Kandji, 2003	From GPG 2006
		North and South America					
		Asia	Improved fallow (SE Asia: 6 year fallow (average))	16.0	4 - 64	Lasco and Suson, 1999	From GPG 2006
			Alley cropping (SE Asia)	2.9	1.5 - 4.5	Lasco et al., 2001	From GPG 2006
	Tropical mountain systems	Africa	Agrosilvicultural	41.0	29 - 53	Albrecht and Kandji, 2003	
		North and South America	Silvopastoral (N America)	143.5	133 - 154	Albrecht and Kandji, 2003	
		Asia	TBD	TBD	TBD	TBD	

UPDATED TABLE 5.2 (CONTINUED) Potential C storage for agroforestry systems in different ecoregions of the world							
Domain	Ecological zone	Continent	Agroforestry System	Above-ground biomass (tonnes ha ⁻¹)	Range (tonnes ha ⁻¹)	Reference	Note
Subtropical	Subtropical humid forest	North and South America	TBD	TBD	TBD	TBD	
		Asia	India: Gmelina arborea + crops	14.1	TBD	Swamy and Puri (2005)	5 years old
			India: Populus deltoides + crops	203	TBD	Singh and Lodhiyal (2009)	8 years old
			India: Populus deltoides + crops	74.5	TBD	Rizvi et al., (2011)	Average of two values
			India: Populus deltoides + crops	40.3	19.3 to 57.7	Yadava (2010)	
	Subtropical dry forest	North and South America	TBD	TBD	TBD	TBD	
		Asia	TBD	TBD	TBD	TBD	
	Subtropical steppe	North and South America	TBD	TBD	TBD	TBD	
		Asia	TBD	TBD	TBD	TBD	
	Subtropical	North and South America	TBD	TBD	TBD	TBD	
		Asia	TBD	TBD	TBD	TBD	
Temperate	TBD	TBD	TBD	TBD	TBD	TBD	
Boreal	TBD	TBD	TBD	TBD	TBD	TBD	
TBD – To be determined for the second order draft based on information on studies including Dossa et al. (2008), Makumba et al. (2007), Takimoto et al. (2008), Henry et al. (2009), Soto-Pinto et al., (2010), Barrios and Cabo, (2004), Schmitt-Harshet et al. (2012), Somarriba et al. (2013), Swamy and Puri (2005), Singh and Lodhiyal (2009), Rizvi et al., (2011), Yadava (2010), Lasco et al. (2010)]							

237

238

239

First Order Draft

TABLE 5.3 EXAMPLES OF CLASSIFICATION OF AGROFORESTRY SYSTEM

Agroforestry system	Description
Multi-strata	This class includes (i) Home garden systems, which refer to intimate combination of multipurpose trees, shrubs, herbaceous plants with annual and perennial crops growing in or adjacent to a home compound. The garden is managed intensively usually by family labor. (ii) Growing shade-tolerant perennials species such as cacao, coffee under or in between over story shade-, timber-, or other commercial tree crops (USAID 2014). For this study, plantation crops were not included in this class.
Tree Intercropping	Includes (i) Alley cropping systems, which are defined as growing food crops between hedgerows of planted shrubs and trees, preferably leguminous species. The hedges are pruned periodically during the crop's growth to provide biomass (which, when returned to the soil, enhances its nutrient status and physical properties) and to prevent shading of the growing crops. (ii) This class also comprises traditional intercropping systems, which consist of growing crops under scattered or in systematically-planted trees (Nair 1993b).
Silvopastoral	This class refers to grazing under scattered or planted trees or tree-fodder systems: fodder banks, parkland.
Protective Systems	This class refers to (i) Protective systems, which refers to boundary planting, windbreaks, shelterbelts, soil conservation hedges trees used to protect fields from wind damage, sea encroachment, floods, etc.
Plantations	This class refers to plantation crops systems such as oil palm, rubber, coconut, cacao, coffee, and tea, cashew, which may include understory crops production in alternate or in other regular arrangements.
Improved fallow	Defined as land resting for cultivation, but comprises planted and managed trees, preferably leguminous, shrubs and herbaceous cover crops before it is cultivated again. The improved fallow agroforestry system can be implemented prior to the establishment of any other class mentioned above.
Vineyard systems	A plantation of grapevines, typically producing grapes used for winemaking
Orchards systems	Land planted with fruit trees (apple, pear, plum and cherry tree). Understory vegetation is usually mowed or grazed.
Source: Description of agroforestry systems adopted from USAID (2014) and Nair (1993b)	

UPDATED- TABLE 5. 4 DEFAULT MAXIMUM AND TIME-AVERAGED MEAN ABOVE-GROUND BIOMASS FOR VARIOUS TYPES OF PERENNIAL CROPLANDS (TONNES HA ⁻¹)											
Domain	Ecological zone	Continent	Cropping system	Maximum above-ground biomass carbon stock at harvest (L _{max}) (tonnes C ha ⁻¹)	Error L _{max}	Harvest /Maturity cycle (yr)	Above-ground biomass accumulation rate (G) (tonnes C ha ⁻¹ yr ⁻¹)	Error G	Mean biomass carbon stock (L _{mean}) (tonnes C ha ⁻¹ yr ⁻¹)	Error L _{mean}	References
Temperate	All	All	General perennial cropland	TBD	TBD	TBD	TBD	TBD	TBD	TBD	
			Orchard e.g. apple	TBD	TBD	TBD	TBD	TBD	TBD	TBD	
			Shrub e.g. berry	TBD	TBD	TBD	TBD	TBD	TBD	TBD	
			Vine e.g. grape	9.1	TBD	30	0.32	TBD	4.5	TBD	Kroodsma & Field 2006; Morande et al 2017; Buwalda and Smith 1987
			Nurseries e.g. Christmas trees	TBD	TBD	TBD	TBD	TBD	TBD	TBD	
			Short Rotation Coppice	12.69	TBD	4	3.2	TBD	6.35	TBD	Hauk et al 2013, adjustment from Krasuska and Rosenqvist 2011
Tropical	Wet, humid	All	Oil palm <i>Elaeis guineensis</i>	94.75	TBD	25	1.53	TBD	24.24	TBD	Germer and Sauerborn 2008 [Agus et al 2013; Ziegler et al 2012; Khasanah et al 2015; Sanquetta et al 2015...]
			Rubber monoculture <i>Hevea brasiliensis</i>	TBD	TBD	30	TBD	TBD	45.1 [Z et al 56; M et al 44.1; A et al 58]	TBD	Blagodatsky et al 2016 [Ziegler et al 2012; Nizami et al 2014; Margiotto et al 2014; Agus et al 2013...]
			Coffee <i>Coffea</i>	TBD	TBD	TBD	TBD	TBD	TBD	TBD	
			Cacao <i>Theobroma cacao</i>	TBD	TBD	TBD	TBD	TBD	TBD	TBD	
Tropical	Wet, humid	All	Coconut <i>Cocos nucifera</i>	196	TBD	TBD	TBD	TBD	TBD	TBD	Lasco et al., 2002

First Order Draft

UPDATED- TABLE 5. 4 DEFAULT MAXIMUM AND TIME-AVERAGED MEAN ABOVE-GROUND BIOMASS FOR VARIOUS TYPES OF PERENNIAL CROPLANDS (TONNES HA ⁻¹)											
Domain	Ecological zone	Continent	Cropping system	Maximum above-ground biomass carbon stock at harvest (L _{max}) (tonnes C ha ⁻¹)	Error L _{max}	Harvest /Maturity cycle (yr)	Above-ground biomass accumulation rate (G) (tonnes C ha ⁻¹ yr ⁻¹)	Error G	Mean biomass carbon stock (L _{mean}) (tonnes C ha ⁻¹ yr ⁻¹)	Error L _{mean}	References
			Banana <i>Musa</i>	TBD	TBD	TBD	TBD	TBD	TBD	TBD	
			Orchard e.g. fruit, nuts	TBD	TBD	TBD	TBD	TBD	TBD	TBD	
	All	All	Cinnamon	TBD	TBD	TBD	TBD	TBD	TBD	TBD	Siregar & Gintings, 2000 Lakprasadi & Navaratne 2012
	Tropical rainforest	SE Asia	Oil Palm	136.0	62 - 202	TBD	TBD	TBD	TBD	TBD	
		SE Asia	Mature rubber	178.0		TBD	TBD	TBD	TBD	TBD	Palm et al., 1999
		SE Asia	Young rubber	48.0	16 - 80	TBD	TBD	TBD	TBD	TBD	Wasrinet al., 2000
		SE Asia	Young cinnamon (7 years)	68.0	TBD	TBD	TBD	TBD	TBD	TBD	Siregar&Gintings, 2000
		SE Asia	Coconut	196.0	TBD	TBD	TBD	TBD	TBD	TBD	Lasco et al., 2002
	Tropical moist deciduous forest	Africa	Cocoa -21 years	130.0	TBD	TBD	TBD	TBD	TBD	TBD	Kongsager et al., 2013
			Oil Palm-7 years	43.4	TBD	TBD	TBD	TBD	TBD	TBD	Kongsager et al., 2013
			Oil Palm-16 years	56.0	TBD	TBD	TBD	TBD	TBD	TBD	Kongsager et al., 2013
			Oil palm-23 years	90.6	TBD	TBD	TBD	TBD	TBD	TBD	Kongsager et al., 2013
			Rubber -12 years	123.0	TBD	TBD	TBD	TBD	TBD	TBD	Kongsager et al., 2013
			Rubber -44 years	427.2	TBD	TBD	TBD	TBD	TBD	TBD	Kongsager et al., 2013
			Orange -25 years	152.6	TBD	TBD	TBD	TBD	TBD	TBD	Kongsager et al., 2013
Temperate, subtropical, tropical	All	All	Tea <i>Camellia sinensis</i>	TBD	TBD	TBD	TBD	TBD	TBD	TBD	

UPDATED- TABLE 5. 4 DEFAULT MAXIMUM AND TIME-AVERAGED MEAN ABOVE-GROUND BIOMASS FOR VARIOUS TYPES OF PERENNIAL CROPLANDS (TONNES HA⁻¹)

Domain	Ecological zone	Continent	Cropping system	Maximum above-ground biomass carbon stock at harvest (L _{max}) (tonnes C ha ⁻¹)	Error L _{max}	Harvest /Maturity cycle (yr)	Above-ground biomass accumulation rate (G) (tonnes C ha ⁻¹ yr ⁻¹)	Error G	Mean biomass carbon stock (L _{mean}) (tonnes C ha ⁻¹ yr ⁻¹)	Error L _{mean}	References
TBD – to be determined for the second order draft based on available information on recent studies including, Bilas Singh et al, 2015, Yashmita-ulman and S. Avudainayagam, 2012, G. Singh, 2017, Miria A. and Anisa B. Khan, 2015, A. S. Dogra et al, 2014, Y. Ulman and S. Avudainayagam, 2014, Swami K. R. et al, 2012, Anil Kumar Yadava, 2010, Sanjeev K. Chauhan et al, 2010, N. Singh and L. S. Lodhiya, 2016, F. Mohsin et al, 2005, K. C. Singh, 2005, Umrao et al, 2010, S. Goswami et al, 2016, C. E Harwood and EKS Nambiar (CSIRO –ACAIR), Palm et al., 1999, Wasrinet al., 2000, Siregar&Gintings, 2000, Lasco et al., 2002, Kongsager et al., 2013, Kotowska et al 2015, Adachi et al 2011, Corley and Tinker, 2003, Bwalya JM 2012, G. Liguori et al., 2009, Ting Wu et al., 2012, Zhaopeng Ou Yang et al. 2012, Kroodsma, D. A. and Field, C. B., 2006. Morgan, K.T. et al., 2006, Chalmers D.J. and Van Den Ende, B. 1975., Jiménez, C.M. and Diaz, J.B.R. 2003, Jiménez, C.M. and Diaz, J.B.R. 2004., Haynes, R.J. and Goh, K.M., 1980., Palmer, J.W., J.N. Wünsche, M. Meland and A. Hann. 2002., Lovatt, C.J. 1996, Villalobos, F.J., Testi, L., Hidalgo, J., Pastor, M., Orgaz, F., 2006., Murphy, T., Jones, G., Vanclay, J., and Glencross, K., 2013., Nendel, C. and Kersebaum, K.C. 2004., Buwalda, J.G. and Smith, G.S. 1987., BMLFUW (Bundesministerium für Land- und Forstwirtschaft, Umwelt und Wasserwirtschaft) 2000: Splechtna, B. & Glatzel, G. 2005: Wirth, C. et al., 2004, Mokany, K., R.J. Raison & A.S.P. Rokushkin, 2006, Popken, S., 2011, Kandler, G.; Bosch, B. 2013, Gyldenkerne, S. et al., 2005, Moxley, J. et al., 2014, Juhos K. és Tökei, L. 2012., Kort, J. and Turnock, R. 1999, Kerckhoffs, L.H.J. and Reid, J.B. 2007, Pessler C, 2012, McConkey, B. et al., 2006, Milne, R. and Brown, T. A. 1997, Walter 2012, Zanotelli et al 2015, Montanaro et al 2017, Scandellari et al 2016, Wu et al 2012 and Japan's studies (on-going work).]											

First Order Draft

Below-ground biomass accumulation**Tier 1**

The default assumption is that there is no change in below-ground biomass of perennial trees in agricultural systems. Default values for below-ground biomass for agricultural systems are not available.

Tier 2

This includes the use of actually measured below-ground biomass data from perennial woody vegetation. Estimating below-ground biomass accumulation is recommended for Tier 2 calculation. Root-to-shoot ratios show wide ranges in values at both individual species (e.g., Anderson *et al.*, 1972) and community scales (e.g., Jackson *et al.*, 1996; Cairns *et al.*, 1997). Limited data is available for below ground biomass thus, as far as possible, empirically-derived root-to-shoot ratios specific to a region or vegetation type should be used.

Tier 3

This includes the use of data from field studies identical to forest inventories and modelling studies, if stock difference method is adopted.

Biomass losses from removal, fuelwood and disturbance**Tier 1**

The default assumption is that all biomass lost is assumed to be emitted in the same year. Biomass removal, fuelwood gathering and disturbance loss data from cropland source are not available. FAO provides total roundwood and fuelwood consumption data, but not separated by source (e.g., Cropland, Forest Land, etc.). It is recognized that statistics on fuelwood are extremely poor and uncertain worldwide. Default removal and fuelwood gathering statistics (discussed in Chapter 4, Section 4.2) may include biomass coming from cropland such as when firewood is harvested from home gardens. Thus, it is necessary to ensure no double counting of losses occurs. If no data are available for roundwood or fuelwood sources from Cropland, the default approach will include losses in Forest Land (Section 4.2) and will exclude losses from Cropland. Updated Tables 5.1 to 5.4 provides default values of maximum carbon stock per area (L_{\max}) and mean carbon stock per area (L_{mean}). Countries should use L_{\max} in updated Table 5.1 to 5.4 in the case that perennial woody biomass is replaced at or over the year of harvest/maturity under a nominal harvest/maturity cycle assuming that perennial cropland is harvested and regenerated back into perennial cropland. Carbon losses are estimated by multiplying annual area of harvested/replaced cropland by L_{\max} . Countries should use L_{mean} in updated Table 5.1 to 5.4 in the case that carbon removal has occurred by land use change where the age of the perennial crop removed is unknown. Carbon losses are estimated by multiplying the annual area of land conversion by L_{mean} . When perennial cropland is converted to another type of cropland, losses are reported in cropland remaining cropland. When perennial cropland is converted to non-cropland land uses, losses are reported in relevant land converted categories

Tiers 2 and 3

National level data at a finer scale, based on inventory studies or production and consumption studies according to different sources, including agricultural systems, can be used to estimate biomass loss. These can be obtained through a variety of methods, including estimating density (crown coverage) of woody vegetation from air photos (or high resolution satellite imagery) and ground-based measurement plots. Species composition, density and above-ground vs. below-ground biomass can vary widely for different cropland types and conditions and thus it may be most efficient to stratify sampling and survey plots by cropland types. General guidance on survey and sampling techniques for biomass inventories is given in Chapter 3, Annex 3A.3.

5.2.1.3 CHOICE OF ACTIVITY DATA

This section has an elaboration on the methods.

Activity data in this section refer to estimates of land areas of growing stock and harvested land with perennial woody crops. The area data are estimated using the approaches described in Chapter 3. They should be regarded as strata within the total cropland area (to keep land-use data consistent) and should be disaggregated depending on the tier used and availability of growth and loss factors. Examples of Cropland subcategories are given in updated Table 5.5.

UPDATED TABLE 5. 5 EXAMPLES OF PERENNIAL CROPLAND SUBCATEGORIES WHICH A COUNTRY MAY HAVE

Broad subcategories	Specific subcategories
Fruit orchards	Mango, Citrus, Apple, Vine (Grape, kiwifruit), Shrub (berry)
Plantation crops	Rubber, Coconut, Oil palm, Coffee, Cacao, Christmas trees, Short-rotation coppice plantations
Agroforestry systems	Hedgerow cropping (alley cropping), Improved fallow, Multi-storey systems, Home gardens, Boundary planting, Windbreaks See Table 5.3 as well.

Tier 1

Under Tier 1, annual or periodic surveys are used in conjunction with the approaches outlined in Chapter 3 to estimate the average annual area of established perennial woody crops and the average annual area of perennial woody crops that are harvested or removed. The area estimates are further sub-divided into general climate regions or soil types to match the default biomass gain and loss values. Under Tier 1 calculations, international statistics such as FAO databases, and other sources can be used to estimate the area of land under perennial woody crops.

Tier 2

Under Tier 2, more detailed annual or periodic surveys are used to estimate the areas of land in different classes of perennial woody biomass crops. Areas are further classified into relevant sub categories such that all major combinations of perennial woody crop types and climatic regions are represented with each area estimate. These area estimates must match any country-specific biomass carbon increment and loss values developed for the Tier 2 method. If country-specific finer resolution data are only partially available, countries are encouraged to extrapolate to the entire land base of perennial woody crops using sound assumptions from best available knowledge.

Tier 3

Tier 3 requires high-resolution activity data disaggregated at sub-national to fine grid scales. Similar to Tier 2, land area is classified into specific types of perennial woody crops by major climate and soil categories and other potentially important regional variables (e.g., regional patterns of management practices). Furthermore, it is *good practice* to relate spatially explicit area estimates with local estimates of biomass increment, loss rates, and management practices to improve the accuracy of estimates.

5.2.1.4 CALCULATION STEPS FOR TIER 1 AND TIER 2

No Refinement

5.2.1.5 UNCERTAINTY ASSESSMENT

No Refinement

5.2.2 Dead organic matter

No refinement

5.2.3 Soil carbon

No Refinement in the Introduction

First Order Draft

Cropland management modifies soil C stocks to varying degrees depending on how specific practices influence C input and output from the soil system (Paustian *et al.*, 1997; Bruce *et al.*, 1999; Ogle *et al.*, 2005). The main management practices that affect soil C stocks in croplands are the type of residue management, tillage management, fertilizer management (both mineral fertilizers and organic amendments), choice of crop and intensity of cropping management (e.g., continuous cropping versus cropping rotations with periods of bare fallow), irrigation management, and mixed systems with cropping and pasture or hay in rotating sequences. In addition, drainage and cultivation of organic soils reduces soil C stocks (Armentano and Menges, 1986).

General information and guidance for estimating changes in soil C stocks are found in Section 2.3.3 of Chapter 2 (including equations). That section should be read before proceeding with specific guidelines dealing with Cropland soil C stocks. The total change in soil C stocks for Cropland is estimated using Equation 2.24 (Chapter 2), which combines the change in soil organic C stocks for mineral soils and organic soils; and stock changes associated with soil inorganic C pools (Tier 3 only). This section provides specific guidance for estimating soil organic C stock changes. Soil inorganic C is fully covered by Section 2.3.3.1.

To account for changes in soil C stocks associated with *Cropland Remaining Cropland*, countries need at a minimum, estimates of the Cropland area at the beginning and end of the inventory time period. If land-use and management data are limited, aggregate data, such as FAO statistics on Cropland, can be used as a starting point, along with expert knowledge about the approximate distribution of land management systems (e.g., medium, low and high input cropping systems, etc.). Cropland management classes must be stratified according to climate regions and major soil types, which can either be based on default or country-specific classifications. This can be accomplished with overlays of land use on suitable climate and soil maps.

5.1.1.1 5.2.3.1 CHOICE OF METHOD

This section contains further elaboration on methods, updates and new guidance.

Inventories can be developed using a Tier 1, 2, or 3 method, with each successive Tier requiring more detail and resources than the previous one. It is also possible that countries will use different tiers to prepare estimates for the separate subcategories of soil C (i.e., soil organic C stocks changes in mineral soils and organic soils, and stock changes associated with soil inorganic C pools). Decision trees are provided for mineral soils (Figure 2.5) and organic soils (Figure 2.6) in Section 2.3.3.1 (Chapter 2) to assist inventory compilers with selection of the appropriate tier for their soil C inventory.

Mineral soils

Tier 1

For mineral soils, the estimation method is based on changes in soil organic C stocks over a finite period following changes in management that impact soil organic C. Equation 2.25 (Chapter 2) is used to estimate change in soil organic C stocks in mineral soils by subtracting the C stock in the last year of an inventory time period (SOC_0) from the C stock at the beginning of the inventory time period ($SOC_{(0-T)}$) and dividing by the time dependence of the stock change factors (D). In practice, country-specific data on land use and management must be obtained and classified into appropriate land management systems (e.g., high, medium and low input cropping), including tillage management, and then stratified by IPCC climate regions and soil types. Soil organic C stocks (SOC) are estimated for the beginning and end of the inventory time period using default reference carbon stocks (SOC_{ref}) and default stock change factors (F_{LU} , F_{MG} , F_1).

Tier 2

Refining Application of the Default Equations

For Tier 2, the same basic equations are used as in Tier 1 (Equation 2.25), but country-specific information is incorporated to specify better the stock change factors and reference C stocks with more disaggregation of climate regions, soil types, and/or the land management classification.

Three-Pool Steady-State C Model

The three-pool steady-state soil C model is based on estimating C inputs to soils and applying soil carbon pool specific decomposition rates that are modified by given environmental conditions and management practices. This model embraces more of the heterogeneity in soils, by subdividing soil C pool into different rates of turnover, i.e., fast (Active Pool), intermediate (Slow Pool), and long turnover times (Passive Pool).

Tier 3

Tier 3 approaches may use dynamic models and/or detailed soil C inventory measurements as the basis for estimating annual stock changes. Estimates from models are computed using coupled equations that estimate the net change of soil C. A variety of models exist (e.g., see reviews by McGill *et al.*, 1996; and Smith *et al.*, 1997). Key criteria in selecting an appropriate model include its capability of representing all of the relevant management practices/systems for croplands; model inputs (i.e., driving variables) are compatible with the availability of country-wide input data; and verification against experimental data.

A Tier 3 approach may also be developed using a measurement-based approach in which a monitoring network is sampled periodically to estimate soil organic C stock changes. A much higher density of benchmark sites will likely be needed than with models to represent adequately the combination of land-use and management systems, climate, and soil types. Additional guidance is provided in Section 2.3.3.1 of Chapter 2.

Organic soils

No Refinement. See 2013 Wetlands Supplement.

Biochar C Amendments to Mineral Soils**Tier 1**

This methodology utilizes a top-down approach in which the total amount of biochar generated and added to mineral soil is used to estimate the change in soil organic C stocks. Use Equation 2.27 to estimate the change in C stock from biochar amendments in Chapter 2, Section 2.3.3.1, Volume IV.

Tier 2

Tier 2 methods use the same definitions and equations as Tier 1, but with country-specific factors. See Section 2.3.3.1, Chapter 2, Volume IV for more information.

Tier 3

Tier 3 methods can be used to account for GHG sources and sinks not captured in Tiers 1 or 2, such as priming, changes to N₂O or CH₄ fluxes from soils, and changes to net primary production. More information on Tier 3 methods is provided in Section 2.3.3.1 of Chapter 2, Volume IV.

5.2.3.2 CHOICE OF STOCK CHANGE AND EMISSION FACTORS

This section contains further elaboration on methods, updates and new guidance.

Mineral soils**Tier 1**

Table 5.6 provides Tier 1 approach default stock change factors for land use (F_{LU}), input (F_I) and management (F_{MG}). The method and studies that were used to derive the default stock change factors are provided in Annex 5A.1 and References. The default time period for stock changes (D) is 20 years and management practice is assumed to influence stocks to a depth of 30 cm, which is also the depth for the reference soil C stocks in Table 2.3 (Chapter 2).

Tier 2*Refining Application of Default Equations*

A Tier 2 approach entails the estimation of country-specific stock change factors. Derivation of input (F_I) and management factors (F_{MG}) are based on comparisons to medium input and intensive tillage, respectively, because they are considered the nominal practices in the IPCC default management classification (see Choice of Activity Data). It is *good practice* to derive values for a higher resolution classification of management, climate and soil types if there are significant differences in the stock change factors among more disaggregated categories based on an empirical analysis. Reference C stocks can also be derived from country-specific data in a Tier 2 approach. Additional guidance is provided in Chapter 2, Section 2.3.3.1.

Reference C stocks can also be derived from country-specific data in a Tier 2 approach. However, reference values must be consistent across land-use sectors (i.e., Forest Land, Cropland, Grassland, Settlements, Other land), which requires coordination among the various teams conducting soil C inventories for AFOLU.

The depth for evaluating soil C stock changes can also be extended with the Tier 2 method. This will require extending the depth of the reference C stocks (SOC_{REF}) and stock change factors for all land uses (i.e., F_{LU} , F_I , and F_{MG}) to ensure

First Order Draft

consistency. Variable depths between reference stocks and stock change factors are likely to introduce biases into the inventory estimates that are computed using Equation 2.25.

For the case of land use change to a system that is increasing in C, such as Croplands converted to Grasslands, a Tier 2 method may be a more accurate way to estimate the increase of soil C stocks to native levels. The Tier 1 method may overestimate soil C stock increases on an annual basis (e.g., Villarino et al., 2014).

Furthermore, inventories may be improved by estimating carbon stocks on a mass equivalency basis when deriving country-specific factors for F_{LU} . This is because the soil weight in a certain soil depth changes with the various operations associated with land use change, for example uprooting, land leveling, and rain compaction due to the disappearance of the cover of tree canopy. In addition, cropland soils usually tend to have relatively higher density than the soils in forest land and possibly grasslands or wetlands. Settlement management may also impact the soil bulk density. In such case, the comparison of the soil carbon stocks between the cropland, settlement, grassland, wetland, or forest land within the same depth is not appropriate. It is more robust to compare the carbon stock on an equivalent mass basis, with the stock change calculated on the same weight soil. This method is also recommended when the bulk density between cropland, grassland, wetland, settlement and forest land is remarkably different even if the site is within close proximity. However, it is important to realize that all measurements and associated stock change factors across all land uses must be on an equivalent mass basis if this method is applied. This will be challenging and possibly not even practical unless done comprehensively for all land uses.

Three-Pool Steady-State C Model

Default parameters are provided for the three-pool steady-state C pool equations (Chapter 2, Section 2.3.3.1, Table 2.4), but parameters may be revised if experimental data are available to test the model. The average lignin and nitrogen contents of the C input is also required to estimate the size of the three C pools (See Table 5.7).

Tier 3

Constant stock change rate factors *per se* are less likely to be estimated in favor of variable rates that more accurately capture land-use and management effects. See Chapter 2, Section 2.3.3.1 for further discussion.

Organic soils

No Refinement. See 2013 Wetlands Supplement.

Biochar C Amendments to Mineral Soils

Tier 1

Default emission factors are provided in Section 2.3.3.1, Chapter 2, Volume IV.

Tier 2

Tier 2 emission factors may be further disaggregated relative to the default factors based on variation in environmental conditions, such as the climate and soil types, in addition to variation associated with the biochar production methods. See Section 2.3.3.1, Chapter 2, Volume IV for more information.

Tier 3

Tier 3 methods are country-specific and may involve empirical or process-based models to account for a broader set of impacts of biochar amendments. More information on Tier 3 methods is provided in Section 2.3.3.1, Chapter 2, Volume IV.

UPDATED TABLE 5. 6 RELATIVE STOCK CHANGE FACTORS (FLU, FMG, AND FI) (OVER 20 YEARS) FOR DIFFERENT MANAGEMENT ACTIVITIES ON CROPLAND						
Factor value type	Level	Temperature regime	Moisture regime ¹	IPCC defaults	Error ^{2,3}	Description
Land use (F _{LU})	Long-term cultivated	Temperate/Boreal	Dry	TBD	TBD	Represents area that has been continuously managed for >20 yrs, to predominantly annual crops. Input and tillage factors are also applied to estimate carbon stock changes. Land-use factor was estimated relative to use of full tillage and nominal ('medium') carbon input levels.
			Moist	TBD	TBD	
		Tropical	Dry	TBD	TBD	
			Moist/Wet	TBD	TBD	
		Tropical montane ⁴	n/a	TBD	TBD	
Land use (F _{LU})	Paddy rice	All	Dry and Moist/Wet	TBD	TBD	Long-term (> 20 year) annual cropping of wetlands (paddy rice). Can include double-cropping with non-flooded crops. For paddy rice, tillage and input factors are not used.
Land use (F _{LU})	Perennial/Tree Crop	All	Dry and Moist/Wet	TBD	TBD	Long-term perennial tree crops such as fruit and nut trees, coffee and cacao.
Land use (F _{LU})	Set aside (< 20 yrs)	Temperate/Boreal and Tropical	Dry	TBD	TBD	Represents temporary set aside of annually cropland (e.g., conservation reserves) or other idle cropland that has been revegetated with perennial grasses.
			Moist/Wet	TBD	TBD	
		Tropical montane	n/a	TBD	TBD	
Tillage (F _{MG})	Full	All	Dry and Moist/Wet	TBD	TBD	Substantial soil disturbance with full inversion and/or frequent (within year) tillage operations. At planting time, little (e.g., <30%) of the surface is covered by residues.
Tillage (F _{MG})	Reduced	Temperate/Boreal	Dry	TBD	TBD	Primary and/or secondary tillage but with reduced soil disturbance (usually shallow and without full soil inversion). Normally leaves surface with >30% coverage by residues at planting.
			Moist	TBD	TBD	
		Tropical	Dry	TBD	TBD	
			Moist/Wet	TBD	TBD	
		Tropical montane ⁴	n/a	TBD	TBD	
Tillage (F _{MG})	No-till	Temperate/Boreal	Dry	TBD	TBD	Direct seeding without primary tillage, with only minimal soil disturbance in the seeding zone. Herbicides are typically used for weed control.
			Moist	TBD	TBD	
		Tropical	Dry	TBD	TBD	
			Moist/Wet	TBD	TBD	
		Tropical montane ⁴	n/a	TBD	TBD	

First Order Draft

UPDATED - TABLE 5.6 (CONTINUED)						
Relative stock change factors (FLU, FMG, AND FI) (over 20 years) for different management activities on cropland						
Factor value type	Level	Temperature regime	Moisture regime ¹	IPCC defaults	Error ^{2,3}	Description
Input (Fi)	Low	Temperate/Boreal	Dry	TBD	TBD	Low residue return occurs when there is due to removal of residues (via collection or burning), frequent bare-fallowing, production of crops yielding low residues (e.g., vegetables, tobacco, cotton), no mineral fertilization or N-fixing crops.
			Moist	TBD	TBD	
		Tropical	Dry	TBD	TBD	
			Moist/Wet	TBD	TBD	
		Tropical montane ⁴	n/a	TBD	TBD	
Input (Fi)	Medium	All	Dry and Moist/Wet	TBD	TBD	Representative for annual cropping with cereals where all crop residues are returned to the field. If residues are removed then supplemental organic matter (e.g., manure) is added. Also requires mineral fertilization or N-fixing crop in rotation.
Input (Fi)	High without manure	Temperate/Boreal and Tropical	Dry	TBD	TBD	Represents significantly greater crop residue inputs over medium C input cropping systems due to additional practices, such as production of high residue yielding crops, use of green manures, cover crops, improved vegetated fallows, irrigation, frequent use of perennial grasses in annual crop rotations, but without manure applied (see row below).
			Moist/Wet	TBD	TBD	
		Tropical montane ⁴	n/a	TBD	TBD	
Input (Fi)	High – with manure	Temperate/Boreal and Tropical	Dry	TBD	TBD	Represents significantly higher C input over medium C input cropping systems due to an additional practice of regular addition of animal manure.
			Moist/Wet	TBD	TBD	
		Tropical montane ⁴	n/a	TBD	TBD	
TBD – To be determined based on literature review for second order draft. These updates to may also require some changes to the descriptions of practices. See Annex 5A1 for more information.						
¹ Where data were sufficient, separate values were determined for temperate and tropical temperature regimes; and dry, moist, and wet moisture regimes. Temperate and tropical zones correspond to those defined in Chapter 3; wet moisture regime corresponds to the combined moist and wet zones in the tropics and moist zone in temperate regions.						
² ± two standard deviations, expressed as a percent of the mean; where sufficient studies were not available for a statistical analysis to derive a default, uncertainty was assumed to be ± 50% based on expert opinion. NA denotes ‘Not Applicable’, where factor values constitute defined reference values, and the uncertainties are reflected in the reference C stocks and stock change factors for land use.						
³ This error range does not include potential systematic error due to small sample sizes that may not be representative of the true impact for all regions of the world.						
⁴ There were not enough studies to estimate stock change factors for mineral soils in the tropical montane climate region. As an approximation, the average stock change between the temperate and tropical regions was used to approximate the stock change for the tropical montane climate.						

NEW GUIDANCE TABLE 5.7 DEFAULT VALUES FOR NITROGEN AND LIGNIN CONTENTS IN CROPS FOR THREE-POOL STEADY-STATE C MODEL				
Crops	N content of above-ground residues	N content of below-ground residues	Lignin content of above-ground residues	Lignin content of below-ground residues
Generic value for crops not indicated below	TBD	TBD	TBD	TBD
Generic Grains	0.006	0.009	TBD	TBD
Winter Wheat	0.006	0.009	TBD	TBD
Spring Wheat	0.006	0.009	TBD	TBD
Barley	0.007	0.014	TBD	TBD
Oats	0.007	0.008	TBD	TBD
Maize	0.007	0.007	TBD	TBD
Rye	0.005	0.011	TBD	TBD
Rice	0.007	TBD	TBD	TBD
Millet	0.007	TBD	TBD	TBD
Sorghum	0.007	0.006	TBD	TBD
Beans and Pulses	0.01	0.01	TBD	TBD
Soybeans	0.008	0.008	TBD	TBD
Potatoes and Tubers	0.019	0.014	TBD	TBD
Peanuts	0.016	TBD	TBD	TBD
Alfalfa and Legume Hay	0.027	0.019	TBD	TBD
Non-legume hay	0.015	0.012	TBD	TBD
TBD – To be determined based on literature review for second order draft. These updates to may also require some changes to the descriptions of practices.				

5.2.3.3 CHOICE OF ACTIVITY DATA

This section contains further elaboration on methods, updates and new guidance.

Mineral soils

Tier 1

Cropland systems are classified by practices that influence soil C storage. The default management classification system is provided in Figure 5.1. Inventory compilers should use this classification to categorize management systems in a manner consistent with the default Tier 1 stock change factors. This classification may be further developed for Tiers 2 and 3 approaches. In general, practices that are known to increase C storage, such as irrigation, mineral fertilization, organic amendments, cover crops and high residue yielding crops, have higher inputs, while practices that decrease C storage, such as residue burning/removal, bare fallow, and low residue crop varieties, have lower inputs. These practices are used to categorize management systems and then estimate the change in soil organic C stocks. Practices should not be considered that are used in less than 1/3 of a given cropping sequence (i.e., crop rotation), which is consistent with the classification of experimental data used to estimate the default stock change

First Order Draft

factors. Rice production, perennial croplands, and set-aside lands (i.e., lands removed from production) are considered unique management systems (see below).

Each of the annual cropping systems (low input, medium input, high input, and high input w/organic amendment) are further subdivided based on tillage management. Tillage practices are divided into no-till (direct seeding without primary tillage and only minimal soil disturbance in the seeding zone; herbicides are typically used for weed control), reduced tillage (primary and/or secondary tillage but with reduced soil disturbance that is usually shallow and without full soil inversion; normally leaves surface with >30% coverage by residues at planting) and full tillage (substantial soil disturbance with full inversion and/or frequent, within year tillage operations, while leaving <30% of the surface covered by residues at the time of planting). It is *good practice* only to consider reduced and no-till if they are used continuously (every year) because even an occasional pass with a full tillage implement will significantly reduce the soil organic C storage expected under the reduced or no-till regimes (Pierce *et al.*, 1994; Smith *et al.*, 1998). Assessing the impact of rotational tillage systems (i.e., mixing reduced, no-till and/or full tillage practices) on soil C stocks will require a Tier 2 method.


The main types of land-use activity data are: i) aggregate statistics (Approach 1), ii) data with explicit information on land-use conversions but without specific geo-referencing (Approach 2), or iii) data with explicit information on land-use conversions and geo-referencing (Approach 3), such as land-use and management inventories making up a statistically-based sample of a country's land area (see Chapter 3 for discussion of approaches). At a minimum, globally available land-use and crop production statistics, such as FAO databases (<http://faostat.fao.org/>), provide annual compilations of total land area by major land-uses, select management data (e.g., irrigated vs. non-irrigated cropland), land area in 'perennial' crops (i.e., vineyards, perennial herbaceous crops, and tree-based crops such as orchards) and annual crops (e.g., wheat, rice, maize, sorghum, etc.). FAO databases would be an example of aggregate data (Approach 1).

Management activity data supplement the land-use data, providing information to classify management systems, such as crop types and rotations, tillage practices, irrigation, manure application, residue management, etc. These data can also be aggregate statistics (Approach 1) or information on explicit management changes (Approach 2 or 3). Where possible, it is *good practice* to determine the specific management practices for land areas associated with cropping systems (e.g., rotations and tillage practice), rather than only area by crop. Remote sensing data are a valuable resource for land-use and management activity data, and potentially, expert knowledge is another source of information for cropping practices. It is *good practice* to elicit expert knowledge using methods provided in Volume 1, Chapter 2 (eliciting expert knowledge).

National land-use and resource inventories, based on repeated surveys of the same locations, constitute activity data gathered using Approach 2 or 3, and have some advantages over aggregated land-use and cropland management data (Approach 1). Time series data can be more readily associated with a particular cropping system (i.e., combination of crop type and management over a series of years), and the soil type can be determined by sampling or by referencing the location to a suitable soil map. Inventory points that are selected based on an appropriate statistical design also enable estimates of the variability associated with activity data, which can be used as part of a formal uncertainty analysis. An example of a survey using Approach 3 is the National Resource Inventory in the U.S. (Nusser and Goebel, 1997).

Activity data require additional in-country information to stratify areas by climate and soil types. If such information has not already been compiled, an initial approach would be to overlay available land cover/land-use maps (of national origin or from global datasets such as IGBP_DIS) with soil and climate maps of national origin or global sources, such as the FAO Soils Map of the World and climate data from the United Nations Environmental Program. A detailed description of the default climate and soil classification schemes is provided in Chapter 3, Annex 3A.5. The soil classification is based on soil taxonomic description and textural data, while climate regions are based on mean annual temperatures and precipitation, elevation, occurrence of frost, and potential evapotranspiration.

Figure 5. 1 Classification scheme for cropping systems. In order to classify cropland management systems, the inventory compiler should start at the top and proceed through the diagram answering questions (move across branches if answer is yes) until reaching a terminal point on the diagram. The classification diagram is consistent with default stock change factors in Table 5.5. C input classes (i.e., low, medium, high and high with organic amendment) are further subdivided by tillage practice.



Note:

1: Does not typically include grazing of residues in the field.

2: e.g. cotton, vegetables and tobacco.

3: Practices that increase C input above the amount typically generated by the low residues yielding varieties such as using organic amendments, cover crops/green manures, and mixed crop/grass systems.

4: Practices that increase C input by enhancing residue production, such as using irrigation, cover crops/green manures, vegetated fallows, high residue yielding crops, and mixed crop/grass systems.

5 Perennial cover without frequent harvest.

Note: Only consider practices, such as irrigation, residue burning/removal, mineral fertilizers, N-fixing crops, organic amendment, cover crops/green manures, low residue crop, or fallow, if used in at least 1/3 of cropping rotation sequence.

First Order Draft

Tier 2*Refining Application of Default Equations*

Tier 2 approaches are likely to involve a more detailed stratification of management systems than in Tier 1 (see Figure 5.1) if sufficient data are available. This can include further subdivisions of annual cropping input categories (i.e., low, medium, high, and high with amendment), rice cultivation, perennial cropping systems, and set-asides. It is *good practice* to further subdivide default classes based on empirical data that demonstrates significant differences in soil organic C storage among the proposed categories. In addition, Tier 2 approaches can involve a finer stratification of climate regions and soil types.

For Tier 2, the specific definitions of management and input factors are typically made to match available activity data on how an activity affects C stocks. For example, if a country has management factors related to specific tillage practices that involve a mix of intensities over time, then the country will also need activity data on those specific tillage practices to apply the country-specific factors.

Three-Pool Steady-State C Model

This method requires soil C input data based on the amount of biomass that is converted to dead organic matter annually. This rate will vary depending on the crop production, management activity, and other environmental variables. Removals or reductions in dead organic matter are subtracted from the C input amount, which could occur with livestock grazing, grassland burning, or harvesting of grass for feed or bioenergy. Additions of C, particularly organic amendments such as manure, are included in the estimate of C input.

It is good practice to estimate C input using country-specific methods in order to produce more accurate estimates. If country-specific methods are not available, Equation 5.1 can be used to estimate C inputs with factors provided in Section 11.2.1.2 of Chapter 11, Volume 4 (See Section 11.2.1.2 for more information).

Equation 5.1 Cropland litter carbon input for three-pool steady-state C model

$$C_{input} = \sum_T [AGR_{(T)} * (1 - Frac_{Remove(T)} - (Frac_{Burnt(T)} * C_f))] + [BGR_{(T)}]$$

$$AGR_{(T)} = Crop_{(T)} * R_{AG(T)} * Area_{(T)} * Frac_{Renew(T)}$$

$$BGR_T = Crop_T * (1 + R_{AG(T)}) * R:S_{(T)} * Area_{(T)} * Frac_{Renew(T)}$$

Where:

C_{input} = annual amount of N in crop residues (above and below ground), including N-fixing crops, and from forage/pasture renewal, returned to soils annually, kg N yr⁻¹

$AGR_{(T)}$ = annual total amount of above-ground crop residue for crop T , kg d.m. ha⁻¹. (Use factors in Table 11.2, Chapter 11, or alternatively, the amount can be calculated using the method and data in Table 11.3, Chapter 11)

$Frac_{Remove(T)}$ = fraction of above-ground residues of crop T removed annually for purposes such as feed, bedding and construction, dimensionless. Survey of experts in country is required to obtain data. If data for $Frac_{Remove}$ are not available, assume no removal

$Frac_{Burnt(T)}$ = fraction of annual harvested area of crop T burnt, dimensionless

C_f = combustion factor (dimensionless) (refer to Chapter 2, Table 2.7)

$BGR_{(T)}$ = annual total amount of belowground crop residue for crop T , kg d.m. ha⁻¹

$Crop_{(T)}$ = harvested annual dry matter yield for crop T , kg d.m. ha⁻¹

$R_{AG(T)}$ = ratio of above-ground residues dry matter ($AG_{DM(T)}$) to harvested yield for crop T ($Crop_{(T)}$), kg d.m. (kg d.m.)⁻¹, (Table 11.2)

Area_(T) = total annual area harvested of crop *T*, ha yr⁻¹

Frac_{Renew}(*T*) = fraction of total area under crop *T* that is renewed annually ¹, dimensionless. For countries where pastures are renewed on average every *X* years, Frac_{Renew} = 1/*X*. For annual crops Frac_{Renew} = 1

R:S(*T*) = ratio of below-ground root biomass to above-ground biomass for crop *T*, kg d.m. (kg d.m.)⁻¹, (Table 11.2)

T = crop or forage type

Data on crop yield statistics (yields and area harvested, by crop) may be obtained from national sources. If such data are not available, FAO publishes data on crop production: (<http://faostat.fao.org/>). Tillage management data are also required (proportion of full tillage, reduced tillage and no-till), and irrigation data for any lands that are provided supplement water (proportion of land). Monthly average temperature, precipitation and potential evapotranspiration is needed for each grid cell or region. This information is available from global datasets, such as the CRU climate dataset (<https://crudata.uea.ac.uk/cru/data/hrg/>), if country-specific data are not available. The average sand content is needed for each grid cell or region, which is available from Harmonized World Soil Database (<http://webarchive.iiasa.ac.at/Research/LUC/External-World-soil-database/HTML/>).

Tier 3

For application of dynamic models and/or a direct measurement-based inventory in Tier 3, similar or more detailed data on the combinations of climate, soil, topographic and management data are needed, relative to the Tiers 1 and 2 methods, but the exact requirements will depend on the model or measurement design.

Organic soils

No Refinement. See 2013 Wetlands Supplement.

Biochar C Amendments to Mineral Soils

Tier 1

The activity data required for the Tier 1 method includes the total quantities of biochar distributed for amendment to mineral soils. These data must be disaggregated by production type, where production type is defined as a process utilizing a specific feedstock type, and a specific conversion process (gasification, or high-, medium-, or low-temperature pyrolysis; Tables 2.4 and 2.5). In case data on the temperature of pyrolysis are unavailable, default factors for uncontrolled or unspecified pyrolysis temperatures are provided in Section 2.3.3.1 of Chapter 2, Volume IV. Changes in soil C associated with biochar amendments is considered to occur where it is incorporated into soil. However, due to the distributed nature of the land sector in which this can take place, inventory compilers may not have access to data on when or where biochar C amendments occur. Therefore, for the purposes of Tier 1 method, inventory compilers can rely on centralized records from biochar producers, importers, exporters or distributors, recording the quantity of biochar that has been provided to the land use sector for use as a soil amendment in the country. Note that exported biochar is not included in the total amount of biochar amended to soils in the country. Inventory compilers may further disaggregate amendments by land use if the data are available.

Tier 2

Tier 2 methods have the same activity data requirements as Tier 1 (quantities of biochar distributed for incorporation into mineral soils, disaggregated by production type). Additionally, activity data on the amount of biochar amendments may be disaggregated by climate zones and/or soil types if country-specific factors are disaggregated by these environmental variables. The additional climate and soil activity data may be obtained with a survey of biochar distributors and land managers.

Tier 3

The additional activity data required to support a Tier 3 method will depend on which processes are represented and environmental variables that are required as input to the model. Priming, soil GHG emissions, and plant production responses to biochar all vary with biochar type, climate, and soil type. Furthermore, soil GHG emissions and plant production responses also vary with crop type and management. Therefore, Tier 3 methods may require environmental data on climate zones, soil types, crop types and crop management systems (such as nitrogen fertilizer application

¹ This term is included in the equation to account for N release and the subsequent increases in N₂O emissions (e.g., van der Weerden *et al.*, 1999; Davies *et al.*, 2001), from renewal/cultivation of grazed grass or grass/clover pasture and other forage crops.

First Order Draft

rates, and whether soils are flooded for paddy rice production), in addition to the amount of biochar amendments in each of the individual combinations of strata for the environmental variables. More detailed activity data specifying the process conditions for biochar production or the physical and chemical characteristics of the biochar may also be required (such as surface area, cation exchange capacity, pH, and ash content).

5.2.3.4 CALCULATION STEPS FOR TIER 1

This section will be updated for the second order draft based on new factors, and also will provide new guidance.

Mineral soils

The steps for estimating SOC_0 and $SOC_{(0-T)}$ and net soil C stock change per ha for *Cropland Remaining Cropland* on mineral soils are as follows:

Step 1: Organize data into inventory time periods based on the years in which activity data were collected (e.g., 1990 to 1995, 1995 to 2000, etc.)

Step 2: Determine the amount *Cropland Remaining Cropland* by mineral soil types and climate regions in the country at the beginning of the first inventory time period. The first year of the inventory time period will depend on the time step of the activity data (0-T; e.g., 5, 10 or 20 years ago).

Step 3: Classify each Cropland into the appropriate management system using Figure 5.1.

Step 4: Assign a native reference C stock values (SOC_{REF}) from Table 2.3 based on climate and soil type.

Step 5: Assign a land-use factor (F_{LU}), management factor (F_{MG}) and C input levels (F_I) to each Cropland based on the management classification (Step 2). Values for F_{LU} , F_{MG} and F_I are given in Table 5.6.

Step 6: Multiply the factors (F_{LU} , F_{MG} , F_I) by the reference soil C stock (SOC_{REF}) to estimate an ‘initial’ soil organic C stock ($SOC_{(0-T)}$) for the inventory time period.

Step 7: Estimate the final soil organic C stock (SOC_0) by repeating Steps 1 to 5 using the same native reference C stock (SOC_{REF}), but with land-use, management and input factors that represent conditions for each cropland in the last (year 0) inventory year.

Step 8: Estimate the average annual change in soil organic C stocks for *Cropland Remaining Cropland* ($\Delta C_{Mineral}$) by subtracting the ‘initial’ soil organic C stock ($SOC_{(0-T)}$) from the final soil organic C stock (SOC_0), and then dividing by the time dependence of the stock change factors (i.e., 20 years using the default factors). If an inventory time period is greater than 20 years, then divide by the difference in the initial and final year of the time period.

Step 9: Repeat steps 2 to 8 if there are additional inventory time periods (e.g., 1990 to 2000, 2001 to 2010, etc.).

A numerical example is given below for *Cropland Remaining Cropland* on mineral soils, using Equation 2.25 and default reference C stocks (Table 2.3) and stock change factors (Table 5.6).

Example: The following example shows calculations for aggregate areas of cropland soil carbon stock change. In a warm temperate wet climate on Mollisol soils, there are 1Mha of permanent annual cropland. The native reference carbon stock (SOCref) for the region is 88 tonnes C ha⁻¹. At the beginning of the inventory calculation period (in this example, 10 yrs earlier in 1990), the distribution of cropland systems were 400,000 ha of annual cropland with low carbon input levels and full tillage and 600,000 ha of annual cropland with medium input levels and full tillage. Thus, initial soil carbon stocks for the area were: 400,000 ha • (88 tonnes C ha⁻¹ • 0.69 • 1 • 0.92) + 600,000 ha • (88 tonnes C ha⁻¹ • 0.69 • 1 • 1) = 58.78 million tonnes C. In the last year of the inventory time period (in this example, the last year is 2000), there are: 200,000 ha of annual cropping with full tillage and low C input, 700,000 ha of annual cropping with reduced tillage and medium C input, and 100,000 ha of annual cropping with no-till and medium C input. Thus, total soil carbon stocks in the inventory year are: 200,000 ha • (88 tonnes C ha⁻¹ • 0.69 • 1 • 0.92) + 700,000 ha • (88 tonnes C ha⁻¹ • 0.69 • 1.08 • 1) + 100,000 ha • (88 tonnes C ha⁻¹ • 0.69 • 1.15 • 1) = 64.06 million tonnes C. Thus, the average annual stock change over the period for the entire area is: 64.06 – 58.78 = 5.28 million tonnes/20 yr = 264,000 tonnes C per year soil C stock increase (Note: 20 years is the time dependence of the stock change factor, i.e., factor represents annual rate of change over 20 years).

Organic soils

No Refinement. See 2013 Wetlands Supplement.

Biochar C Amendments to Mineral Soils

Will be provided in second order draft after factors are derived.

5.2.3.5 UNCERTAINTY ASSESSMENT

No Refinement

5.2.4 Non-CO₂ greenhouse gas emissions from biomass burning

No Refinement

5.3 LAND CONVERTED TO CROPLAND

No Refinement in the Introduction

5.3.1 Biomass

5.3.1.1 CHOICE OF METHOD

This section provides elaboration on how to calculate ΔC_G .

This section provides guidance on methods for calculating carbon stock change in biomass due to the conversion of land from natural conditions and other uses to Cropland, including deforestation and conversion of pasture and grazing lands to Cropland. The methods require estimates of carbon in biomass stocks prior to and following conversion, based on estimates of the areas of lands converted during the period between land-use surveys. As a result of conversion to Cropland, it is assumed (in Tier 1) that the dominant vegetation is removed entirely leading to emissions, resulting in near zero amounts of carbon remaining in biomass. Some type of cropping system is planted soon thereafter increasing the amount of carbon stored in biomass. The difference between initial and final biomass carbon pools is used to calculate carbon stock change from land-use conversion; and in subsequent years accumulations and losses in perennial woody biomass in Cropland are counted using methods in Section 5.2.1 (Cropland Remaining Cropland).

It is *good practice* to consider all carbon pools (i.e., above ground and below ground biomass, dead organic matter, and soils) in estimating changes in carbon stocks in *Land Converted to Cropland*. Currently, there is insufficient information to provide a default approach with default parameters to estimate carbon stock change in dead organic matter (DOM) pools². DOM is unlikely to be important except in the year of conversion. It is assumed that there will be no DOM in Cropland. In addition, the methodology below considers only carbon stock change in above-ground biomass since limited data are available on below-ground carbon stocks in perennial Cropland.

The *IPCC Guidelines* describe increasingly sophisticated alternatives that incorporate greater detail on the areas of land converted, carbon stocks on lands, and loss of carbon resulting from land conversions. It is *good practice* to adopt the appropriate tier depending on key source analysis, data availability and national circumstances. All countries should strive for improving inventory and reporting approaches by advancing to the highest tier possible given national circumstances. It is *good practice* for countries to use a Tier 2 or Tier 3 approach if carbon emissions and removals in *Land Converted to Cropland* is a key category and if the sub-category of biomass is considered significant based on principles outlined in Volume 1, Chapter 4. Countries should use the decision tree in Figure 1.3 to help with the choice of method. *Land Converted to Cropland* is likely to be a key category for many countries and further biomass is likely to be a key source.

Tier 1

The Tier 1 method follows the approach in Chapter 4 (Forest Land) where the amount of biomass that is cleared for cropland is estimated by multiplying the area converted in one year by the average carbon stock in biomass in the Forest Land or Grassland prior to conversion. It is *good practice* to account completely for all land conversions to Cropland. Thus, this section elaborates on the method such that it includes different initial uses, including but not limited to forests.

Equation 2.15 in Chapter 2 summarises the major elements of a first-order estimation of carbon stock change from land-use conversion to Cropland. Average carbon stock change on a per hectare basis is estimated for each type of conversion. The average carbon stock change is equal to the carbon stock change due to the removal of biomass from the initial land use (i.e., carbon in biomass immediately after conversion minus the carbon in biomass prior to conversion), plus carbon stocks from one year of growth in Cropland following conversion. It is necessary to account only for any woody vegetation that replaces the vegetation that was cleared during land-use conversion. The *GPG-LULUCF* combines carbon in biomass after conversion and carbon in biomass that grows on the land following

² Any litter and dead wood pools (estimated using the methods described in Chapter 2, Section 2.3.2) should be assumed oxidized following land conversion.

conversion into a single term. In this method, they are separated into two terms, B_{AFTER} and ΔC_G to increase transparency.

As described in section 5.3.1.1., at Tier 1, carbon stocks in biomass immediately after conversion (B_{AFTER}) are assumed to be zero, since the land is cleared of all vegetation before planting crops. Average carbon stock change per hectare for a given land-use conversion is multiplied by the estimated area of lands undergoing such a conversion in a given year. In subsequent years, change in biomass of annual crops is considered zero because carbon gains in biomass from annual growth are offset by losses from harvesting. Changes in biomass of perennial woody crops are counted following the methodology in Section 2.3.1.1 (Change in carbon stocks in biomass in land remaining in a land-use category) and Section 5.2.1 (Change in carbon stocks in biomass in cropland remaining cropland). Thus, carbon gain of an annual crop is estimated only for the first year following a conversion, whereas, carbon gains and losses of perennial woody crop may also occur in subsequent years up to 20 years (at maximum).

The default assumption for Tier 1 is that all carbon in biomass removed is lost to the atmosphere through burning or decay processes either on-site or off-site. Tier 1 calculations do not differentiate immediate emissions from burning and other conversion related losses.

Tier 2

The Tier 2 calculations are structurally similar to Tier 1, with the following distinctions. First, Tier 2 relies largely on country-specific estimates of the carbon stocks in initial and final land uses rather than the default data. Area estimates for *Land Converted to Cropland* are disaggregated according to original vegetation (e.g., from Forest Land or Grassland) at finer spatial scales to capture regional and crop systems variations in country-specific carbon stocks values.

Second, Tier 2 may modify the assumption that carbon stocks immediately following conversion are zero. This enables countries to take into account land-use transitions where some, but not all, vegetation from the original land use is removed.

Third, under Tier 2, it is *good practice* to apportion carbon losses to burning and decay processes if applicable. Emissions of carbon dioxide occur as a result of burning and decay in land-use conversions. Further, non- CO_2 trace gas emissions occur as a result of burning. By partitioning losses to burning and decay, countries can also calculate non- CO_2 trace gas emissions from burning (Section 5.3.4).

The immediate impacts of land conversion activities on the five carbon stocks can be summarized in a disturbance matrix, which describes the retention, transfers and releases of carbon in the pools in the original ecosystem following conversion to Cropland. A disturbance matrix defines for each pool the proportion that remains in that pool and the proportion that is transferred to other pools. A small number of transfers are possible, and are outlined in a disturbance matrix in Table 5.9. The disturbance matrix ensures consistency of the accounting of all carbon pools.

Biomass transfers to dead wood and litter can be estimated using Equation 2.20.

Tier 3

The Tier 3 method is similar to Tier 2, with the following distinctions: i) rather than relying on average annual rates of conversion, countries can use direct estimates of spatially disaggregated areas converted annually for each initial and final land use; ii) carbon densities and soil carbon stock change are based on locally specific information, which makes possible a dynamic link between biomass and soil; and iii) biomass volumes are based on actual inventories. The transfer of biomass, to dead wood and litter following land-use conversion can be estimated using Equation 2.20.

Table 5. 8 Text to be provided in SOL

TABLE 5.9 EXAMPLE OF A SIMPLE DISTURBANCE MATRIX (TIER 2) FOR THE IMPACTS OF LAND CONVERSION ACTIVITIES ON CARBON POOLS								
From \ To	Above-ground biomass	Below-ground biomass	Dead wood	Litter	Soil organic matter	Harvested wood products	Atmosphere	Sum of row (must equal 1)
Above-ground biomass								
Below-ground biomass								
Dead wood								
Litter								
Soil organic matter								
Enter the proportion of each pool on the left side of the matrix that is transferred to the pool at the top of each column. All of the pools on the left side of the matrix must be fully accounted, so the values in each row must sum to 1. Impossible transitions are blacked out.								

5.3.1.2 CHOICE OF EMISSION FACTORS

This section provides elaboration on methods and updates.

The emission/removal factors needed for the default method are: carbon stocks before conversion in the initial land use and after conversion to Cropland; and growth in biomass carbon stock from one year of cropland growth.

Tier 1

Default biomass carbon stock in initial land-use categories (B_{BEFORE}) mainly Forest Land and Grassland are provided in Updated Table 5.10. Initial land-use based carbon stocks should be obtained for different Forest Land or Grassland categories based on biome type, climate, soil management systems, etc. It is assumed that all biomass is cleared when preparing a site for cropland use, thus, the default for B_{AFTER} is 0 tonne C ha⁻¹.

In addition, a value is needed for carbon stocks after one year of growth in crops planted after conversion (ΔC_G). Updated Table 5.11 provides general defaults for ΔC_G while updated table 5.3 provides defaults for specific crops. Separate defaults are provided for annual non-woody crops and perennial woody crops. For lands planted in annual crops, the default value of ΔC_G is [5.0] [4.7] tonnes of C per hectare, based on the original *IPCC Guidelines* recommendation of 10 tonnes of dry biomass per hectare (dry biomass has been converted to tonnes carbon in Table 5.11). The total accumulation of carbon in perennial woody biomass will, over time, exceed that of the default carbon stock for annual cropland. However, default values provided in this section are for one year of growth immediately following conversion, which usually give lower carbon stocks for perennial woody crops compared to annual crops.

UPDATED TABLE 5. 10 DEFAULT BIOMASS CARBON STOCKS REMOVED DUE TO LAND CONVERSION TO CROPLAND

Land-use category	Carbon stock in biomass before conversion (B_{Before}) (tonnes C ha ⁻¹)	Error range #
Forest Land	See Chapter 4 Tables 4.7 to 4.12 for carbon stocks in a range of forest types by climate regions. Stocks are in terms of dry matter. Multiply values by a carbon fraction (CF) in Table 4.3 consistent with what used in forest land estimation to convert dry matter to carbon.	See Section 4.3 (Land Converted to Forest Land)
Grassland	See Chapter 6 Table 6.4 for carbon stocks in a range of grassland types by climate regions. Multiply default carbon fraction (CF) 0.47 (for herbaceous biomass for Grassland, see page 6.29, Chapter 6 of the 2006 guidelines to convert dry matter to carbon.	$\pm 75\%$ [This range may change based on updated Table 6.4]
# Represents a nominal estimate of error, equivalent to two times standard deviation, as a percentage of the mean.		

UPDATED TABLE 5. 11 DEFAULT BIOMASS CARBON STOCKS PRESENT ON LAND CONVERTED TO CROPLAND IN THE YEAR FOLLOWING CONVERSION

Crop type by climate region	Ecological zone	Continent	Cropping system	Carbon stock in biomass after one year (ΔC_G) (tonnes C ha ⁻¹)	Error range#
Annual cropland	All	All	Annual cropland	[5.0][4.7]	$\pm 75\%$
[others]	TBD	TBD	TBD	TBD	TBD
Perennial cropland					
Tropical	Highland	All	General perennial agroforestry	3.9	$\pm 75\%$
Tropical	Dry	All	General perennial agroforestry	2.6	$\pm 75\%$
Tropical	Moist	All	General perennial agroforestry	6.1	$\pm 75\%$
Tropical	Wet	All	General perennial agroforestry	10.0	$\pm 75\%$
Temperate and Subtropical	All	All	General perennial cropland	TBD	TBD
# Represents a nominal estimate of error, equivalent to two times standard deviation, as a percentage of the mean.					
[Placeholder: Table 5.11 will be updated in the SOD consistent with default values in updated tables 5.1 to 5.4 for perennial crop and relevant tables in chapter 2 (Table 2.4 and Table 2.6) and chapter 11 (Table 11.2) for annual crop.]					

Tier 2

Tier 2 methods should include some country-specific estimates for biomass stocks and removals due to land conversion, and also include estimates of on-site and off-site losses due to burning and decay following land conversion to Cropland. These improvements can take the form of systematic studies of carbon content and emissions and removals associated with land uses and land-use conversions within the country and a re-examination of default assumptions in light of country-specific conditions.

Default parameters for emissions from burning and decay are provided. However, countries are encouraged to develop country-specific coefficients to improve the accuracy of estimates. The *IPCC Guidelines* use a general default of 0.5 for the proportion of biomass burnt on-site for both Forest Land and Grassland conversions. Research studies suggest that the fraction is highly variable and could be as low as 0.2 (Fearnside, 2000; Barbosa and Fearnside, 1996; and Fearnside, 1990). Updated default proportions of biomass burnt on-site are provided in Chapter 4 (Forest Land) for a range of forest vegetation classes. These defaults should be used for transitions from Forest Land to Cropland. For non-forest initial land uses, the default proportion of biomass left on-site and burnt is 0.35. This default takes into consideration research, which suggests the fraction should fall within the range 0.2 to 0.5 (e.g., Fearnside, 2000; Barbosa and Fearnside, 1996; and Fearnside, 1990). It is *good practice* for countries to use 0.35 or another value within this range, provided that the rationale for the choice is documented. There is no default value for the amount of biomass taken off-site and burnt; countries will need to develop a proportion based on national data sources. In

First Order Draft

Chapter 4 (Forest Land), the default proportion of biomass oxidized as a result of burning is 0.9, as originally stated in the *GPG-LULUCF*.

The method for estimating emissions from decay assumes that all biomass decays over a period of 10 years. For reporting purposes countries have two options: 1) report all emissions from decay in one year, recognizing that in reality they occur over a 10 year period, and 2) report all emission from decay on an annual basis, estimating the rate as one tenth of the totals. If countries choose the latter option, they should add a multiplication factor of 0.10 to the equation.

Tier 3

Under Tier 3, all parameters should be country-defined using measurements and monitoring for more accurate values rather than the defaults. Process based models and decay functions can also be used.

5.3.1.3 CHOICE OF ACTIVITY DATA

This section provides an elaboration clarifying the activity data required for carbon gain estimation.

All tiers require estimates of land areas converted to Cropland. The same area estimates should be used for both biomass and soil C calculations on *Land Converted to Cropland*. Higher tiers require greater specificity of areas. At a minimum, the area of Forest Land and natural Grassland converted to Cropland should be identified separately for all tiers. This implies at least some knowledge of the land uses prior to conversion. This may also require expert judgment if Approach 1 in Chapter 3 of these guidelines is used for land area identification.

Tier 1

Separate estimates are required of areas converted to Cropland from initial land uses (i.e., Forest Land, Grassland, Settlements, etc.) to final crop land type (i.e., annual or perennial) ($A_{TO OTHERS}$). For example, countries should estimate separately the area of tropical moist forest converted to annual cropland, tropical moist forest converted to perennial cropland, tropical moist Grassland converted to perennial cropland, etc. Although, to allow other pools to equilibrate and for consistency with land area estimation overall, land areas should remain in the conversion category for 20 years (or other period reflecting national circumstances) following conversion. The methodology assumes that area estimates are based on a one-year time frame, which is likely to require estimation on the basis of average rates on land-use conversion, determined by measurements estimates made at longer intervals. If countries do not have these data, partial samples may be extrapolated to the entire land base or historic estimates of conversions may be extrapolated over time based on the judgement of country experts. Under Tier 1 calculations, international statistics such as FAO databases, *IPCC GPG Reports* and other sources, supplemented with sound assumptions, can be used to estimate the area of *Land Converted to Cropland* from each initial land use. For higher tier calculations, country-specific data sources are used to estimate all possible transitions from initial land use to final crop type. For perennial woody cropland, the total area of planted perennial woody crops for the age classes within the maturing/harvesting cycle (up to 20 years) is required to estimate all biomass carbon change (ΔC_G). See section 5.2.1.3 for details.

Tier 2

It is *good practice* for countries to use actual area estimates for all possible transitions from initial land use to final crop type. Full coverage of land areas can be accomplished either through analysis of periodic remotely sensed images of land-use and land cover patterns, through periodic ground-based sampling of land-use patterns, or hybrid inventory systems. If finer resolution country-specific data are partially available, countries are encouraged to use sound assumptions from best available knowledge to extrapolate to the entire land base. Historic estimates of conversions may be extrapolated over time based on the judgment of country experts.

Tier 3

Activity data used in Tier 3 calculations should be a full accounting of all land-use transitions to Cropland and be disaggregated to account for different conditions within a country. Disaggregation can occur along political (county, province, etc.), biome, climate, or on a combination of such parameters. In many cases, countries may have information on multi-year trends in land conversion (from periodic sample-based or remotely sensed inventories of land use and land cover). Periodic land-use change matrix need to be developed giving the initial and final land-use areas at disaggregated level based on remote sensing and field surveys.

5.1.1.2 CALCULATION STEPS FOR TIER 1 AND TIER 2

No Refinement

5.1.1.3 UNCERTAINTY ASSESSMENT

No Refinement

5.3.2 Dead Organic Matter

No Refinement

5.3.3 Soil carbon

No Refinement in the Introduction

Land is typically converted to Cropland from native lands, managed Forest Land and Grassland, but occasionally conversions can occur from Wetlands and seldom Settlements. Regardless of soil type (i.e., mineral or organic), the conversion of land to Cropland will, in most cases, result in a loss of soil C for some years following conversion (Mann, 1986; Armentano and Menges, 1986; Davidson and Ackerman, 1993). Possible exceptions are irrigation of formerly arid lands and conversion of degraded lands to Cropland.

General information and guidance for estimating changes in soil C stocks are provided in Section 2.3.3 of Chapter 2 (including equations), and that section needs to be read before proceeding with a consideration of specific guidelines dealing with cropland soil C stocks. The total change in soil C stocks for *Land Converted to Cropland* is estimated using Equation 2.24 (Chapter 2), which combines the change in soil organic C stocks (SOC stocks) for mineral soils and organic soils; and stock changes associated with soil inorganic C pools (Tier 3 only). This section provides specific guidance for estimating soil organic C stock changes; see Section 2.3.3.1 for discussion on soil inorganic C (no additional guidance is provided in the Cropland section below).

To account for changes in soil C stocks associated with *Land Converted to Cropland*, countries need to have, at a minimum, estimates of the areas of *Land Converted to Cropland* during the inventory time period. If land-use and management data are limited, aggregate data, such as FAO statistics, can be used as a starting point, along with knowledge of country experts of the approximate distribution of land-use types being converted and their associated management. If the previous land uses and conversions are not unknown, SOC stocks changes can still be computed using the methods provided in *Cropland Remaining Cropland*, but the land base area will likely be different for croplands in the current year relative to the initial year in the inventory. It is critical, however, that the total land area across all land-use sectors be equal over the inventory time period (e.g., 7 million ha may be converted from Forest Land and Grassland to Cropland during the inventory time period, meaning that croplands will have an additional 7 Million ha in the last year of the inventory, while grasslands and forests will have a corresponding loss of 7 Million ha in the last year). *Land Converted to Cropland* is stratified according to climate regions and major soil types, which could either be based on default or country-specific classifications. This can be accomplished with overlays of climate and soil maps, coupled with spatially-explicit data on the location of land conversions.

5.3.3.1 CHOICE OF METHOD

This section contains elaboration on methods and new guidance.

Inventories can be developed using a Tier 1, 2 or 3 approach with each successive tier requiring more detail and resources than the previous one. It is also possible that countries will use different tiers to prepare estimates for the separate subcategories of soil C (i.e., soil organic C stocks changes in mineral soils and organic soils; and stock changes associated with soil inorganic C pools). Decision trees are provided for mineral soils (Figure 2.5) and organic soils (Figure 2.6) in Section 2.3.3.1 (Chapter 2) to assist inventory compilers with selection of the appropriate tier for their soil C inventory.

First Order Draft

Mineral soils**Tier 1**

Soil organic C stock changes for mineral soils can be estimated for land-use conversion to Cropland using Equation 2.25 in Chapter 2. For Tier 1, the initial (pre-conversion) soil organic C stock ($SOC_{(0-T)}$) and C stock in the last year of the inventory time period (SOC_0) are computed from the default reference soil organic C stocks (SOC_{REF}) and default stock change factors (F_{LU} , F_{MG} , F_I). Annual rates of stock changes are estimated as the difference in stocks (over time) divided by the time dependence (D) of the Cropland stock change factors (default is 20 years).

Tier 2*Refining Application of Default Equations*

The Tier 2 method for mineral soils also uses Equation 2.25, but involves country-specific or region-specific reference C stocks and/or stock change factors and more disaggregated land-use activity and environmental data.

Three-Pool Steady-State C Model

The three-pool steady-state soil C model is based on estimating C inputs to soils and applying soil carbon pool specific decomposition rates that are modified by given environmental conditions and management practices. This model embraces more of the heterogeneity in soils, by subdividing soil C pool into different rates of turnover, i.e., fast (Active Pool), intermediate (Slow Pool), and long turnover times (Passive Pool).

Tier 3

Tier 3 methods will involve more detailed and country-specific models and/or measurement-based approaches along with highly disaggregated land-use and management data. Tier 3 approaches estimate soil C change from land-use conversions to Cropland, and may employ models, data sets and/or monitoring networks. If possible, it is recommended that Tier 3 methods be integrated with estimates of biomass removal and the post-clearance treatment of plant residues (including woody debris and litter), as variation in the removal and treatment of residues (e.g., burning, site preparation) will affect C inputs to soil organic matter formation and C losses through decomposition and combustion. It is important that models be evaluated with independent observations from country-specific or region-specific field locations that are representative of the interactions of climate, soil and cropland management on post-conversion change in soil C stocks.

Organic soils

No Refinement. See 2013 Wetlands Supplement.

Biochar C Amendments to Mineral Soils**Tier 1**

This methodology utilizes a top-down approach in which the total amount of biochar generated and added to mineral soil is used to estimate the change in soil organic C stocks. Use Equation 2.27 to estimate the change in C stock from biochar amendments in Chapter 2, Section 2.3.3.1, Volume IV.

Tier 2

Tier 2 methods use the same definitions and equations as Tier 1, but with country-specific factors. See Section 2.3.3.1, Chapter 2, Volume IV for more information.

Tier 3

Tier 3 methods can be used to account for GHG sources and sinks not captured in Tiers 1 or 2, such as priming, changes to N_2O or CH_4 fluxes from soils, and changes to net primary production. More information on Tier 3 methods is provided in Section 2.3.3.1 of Chapter 2, Volume IV.

5.3.3.2 CHOICE OF STOCK CHANGE AND EMISSION FACTORS

This section contains elaboration on methods and new guidance.

Mineral soils**Tier 1**

For native unmanaged land, as well as for managed forest lands, settlements and nominally managed grasslands with low disturbance regimes, soil C stocks are assumed equal to the reference values (i.e., land-use, disturbance (forests only), management and input factors equal 1), while it will be necessary to apply the appropriate stock change factors

to represent previous land-use systems that are not the reference condition, such as improved and degraded grasslands. It will also be necessary to apply the appropriate stock change factor to represent input and management effects on soil C storage in the new cropland system. Default reference C stocks are found in Table 2.3 (Chapter 2). See the appropriate land-use chapter for default stock change factors.

In the case of transient land-use conversions to Cropland, the stock change factors are given in Table 5.12, and depend on the length of the fallow (vegetation recovery) cycle in a shifting cultivation system, representing an average soil C stock over the crop-fallow cycle. Mature fallow denotes situations where the non-cropland vegetation (e.g., forests) recovers to a mature or near mature state prior to being cleared again for cropland use, whereas in shortened fallow, vegetation recovery is not attained prior to re-clearing. If land already in shifting-cultivation is converted to permanent Cropland (or other land uses), the stock change factors representing shifting cultivation would provide the 'initial' C stocks ($SOC_{(0-T)}$) in the calculations using Equation 2.25 (Chapter 2).

TABLE 5. 12 SOIL STOCK CHANGE FACTORS (FLU, FMG, FI) FOR LAND-USE CONVERSIONS TO CROPLAND					
Factor value type	Level	Climate regime	IPCC default	Error #	Definition
Land use	Native forest or grassland (non-degraded)	All	1	NA	Represents native or long-term, non-degraded and sustainably managed forest and grasslands.
		Tropical	1	NA	
Land use	Shifting cultivation – Shortened fallow	Tropical	0.64	± 50%	Permanent shifting cultivation, where tropical forest or woodland is cleared for planting of annual crops for a short time (e.g., 3-5 yr) period and then abandoned to regrowth.
	Shifting cultivation – Mature fallow	Tropical	0.8	± 50%	
Land-use, Management, & Input	Managed forest	(default value is 1)			
Land-use, Management, & Input	Managed grassland	(See default values in Table 6.2)			
Land-use, Management, & Input	Cropland	(See default values in Table 5.5)			
# Represents a nominal estimate of error, equivalent to two times standard deviation, as a percentage of the mean. NA denotes ‘Not Applicable’, where factor values constitute defined reference values.					

Tier 2

Refining Application of Default Equations

Estimation of country-specific stock change factors is probably the most important development associated with the Tier 2 approach. Differences in soil organic C stocks among land uses are computed relative to a reference condition, using land-use factors (F_{LU}). Input factors (F_i) and management factors (F_{MG}) are then used to further refine the C stocks of the new cropland system. Additional guidance on how to derive these stock change factors is given in *Croplands Remaining Croplands*, Section 5.2.3.2. See the appropriate chapter for specific information regarding the derivation of stock change factors for other land-use categories (Forest Land in Section 4.2.3.2, Grassland in 6.2.3.2, Settlements in 8.2.3.2, and Other Land in 9.3.3.2).

Reference C stocks can also be derived from country-specific data in a Tier 2 approach. However, reference values should be consistent across the land uses (i.e., Forest Land, Cropland, Grassland, Settlements, Other Land), and thus must be coordinated among the various teams conducting soil C inventories for AFOLU.

The depth for evaluating soil C stock changes can also be extended with the Tier 2 method. This will require extending the depth of the reference C stocks (SOC_{REF}) and stock change factors for all land uses (i.e., F_{LU} , F_i , and F_{MG}) to ensure consistency. Variable depths between reference stocks and stock change factors are likely to introduce biases into the inventory estimates that are computed using Equation 2.25.

First Order Draft

For the case of land use change to a system that is increasing in C, such as Croplands converted to Grasslands, a Tier 2 method may be a more accurate way to estimate the increase of soil C stocks to native levels. The Tier 1 method may overestimate soil C stock increases on an annual basis (e.g., Villarino et al., 2014).

Furthermore, inventories may be improved by estimating carbon stocks on a mass equivalency basis when deriving country-specific factors for F_{LU} . This is because the soil weight in a certain soil depth changes with the various operations associated with land use change, for example uprooting, land leveling, and rain compaction due to the disappearance of the cover of tree canopy. In addition, cropland soils usually tend to have relatively higher density than the soils in forest land and possibly grasslands or wetlands. Settlement management may also impact the soil bulk density. In such case, the comparison of the soil carbon stocks between the cropland, settlement, grassland, wetland, or forest land within the same depth is not appropriate. It is more robust to compare the carbon stock on an equivalent mass basis, with the stock change calculated on the same weight soil. This method is also recommended when the bulk density between cropland, grassland, wetland, settlement and forest land is remarkably different even if the site is within close proximity. However, it is important to realize that all measurements and associated stock change factors across all land uses must be on an equivalent mass basis if this method is applied. This will be challenging and possibly not even practical unless done comprehensively for all land uses.

Three-Pool Steady-State C Model

Default parameters are provided for the three-pool steady-state C pool equations (Chapter 2, Section 2.3.3.1, Table 2.6), but parameters may be revised if experimental data are available to test the model. Lignin and nitrogen contents are also needed for the C input data (See Section 5.2.3.2 for crop data, and Section 6.2.3.2 for grass data).

Tier 3

Constant stock change rate factors *per se* are less likely to be estimated in favor of variable rates that more accurately capture land-use and management effects. See Chapter 2, Section 2.3.3.1 for further discussion.

Organic soils

No Refinement. See 2013 Wetlands Supplement.

Biochar C Amendments to Mineral Soils

Tier 1

Default emission factors are provided in Chapter 2, Section 2.3.3.1, Volume IV.

Tier 2

Tier 2 emission factors may be further disaggregated relative to the default factors based on variation in environmental conditions, such as the climate and soil types, in addition to variation associated with the biochar production methods. See Section 2.3.3.1, Chapter 2, Volume IV for more information.

Tier 3

Tier 3 methods are country-specific and may involve empirical or process-based models to account for a broader set of impacts of biochar amendments. More information on Tier 3 methods is provided in Section 2.3.3.1, Chapter 2, Volume IV.

5.3.3.3 CHOICE OF ACTIVITY DATA

This section contains elaboration on methods and new guidance.

Mineral soils

Tier 1 and Tier 2 - Refinement of Default Equations

For purposes of estimating soil carbon stock change, area estimates of *Land Converted to Cropland* should be stratified according to major climate regions and soil types. This can be based on overlays with suitable climate and soil maps and spatially-explicit data of the location of land conversions. Detailed descriptions of the default climate and soil classification schemes are provided in Chapter 3, Annex 3A.5. Specific information is provided in each of the land-use chapters regarding treatment of land-use/management activity data (Forest Land in Section 4.2.3.3, Cropland in 5.2.3.3, Grassland in 6.2.3.3, Settlements in 8.2.3.3, and Other Land in 9.3.3.3).

One critical issue in evaluating the impact of *Land Converted to Cropland* on soil organic C stocks is the type of land-use and management activity data. Activity data gathered using Approach 2 or 3 (see Chapter 3 for discussion about

approaches) provide the underlying basis for determining the previous land use for *Land Converted to Cropland*. In contrast, aggregate data (Approach 1, Chapter 3) only provide the total amount of area in each land at the beginning and end of the inventory period (e.g., 1985 and 2005). Approach 1 data are not sufficient to determine specific transitions. In this case all Cropland will be reported in the *Cropland Remaining Cropland* category and in effect transitions become step changes across the landscape. This makes it particularly important to achieve coordination among all land sectors to ensure that the total land base is remaining constant over time, given that some land area will be lost and gained within individual sectors during each inventory year due to land-use change.

Tier 2 – Three-Pool Steady-State C Model

This method requires soil C input data based on the amount of biomass that is converted to dead organic matter annually. This rate will vary depending on plant production, management activity, natural disturbances, and other environmental variables. Removals or reductions in dead organic matter are subtracted from the C input, which could occur with practices such as collection of coarse woody debris or crop residues, burning of grasslands, field burning of agricultural residues, livestock grazing, and other practices. Disturbance events, such as pest outbreaks, may increase the dead organic matter, and therefore the C input to soils. It is good practice to use country-specific methods for estimating C input to soils, but defaults approaches are provided for cropland (Section 5.2.3.3) and grassland (Section 6.2.3.3). Tillage management data are also required (proportion of full tillage, reduced tillage and no-till), and irrigation data for any lands that are provided supplement water (proportion of land).

Additional ancillary data for this method include monthly weather data and soil texture (i.e., sand content), which are available from global weather and soils datasets if country-specific data are not available, such as the CRU climate dataset (<https://crudata.uea.ac.uk/cru/data/hrg/>), and the Harmonized World Soil Database (<http://webarchive.iiasa.ac.at/Research/LUC/External-World-soil-database/HTML/>), respectively.

Tier 3

For application of dynamic models and/or a direct measurement-based inventory in Tier 3, similar or more detailed data on the combinations of climate, soil, topographic and management data are needed, relative to Tier 1 or 2 methods, but the exact requirements will be dependent on the model or measurement design.

Organic soils

No Refinement. See 2013 Wetlands Supplement.

Biochar C Amendments to Mineral Soils

Tier 1

The activity data required for the Tier 1 method includes the total quantities of biochar distributed for amendment to mineral soils. These data must be disaggregated by production type, where production type is defined as a process utilizing a specific feedstock type, and a specific conversion process (gasification, or high-, medium-, or low-temperature pyrolysis; Tables 2.4 and 2.5). In case data on the temperature of pyrolysis are unavailable, default factors for uncontrolled or unspecified pyrolysis temperatures are provided in Section 2.3.3.1 of Chapter 2, Volume IV. Changes in soil C associated with biochar amendments is considered to occur where it is incorporated into soil. However, due to the distributed nature of the land sector in which this can take place, inventory compilers may not have access to data on when or where biochar C amendments occur. Therefore, for the purposes of Tier 1 method, inventory compilers can rely on centralized records from biochar producers, importers, exporters or distributors, recording the quantity of biochar that has been provided to the land use sector for use as a soil amendment in the country. Note that exported biochar is not included in the total amount of biochar amended to soils in the country. Inventory compilers may further disaggregate amendments by land use if the data are available.

Tier 2

Tier 2 methods have the same activity data requirements as Tier 1 (quantities of biochar distributed for incorporation into mineral soils, disaggregated by production type). Additionally, activity data on the amount of biochar amendments may be disaggregated by climate zones and/or soil types if country-specific factors are disaggregated by these environmental variables. The additional climate and soil activity data may be obtained with a survey of biochar distributors and land managers.

Tier 3

The additional activity data required to support a Tier 3 method will depend on which processes are represented and environmental variables that are required as input to the model. Priming, soil GHG emissions, and plant production responses to biochar all vary with biochar type, climate, and soil type. Furthermore, soil GHG emissions and plant production responses also vary with crop type and management. Therefore, Tier 3 methods may require environmental data on climate zones, soil types, crop types and crop management systems (such as nitrogen fertilizer application rates, and whether soils are flooded for paddy rice production), in addition to the amount of biochar amendments in each of the individual combinations of strata for the environmental variables. More detailed activity data specifying the process conditions for biochar production or the physical and chemical characteristics of the biochar may also be required (such as surface area, cation exchange capacity, pH, and ash content).

5.3.3.4 CALCULATION STEPS FOR TIER 1

This section will be updated for the second order draft based on new factors, and also will provide new guidance.

Mineral soils

The steps for estimating SOC_0 and $SOC_{(0-T)}$ and net soil C stock change per ha of *Land Converted to Cropland* on mineral soils are as follows:

Step 1: Organize data into inventory time periods based on the years in which activity data were collected (e.g., 1990 to 1995, 1995 to 2000, etc.)

Step 2: Determine the amount of *Land Converted to Cropland* by mineral soil types and climate regions in the country at the beginning of the first inventory time period. The first year of the inventory time period will depend on the time step of the activity data (0-T; e.g., 5, 10 or 20 years ago).

Step 3: For Grassland converted to Cropland, classify previous grasslands into the appropriate management system using Figure 6.1. No classification is needed for other land uses at the Tier 1 level.

Step 4: Assign native reference C stock values (SOC_{REF}) from Table 2.3 based on climate and soil type.

Step 5: Assign a land-use factor (F_{LU}), management factor (F_{MG}) and C input levels (F_I) to each grassland based on the management classification (Step 2). Values for F_{LU} , F_{MG} and F_I are given in Table 6.2 for grasslands. Values are assumed to be 1 for all other land uses.

Step 6: Multiply the factors (F_{LU} , F_{MG} , F_I) by the reference soil C stock to estimate an ‘initial’ soil organic C stock ($SOC_{(0-T)}$) for the inventory time period.

Step 7: Estimate the final soil organic C stock (SOC_0) by repeating Steps 1 to 5 using the same native reference C stock (SOC_{REF}), but with land-use, management and input factors that represent conditions for the cropland in the last (year 0) inventory year.

Step 8: Estimate the average annual change in soil organic C stocks for land converted to Cropland ($\Delta C_{Mineral}$) by subtracting the ‘initial’ soil organic C stock ($SOC_{(0-T)}$) from the final soil organic C stock (SOC_0), and then dividing by the time dependence of the stock change factors (i.e., 20 years using the default factors). Note: if an inventory time period is greater than 20 years, then divide by the difference in the initial and final year of the time period.

Step 9: Repeat Steps 2 to 8 if there are additional inventory time periods (e.g., 1990 to 2000, 2001 to 2010, etc.). Note that *Land Converted to Cropland* will retain that designation for 20 years. Therefore, inventory time periods that are less than 20 years may need to refer to the previous inventory time period to evaluate if a parcel of land is considered *Land Converted to Cropland* or *Cropland Remaining Cropland*.

A numerical example is given below for Forest Land converted to Cropland on mineral soils, using Equation 2.25 and default reference C stocks (Table 2.3) and stock change factors (Table 5.6).

Example: For a forest on volcanic soil in a tropical moist environment: $SOC_{Ref} = 70 \text{ tonnes C ha}^{-1}$. For all forest soils (and for native grasslands) default values for stock change factors (F_{LU} , F_{MG} , F_I) are all 1; thus $SOC_{(0-T)}$ is $70 \text{ tonnes C ha}^{-1}$. If the land is converted into annual cropland, with intensive tillage and low residue C inputs then $SOC_0 = 70 \text{ tonnes C ha}^{-1} \bullet 0.48 \bullet 1 \bullet 0.92 = 30.9 \text{ tonnes C ha}^{-1}$. Thus the average annual change in soil C stock for the area over the inventory time period is calculated as $(30.9 \text{ tonnes C ha}^{-1} - 70 \text{ tonnes C ha}^{-1}) / 20 \text{ yrs} = -2.0 \text{ tonnes C ha}^{-1} \text{ yr}^{-1}$.

Organic soils

No Refinement. See 2013 Wetlands Supplement.

Biochar C Amendments to Mineral Soils

Will be provided in second order draft after factors are derived.

5.3.3.5 UNCERTAINTY ASSESSMENT

No Refinement

5.3.4 Non-CO₂ greenhouse gas emissions from biomass burning

No Refinement

5.4 COMPLETENESS, TIME SERIES, QA/QC, AND REPORTING

No Refinement

First Order Draft

5.5 METHANE EMISSIONS FROM RICE CULTIVATION

No Refinement in the Introduction.

Anaerobic decomposition of organic material in flooded rice fields produces methane (CH₄), which escapes to the atmosphere primarily by transport through the rice plants (Takai, 1970; Cicerone and Shetter, 1981; Conrad, 1989; Nouchi *et al.*, 1990). The annual amount of CH₄ emitted from a given area of rice is a function of the number and duration of crops grown, water regimes before and during cultivation period, and organic and inorganic soil amendments (Neue and Sass, 1994; Minami, 1995). Soil type, temperature, and rice cultivar also affect CH₄ emissions.

5.5.1 Choice of method

No Refinement

The basic equation to estimate CH₄ emissions from rice cultivation is shown in Equation 5.2. CH₄ emissions are estimated by multiplying daily emission factors by cultivation period³ of rice and annual harvested areas⁴. In its most simple form, this equation is implemented using national activity data (i.e., national average cultivation period of rice and area harvested) and a single emission factor. However, the natural conditions and agricultural management of rice production may be highly variable within a country. It is *good practice* to account for this variability by disaggregating national total harvested area into sub-units (e.g., harvested areas under different water regimes). Harvested area for each sub-unit is multiplied by the respective cultivation period and emission factor that is representative of the conditions that define the sub-unit (Sass, 2002). With this disaggregated approach, total annual emissions are equal to the sum of emissions from each sub-unit of harvested area.

Equation 5. 2 CH₄ emissions from rice cultivation

$$CH_4 \text{ Rice} = \sum_{i,j,k} (EF_{i,j,k} \cdot t_{i,j,k} \cdot A_{i,j,k} \cdot 10^{-6})$$

Where:

CH₄ Rice = annual methane emissions from rice cultivation, Gg CH₄ yr⁻¹

EF_{ijk} = a daily emission factor for *i, j*, and *k* conditions, kg CH₄ ha⁻¹ day⁻¹

t_{ijk} = cultivation period of rice for *i, j*, and *k* conditions, day

A_{ijk} = annual harvested area of rice for *i, j*, and *k* conditions, ha yr⁻¹

i, j, and *k* = represent different ecosystems, water regimes, type and amount of organic amendments, and other conditions under which CH₄ emissions from rice may vary

The different conditions that should be considered include rice ecosystem type, flooding pattern before and during cultivation period, and type and amount of organic amendments. Other conditions such as soil type, and rice cultivar can be considered for the disaggregation if country-specific information about the relationship between these conditions and CH₄ emissions are available. The rice ecosystem types and water regimes during cultivation period are listed in Table 5.14. If the national rice production can be subdivided into climatic zones with different production systems (e.g., flooding patterns), Equation 5.2 should be applied to each region separately. The same applies if rice statistics or expert judgments are available to distinguish management practices or other factors along administrative

³ In the case of a ratoon crop, 'cultivation period' should be extended by the respective number of days.

⁴ In case of multiple cropping during the same year, 'harvested area' is equal to the sum of the area cultivated for each cropping.

units (district or province). In addition, if more than one crop is harvested during a given year, emissions should be estimated for each cropping season taking into account possible differences in cultivation practice (e.g., use of organic amendments, flooding pattern before and during the cultivation period).

The decision tree in Figure 5.2 guides inventory agencies through the process of applying the *good practice* IPCC approach. Implicit in this decision tree is a hierarchy of disaggregation in implementing the IPCC method. Within this hierarchy, the level of disaggregation utilised by an inventory agency will depend upon the availability of activity and emission factor data, as well as the importance of rice as a contributor to its national greenhouse gas emissions. The specific steps and variables in this decision tree, and the logic behind it, are discussed in the text that follows the decision tree.

Tier 1

Tier 1 applies to countries in which either CH₄ emissions from rice cultivation are not a key category or country-specific emission factors do not exist. The disaggregation of the annual harvest area of rice needs to be done for at least three baseline water regimes including irrigated, rainfed, and upland. It is encouraged to incorporate as many of the conditions (*i, j, k*, etc.) that influence CH₄ emissions (summarized in Box 5.2) as possible. Emissions for each sub-unit are adjusted by multiplying a baseline default emission factor (for field with no pre-season flooding for less than 180 days prior to rice cultivation and continuously flooded fields without organic amendments, EF_c) by various scaling factors as shown in Equation 5.3. The calculations are carried out for each water regime and organic amendment separately as shown in Equation 5.3.

Box 5. 1 Text to be provided in SOD

Box 5. 2 Conditions influencing CH₄ emissions from rice cultivation

The following rice cultivation characteristics should be considered in calculating CH₄ emissions as well as in developing emission factors:

Regional differences in rice cropping practices: If the country is large and has distinct agricultural regions with different climate and/or production systems (e.g., flooding patterns), a separate set of calculations should be performed for each region.

Multiple crops: If more than one crop is harvested on a given area of land during the year, and the growing conditions vary among cropping seasons, calculations should be performed for each season.

Water regime: In the context of this chapter, water regime is defined as a combination of (i) ecosystem type and (ii) flooding pattern.

Ecosystem type: At a minimum, separate calculations should be undertaken for each rice ecosystem (i.e., irrigated, rainfed, and deep water rice production).


Flooding pattern: Flooding pattern of rice fields has a significant effect on CH₄ emissions (Sass et al., 1992; Yagi et al., 1996; Wassmann et al., 2000). Rice ecosystems can further be distinguished into continuously and intermittently flooded (irrigated rice), and regular rainfed, drought prone, and deep water (rainfed), according to the flooding patterns during the cultivation period. Also, flooding pattern before cultivation period should be considered (Yagi et al., 1998; Cai et al., 2000; 2003a; Fitzgerald et al., 2000).

Organic amendments to soils: Organic material incorporated into rice soils increases CH₄ emissions (Schütz et al., 1989; Yagi and Minami, 1990; Sass et al., 1991). The impact of organic amendments on CH₄ emissions depends on type and amount of the applied material which can be described by a dose response curve (Denier van der Gon and Neue, 1995; Yan et al., 2005). Organic material incorporated into the soil can either be of endogenous (straw, green manure, etc.) or exogenous origin (compost, farmyard manure, etc.). Calculations of emissions should consider the effect of organic amendments.

Other conditions: It is known that other factors, such as soil type (Sass et al., 1994; Wassmann et al., 1998; Huang et al., 2002), rice cultivar (Watanabe and Kimura, 1998; Wassmann and Aulakh, 2000), sulphate containing amendments (Lindau et al., 1993; Denier van der Gon and Neue, 2002), etc., can significantly influence CH₄ emissions. Inventory agencies are encouraged to make every effort to consider these conditions if country-specific information about the relationship between these conditions and CH₄ emissions is available.

1169

1170

Figure 5.2 Decision tree for CH₄ emissions from rice production

Note:

1: See Volume 1 Chapter 4, "Methodological Choice and Identification of Key Categories" (noting Section 4.1.2 on limited resources), for discussion of *key categories* and use of decision trees.

First Order Draft

Equation 5. 3 Adjusted daily emission factor

$$EF_i = EF_c \bullet SF_w \bullet SF_p \bullet SF_o \bullet SF_{s,r}$$

Where:

EF_i = adjusted daily emission factor for a particular harvested area

EF_c = baseline emission factor for continuously flooded fields without organic amendments (from Table 5.13)

SF_w = scaling factor to account for the differences in water regime during the cultivation period (from Table 5.14)

SF_p = scaling factor to account for the differences in water regime in the pre-season before the cultivation period (from Table 5.15)

SF_o = scaling factor should vary for both type and amount of organic amendment applied (from Equation 5.4 and Table 5.16)

$SF_{s,r}$ = scaling factor for soil type, rice cultivar, etc., if available

Tier 2

Tier 2 applies the same methodological approach as Tier 1, but country-specific emission factors and/or scaling factors should be used. These country-specific factors are needed to reflect the local impact of the conditions (i, j, k , etc.) that influence CH_4 emissions, preferably being developed through collection of field data. As for Tier 1 approach, it is encouraged to implement the method at the most disaggregated level and to incorporate the multitude of conditions (i, j, k , etc.) that influence CH_4 emissions.

Tier 3

Tier 3 includes models and monitoring networks tailored to address national circumstances of rice cultivation, repeated over time, driven by high-resolution activity data and disaggregated at sub-national level. Models can be empirical or mechanistic, but must in either case be validated with independent observations from country or region-specific studies that cover the range of rice cultivation characteristics (Cai *et al.*, 2003b; Li *et al.*, 2004; Huang *et al.*, 2004). Proper documentation of the validity and completeness of the data, assumptions, equations and models used is therefore critical. Tier 3 methodologies may also take into account inter-annual variability triggered by typhoon damage, drought stress, etc. Ideally, the assessment should be based on recent satellite data.

5.5.2 Choice of emission and scaling factors

This section contains updates and new guidance.

Tier 1

A baseline emission factor for non-flooded fields for less than 180 days prior to rice cultivation and continuously flooded during the rice cultivation period without organic amendments (EF_c) is used as a starting point. The IPCC default for EF_c is $1.15 \text{ kg } CH_4 \text{ ha}^{-1} \text{ day}^{-1}$ (with error range of 0.77 – 1.71, Table 5.13), estimated by a statistical analysis of available field measurement data.

Scaling factors are used to adjust the EF_c to account for the various conditions discussed in Box 5.2, which result in adjusted daily emission factors (EF_i) for a particular sub-unit of disaggregated harvested area according to Equation 5.3. The most important scaling factors, namely water regime during and before cultivation period and organic amendments, are represented in Tables 5.14, 5.15, and 5.16, respectively, through default values. Country-specific scaling factors should only be used if they are based on well-researched and documented measurement data. It is encouraged to consider soil type, rice cultivar, and other factors if available.

UPDATED- TABLE 5. 13 DEFAULT CH ₄ BASELINE EMISSION FACTOR ASSUMING NO FLOODING FOR LESS THAN 180 DAYS PRIOR TO RICE CULTIVATION, AND CONTINUOUSLY FLOODED DURING RICE CULTIVATION WITHOUT ORGANIC AMENDMENTS					
CH ₄ emission (kg CH ₄ ha ⁻¹ d ⁻¹)	World		Regional		
	Emission factor	Error range	Region	Emission factor	Error range
	1.15	0.77 - 1.71	East Asia	TBD	TBD
			Southeast Asia	TBD	TBD
			South Asia	TBD	TBD
			Europe	TBD	TBD
			North America	TBD	TBD
			South America	TBD	TBD
Source: Emission Factors and Error Ranges with TBD are to be determined based on statistical model and updated database used in developing the IPCC 2006 Guidelines. Updated values are to be provided in the second order draft (SOD).					

New Guidance

Box 5.3 Good practice guidance for developing baseline emission factors (EF) for methane emission from rice cultivation

The following guidelines provide good practices in performing manual measurement of methane emissions using the closed-chamber technique for continuously flooded rice fields with recommended fertilizer application and no organic amendment. The data can be used to develop country- and region-specific EF_c.

Chamber Design: It is good practice to use lightweight material that is break resistant and inert to reactions with CH₄ (e.g., acrylic and PVC). It may be a rectangular or cylindrical chamber, covering at least two rice hills. The chamber height must be higher than the rice plant. If necessary, use a base with a groove that can be filled with water to ensure a gas-tight closure. The chamber is equipped with a small fan, a thermometer, a vent hole with a stopper, and a gas sampling port (e.g., a flexible tube connected to a valve).

Field set up and Experimental Design: Select a field that is homogeneous with respect to soil properties, and use an appropriate experimental design with at least 3 replications.

Sampling Strategies: Sampling can be done 1 or 2 times per day between mid-morning and late morning period, and at least once a week for the whole growing period. More frequent measurements are needed during agricultural management events (e.g., irrigation, drainage, and N fertilization). All treatments would have to be measured at the same time. At each sampling time, it is good practice to obtain 3 to 4 gas samples within 30 minutes after closure of the chamber.

For gas sampling, the use of a syringe or a pump is recommended depending on the required sample volume. Plastic or glass containers can be used for collecting samples, and should be transferred to a laboratory and analyzed within the allowable storage period.

Gas Analysis: Use gas chromatograph (GC) equipped with a flame ionization detector (FID) for analysis. Calibrate the GC before every analysis, using certified standard gases.

Data Processing: Use linear regression of the gas concentration inside the chamber against time to calculate the hourly flux. Identify the reasons of non-linearity (if exists) for the validation and correction of calculated flux. Use trapezoidal integration to calculate cumulative gas emissions from the hourly flux data.

Deriving Emission Factor:

A simple average and standard deviation could be used to derive the country EF with data from several sites in different regions, or environmental conditions that create variation in methane emissions in the continuously flooded rice systems. The compiler could also derive disaggregated EFs using regressions models to predict the values for different regions and/or environmental conditions.

For more details refer to Minamikawa et al. 2015, and Sanders and Wassmann, 2014.

1215

1216 **Water regime during the cultivation period (SF_w):** Table 5.14 provides default scaling factors and error ranges
 1217 reflecting different water regimes. The aggregated case refers to a situation when activity data are only available for
 1218 rice ecosystem types, but not for flooding patterns (see Box 5.2). In the disaggregated case, flooding patterns can be
 1219 distinguished in the form of three subcategories as shown in Table 5.14. It is *good practice* to collect more
 1220 disaggregated activity data and apply disaggregated case SF_w whenever possible.




1221

UPDATED TABLE 5. 14 DEFAULT CH4 EMISSION SCALING FACTORS FOR WATER REGIMES DURING THE CULTIVATION PERIOD RELATIVE TO CONTINUOUSLY FLOODED FIELDS					
Water regime		Aggregated case		Disaggregated case	
		Scaling factor (SF _w)	Error range	Scaling factor (SF _w)	Error range
Upland ^a		0	-	0	-
Irrigated ^b	Continuously flooded	0.79	0.70 - 0.89	1	0.72 - 1.28
	Intermittently flooded – single aeration			0.79	0.59 – 1.06
	Intermittently flooded – multiple aeration			0.53	0.40 - 0.70
Rainfed and deep water ^c	Regular rainfed	0.42	0.37 - 0.47	0.55	0.40 - 0.77
	Drought prone			0.18	0.12 - 0.26
	Deep water			0.06	0.03 – 0.12
ND: not determined					
^a Fields are never flooded for a significant period of time.					
^b Fields are flooded for a significant period of time and water regime is fully controlled.					
• Continuously flooded: Fields have standing water throughout the rice growing season and may only dry out for harvest (end-season drainage).					
• Intermittently flooded : Fields have at least one aeration period of more than 3 days during the cropping season.					
- Single aeration: Fields have a single aeration during the cropping season at any growth stage (except for end-season drainage).					
- Multiple aeration: Fields have more than one aeration period during the cropping season except for end-season drainage, including alternate wetting and drying (AWD).					
^c Fields are flooded for a significant period of time and water regime depends solely on precipitation.					
• Regular rainfed: The water level may rise up to 50 cm during the cropping season.					
• Drought prone: Drought periods occur during every cropping season.					
• Deep water rice: Floodwater rises to more than 50 cm for a significant period of time during the cropping season.					
Note: Other rice ecosystem categories, like swamps and inland, saline or tidal wetlands may be discriminated within each sub-category.					
Source: Scaling Factors and Error Ranges are determined based on statistical model and updated database used in developing the IPCC 2006 Guidelines.					

1222 **Water regime before the cultivation period (SF_p):** Table 5.15 provides default scaling factors for water regime before
 1223 the cultivation period which can be used when country-specific data are unavailable. This table distinguishes three
 1224 different water regimes prior to rice cultivation, namely:

- 1225 1. Non-flooded pre-season < 180 days, which often occurs under double cropping of rice;
- 1226 2. Non-flooded pre-season > 180 days, e.g., single rice crop following a dry fallow period; and
- 1227 3. Flooded pre-season in which the minimum flooding interval is set to 30 days; i.e., shorter flooding periods (usually
 1228 done to prepare the soil for ploughing) will not be included in this category.

1229 When activity data for the pre-season water status are not available, aggregated case factors can be used. It is *good*
 1230 *practice* to collect more disaggregated activity data and apply disaggregated case of SF_p. Scaling factors for additional
 1231 water regimes can be applied if country-specific data are available.

UPDATED - TABLE 5. 15 DEFAULT CH ₄ EMISSION SCALING FACTORS FOR WATER REGIMES BEFORE THE CULTIVATION PERIOD				
Water regime prior to rice cultivation (schematic presentation showing flooded periods as shaded)	Aggregated case		Disaggregated case	
	Scaling factor (SF _p)	Error range	Scaling factor (SF _p)	Error range
Non flooded pre-season <180 d 	1.20	0.91 - 1.58	1	0.88 - 1.12
Non flooded pre-season >180 d 			0.92	0.83 - 1.03
Flooded pre-season (>30 d) ^{a,b} 			2.12	1.86 - 2.41
^a Short pre-season flooding periods of less than 30 d are not considered in selection of SF _p ^b For calculation of pre-season emission see below (section on completeness)				
Source: Scaling Factors and Error Ranges are determined based on statistical model and updated database used in developing the IPCC 2006 Guidelines.				

Organic amendments (SF_o): It is *good practice* to develop scaling factors that incorporate information on the type and amount of organic amendment applied (compost, farmyard manure, green manure, and rice straw). On an equal mass basis, more CH₄ is emitted from amendments containing higher amounts of easily decomposable carbon and emissions also increase as more of each organic amendment is applied. Equation 5.4 and Table 5.16 present an approach to vary the scaling factor according to the amount of different types of amendment applied. Rice straw is often incorporated into the soil after harvest. In the case of a long fallow after rice straw incorporation, CH₄ emissions in the ensuing rice-growing season will be less than the case that rice straw is incorporated just before rice transplanting (Fitzgerald *et al.*, 2000). Therefore, the timing of rice straw application was distinguished. An uncertainty range of 0.54-0.64 can be adopted for the exponent 0.59 in Equation 5.4.

Equation 5. 4 Adjusted CH₄ emission scaling factors for organic amendments

$$SF_o = \left(1 + \sum_i ROA_i \bullet CFOA_i \right)^{0.59}$$

Where:

SF_o = scaling factor for both type and amount of organic amendment applied

ROA_i = application rate of organic amendment *i*, in dry weight for straw and fresh weight for others, tonne ha⁻¹

CFOA_i = conversion factor for organic amendment *i* (in terms of its relative effect with respect to straw applied shortly before cultivation) as shown in Table 5.16.

TABLE 5.16 DEFAULT CONVERSION FACTORS FOR DIFFERENT TYPES OF ORGANIC AMENDMENTS

Organic amendment	Conversion factor (CFOA)	Error range
Straw incorporated shortly (<30 days) before cultivation ^a	1	0.96 - 1.04
Straw incorporated long (>30 days) before cultivation ^a	0.09	0.01 - 0.17
Compost	0.17	0.08 - 0.26
Farm yard manure	0.14	0.08 - 0.20
Green manure	0.42	0.37 - 0.47
^a Straw application means that straw is incorporated into the soil, it does not include case that straw just placed on the soil surface, nor that straw was burnt on the field. Source: Conversion Factors and Error Ranges are determined based on statistical model and updated database used in developing the IPCC 2006 Guidelines.		

Soil type (SF_s) and rice cultivar (SF_r): In some countries emission data for different soil types and rice cultivar are available and can be used to derive SF_s and SF_r, respectively. Both experiments and mechanistic knowledge confirm the importance of these factors, but large variations within the available data do not allow one to define reasonably accurate default values. It is anticipated that in the near future simulation models will be capable of producing specific scaling factors for SF_s and SF_r.

Tier 2

Inventory agencies can use country-specific emission factors from field measurements that cover the conditions of rice cultivation in their respective country. It is *good practice* to compile country-specific data bases on available field measurements which supplement the Emission Factor database⁵ by other measurement programs (e.g., national) not yet included in this data base. However, certain standard QA/QC requirements apply to these field measurements (see Section 5.5.5).

In Tier 2, inventory agencies can define the baseline management according to the prevailing conditions found in their respective country and determine country-specific emission factors for such a baseline. Then, inventory agencies can also determine country-specific scaling factors for management practices other than the baseline. In case where country-specific scaling factors are not available, default scaling factors can be used. However, this may require some recalculation of the scaling factors given in Tables 5.14 to 5.16 if the condition is different from the baseline.

Tier 3

Tier 3 approaches do not require choice of emission factors, but are instead based on a thorough understanding of drivers and parameters (see above).

5.5.3 Choice of activity data

This section contains further elaboration on methods.

New Guidance Box 5.4 Example of how to estimate methane emission from rice cultivation using Tier 1 method

First Order Draft

A country in Southeast Asia has rice area of 3 million hectares, with 50% of the area classified as irrigated, 30% rainfed, 15% upland, and 5% deep water. Irrigated areas are planted for 2 growing seasons annually. Rice growing periods are 120 days, except for deep water rice which has 220 days. For irrigated areas, 50% is continuously flooded and 50% is managed with multiple aerations. Irrigated areas are flooded pre-season >30 days, and 2 tonnes/ha of straw residues are incorporated shortly less than 30 days before cultivation.

Calculations:

Equations 5.2 and 5.3, of the 2006 IPCC Guidelines, are used to estimate methane emission from rice cultivation:

$$CH_4 \text{ Rice} = \sum_{i,j,k} (EF_{i,j,k} \cdot t_{i,j,k} \cdot A_{i,j,k} \cdot 10^{-6})$$

$$EF_i = EF_c \cdot SF_w \cdot SF_p \cdot SF_o \cdot SF_{s,r}$$

Step 1: Calculate annual harvested area for each rice ecosystem as follows:

Irrigated harvested area = 3,000,000 ha x 0.50 x 2 = 3,000,000 ha

- Irrigated, continuously flooded harvested area = 3,000,000 x 0.5 = 1,500,000 ha

- Irrigated, with multiple aeration harvested area = 3,000,000 x 0.5 = 1,500,000 ha

Rainfed harvested area = 3,000,000 ha x 0.30 = 900,000 ha

Upland harvested area = 3,000,000 ha x 0.15 = 450,000 ha

Deepwater harvested area = 3,000,000 ha x 0.05 = 150,000 ha

Step 2: Assign scaling factors from Table 5.14 to account for water management during the cultivation period)SFw(, for each rice ecosystem:

SFw for Irrigated, Continuously Flooded = 1

SFw for Irrigated, Multiple Aeration = 0.53

SFw for Rainfed = 0.55

SFw for Upland = 0

SFw for Deepwater = 0.06

Step 3: Assign scaling factor from Table 5.15 to account for water management before the cultivation period)SFp(, for Irrigated area:

SFp for Irrigated = 2.12

SFp for Rainfed = 1

SFp for Upland = 1

SFp for Deepwater = 2.12

Step 4: Assign scaling factors from Table 5.14 to account for organic amendment)SFo(in irrigated

Rice areas only:

SFo for Irrigated = $1 + ROA \times CFA^{0.59} = 1 + 2 \times 1^{0.59} = 1.91$

Step 5: Calculate adjusted daily emission factors)EFi(, kg CH₄/ha/day(for each rice ecosystem, using emission factor)EFc(for Southeast Asia in Table 5.13, as follows:

EFi for Irrigated, Continuously Flooded = $EF_c \times 1 \times 2.12 \times 1.91 =$

EFi for Irrigated, Multiple Aeration = $EF_c \times 0.53 \times 2.12 \times 1.91 =$

EFi for Rainfed = $EF_c \times 0.55 \times 1 =$

E_{Fi} for Upland = $E_{Fc} \times 0 \times 1 =$
 E_{Fi} for Deepwater = $E_{Fc} \times 0.06 \times 2.12 =$
 Step 6: Estimate methane emissions)Gg CH₄/yr(for each rice ecosystem as follows:
 CH₄ Emission for Irrigated, Continuously Flooded = $E_{Fi} \times t \text{ 240 days} \times 1,500,000 \text{ ha} \times 10^{-6} =$
 CH₄ Emission for Irrigated, Multiple Aeration = $E_{Fi} \times t \text{ 240 days} \times 1,500,000 \text{ ha} \times 10^{-6} =$
 CH₄ Emission for Rainfed = $E_{Fi} \times 120 \text{ days} \times 900,000 \text{ ha} \times 10^{-6} =$
 CH₄ Emission for Upland = $E_{Fi} \times 120 \text{ days} \times 450,000 \text{ ha} \times 10^{-6} =$
 CH₄ Emission for Deepwater = $E_{Fi} \times 220 \times 150,000 \text{ ha} \times 10^{-6} =$
 Step 7: Sum the emissions from each rice ecosystem to estimate total methane emissions for the country.
 Total CH₄ Emission = _____ Gg CH₄/yr (Note: Final estimate will be provided in the SOD, once E_{Fc} for Southeast Asia is made available).

5.5.4 Uncertainty assessment

No Refinement

References

- IPCC (1997). Revised 1996 IPCC Guidelines for National Greenhouse Inventories. Houghton J.T., Meira Filho L.G., Lim B., Tréanton K., Mamaty I., Bonduki Y., Griggs D.J. Callander B.A. (Eds). Intergovernmental Panel on Climate Change (IPCC), IPCC/OECD/IEA, Paris, France.
- IPCC (2000). Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Penman J., Kruger D., Galbally I., Hiraishi T., Nyenzi B., Emmanuel S., Buendia L., Hoppaus R., Martinsen T., Meijer J., Miwa K., Tanabe K. (Eds). Intergovernmental Panel on Climate Change (IPCC), IPCC/OECD/IEA/IGES, Hayama, Japan.
- IPCC (2003). Good Practice Guidance for Land Use, Land-Use Change and Forestry. Penman J., Gytarsky M., Hiraishi T., Krug, T., Kruger D., Pipatti R., Buendia L., Miwa K., Ngara T., Tanabe K., Wagner F. (Eds). Intergovernmental Panel on Climate Change (IPCC), IPCC/IGES, Hayama, Japan.
- BIOMASS**
- From 2006 GL**
- Albrecht, A. and Kandji, S.T. (2003). Carbon sequestration in tropical agroforestry systems. *Agriculture, Ecosystems and Environment* **99**: 15-27.
- Hairiah, K. and Sitompul, S.M. (2000). Assessment and simulation of above-ground and below-ground carbon dynamics. Report to Asia Pacific Network (APN). Brawijaya University, Faculty of Agriculture, Malang, Indonesia.
- Lasco, R.D. and Suson, P.D. (1999). A *Leucaena Leucocephala* -based indigenous fallow system in central Philippines: the Naalad system. *Intl Tree Crops Journal* **10**: 161-174.
- Lasco, R.D., Lales, J.S., Arnuevo, M.T., Guillermo, I.Q., de Jesus, A.C., Medrano, R., Bajar, O.F. and Mendoza, C.V. (2002). Carbon dioxide (CO₂) storage and sequestration of land cover in the Leyte Geothermal Reservation. *Renewable Energy* **25**: 307-315.

First Order Draft

- Lasco, R.D., Sales, R.F., Estrella, R., Saplaco, S.R., Castillo, A.S.A., Cruz, R.V.O. and Pulhin, F.B. (2001). Carbon stocks assessment of two agroforestry systems in the Makiling Forest Reserve, Philippines. *Philippine Agricultural Scientist* **84**: 401-407.
- Millennium Ecosystems Assessment (2005). *Ecosystems and Human Well-being: A Synthesis*. Island Press, Washington DC. 137pp.
- Moore III, B. (2002). Chapter 2 Challenges of a changing earth. In, *Challenges of a Changing Earth* (W. Steffen, J. Jaeger, D.J. Carson, and C. Bradshaw, eds). Berlin: Springer-Verlag. Pp. 7-17.
- Palm, C.A., Woomer, P.L., Alegre, J., Arevalo, L., Castilla, C., Cordeiro, D.G., Feigl, B., Hairiah, K., Kotto-Same, J., Mendes, A., Maukam, A., Murdiyarso, D., Njomgang, R., Parton, W.J., Ricse, A., Rodrigues, V., Sitompus, S.M. and van Noordwijk, M. (1999). Carbon sequestration and trace gas emissions in slash-and-burn and alternative land-uses in the Humid Tropics. ACB Climate Change Working Group. Final Report Phase II, Nairobi, Kenya.
- Siregar, C.A. and Gintings, Ng. (2000). Research activities related to ground biomass measurement at Forestry Research Development Agency. Paper presented at the Workshop on LUCC and Greenhouse Gas Emissions Biophysical Data. Institut Pertanian Bogor. Indonesia, 16 December 2000.
- Tjitrosemito, S. and Mawardi, I. (2000). 'Terrestrial carbon stock in oil palm plantation', Paper presented at the Science Policy Workshop on Terrestrial Carbon Assessment for Possible Trading under CDM Projects, Bogor, Indonesia 28-29 February 2000.
- Tomich, T.P., van Noordwijk, M., Budidarsono, S., Gillison, A., Kusumanto, T., Murdiyarso, D., Stolle, T. and Fagi, A.M. (1998). Alternative to slash and burn in Indonesia. Summary Report and Synthesis of Phase II. ASB-Indonesia, Report No. 8, ICRAF, Bogor, Indonesia.
- Wasrin, U.R., Rohiani, A., Putera, A.E. and Hidayat, A. (2000). Assessment of above-ground C-stock using remote sensing and GIS technique. Final Report, Seameo Biotrop, Bogor, 28p.
- New References**
- Adachi, M., Ito, A., Ishida, A., Kadir, W. R., Ladpala, P. and Yamagata, Y., 2011, Carbon budget of tropical forests in Southeast Asia and the effects of deforestation: an approach using a process-based model and field measurements, *Biogeosciences*, 8, 2635-2647
- Anil Kumar Yadava, 2010, Biomass Production and Carbon Sequestration in Different Agroforestry Systems in Tarai Region of Central Himalaya, *The Indian Forester*, 135(2), 234-232
- Barrios, E., & Cobo, J. G. (2004). Plant growth, biomass production and nutrient accumulation by slash/mulch agroforestry systems in tropical hillsides of Colombia. *Agroforestry Systems*, 60(3), 255–265.
- Blagodatsky, S., Xu, J., Cadisch, G., (2016), Carbon balance of rubber (*Hevea brasiliensis*) plantations: A review of uncertainties at plot, landscape and production level, *Agriculture Ecosystems & Environment* 221, 8-19
- BMLFUW (Bundesministerium für Land- und Forstwirtschaft, Umwelt und Wasserwirtschaft) 2000: Empfehlungen für die sachgerechte Düngung von Christbaumkulturen. Federal Ministry for Agriculture, Forestry, Environment and Water Management, Wien.
- Buwalda, J.G. and Smith, G.S. 1987. Accumulation and partitioning of dry matter and mineral nutrients in developing kiwifruit vines. *Tree Physiology* 295-307.
- Chalmers D.J. and Van Den Ende, B. 1975. Productivity of peach trees: factors affecting dryweight distribution during tree growth. *Annals of Botany*: 423-432.
- Dogra, A. S., Nautiyal, S., Nautiyal, D. P. (2014). Contribution of *Populus Deltoides* to Farm Economy of Punjab, *The Indian Forester*, 140(8), 758-762.
- Dossa, E. L., Fernandes, E. C. M., Reid, W. S., & Ezui, K. (2008). Above- and belowground biomass, nutrient and carbon stocks contrasting an open-grown and a shaded coffee plantation. *Agroforestry Systems*, 72(2), 103–115.
- Germer, J., Sauerborn, J., (2008). Estimation of the impact of oil palm plantation establishment on greenhouse gas balance, *Environment Development and Sustainability* 10(6), 697-716

- 1394 Goswam, S., Verma, K. S., & Kaushal, R. (2014). Biomass and carbon sequestration in different agroforestry systems
1395 of a Western Himalayan watershed. *Biological Agriculture & Horticulture*, 30(2), 88–96.
- 1396 Goswam, S., Verma, K. S., & Pala, N. A., (2016)., Impact of Input Use on Biomass Attributes and Carbon Mitigation
1397 in Agroforestry Systems of Indian Himalaya, *The Indian Forester*, 142(12), 1214-1219.
- 1398 Gyldenkerne, S., Münier, B., Olesen, J., Olesen, S., Petersen, B. & Christensen, B. (2005).Opgørelse af CO2-
1399 emissioner fra arealanvendelse og ændringer i arealanvendelse. LULUCF (Land Use, Land Use Change and
1400 Forestry). Metodebeskrivelse samt opgørelse for 1990 - 2003., Vol. Arbejdsrapport fra DMU, nr. 213. 2005:
1401 Danmarks Miljøundersøgelser.
- 1402 Hauk, S., Knoke, T., Wittkopf, S. (2013), Economic evaluation of short rotation coppice systems for energy from
1403 biomass—A review, *Renewable and Sustainable Energy Reviews* 29, 435-448
- 1404 Harwood C.E. and Nambiar E.K.S. 2014. Sustainable plantation forestry in South- East Asia. ACIAR Technical
1405 Reports 84.
- 1406 Haynes, R.J. and Goh, K.M., 1980. Variation in the nutrient content of leaves and fruit with season and crown position
1407 for two apple varieties. *Australian Journal of Agricultural Research* 31(4) 739-748
- 1408 Henry, M., Titttonell, P., Manlay, R. J., Bernoux, M., Albrecht, A., & Vanlauwe, B. (2009). Biodiversity, carbon stocks
1409 and sequestration potential in aboveground biomass in smallholder farming systems of western Kenya. *Agriculture,*
1410 *Ecosystems and Environment*, 129(1–3), 238–252.
- 1411 Isaac, M. E., Timmer, V. R., & Quashie-Sam, S. J. (2007). Shade tree effects in an 8-year-old cocoa agroforestry
1412 system: Biomass and nutrient diagnosis of *Theobroma cacao* by vector analysis. *Nutrient Cycling in*
1413 *Agroecosystems*, 78(2), 155–165.
- 1414 Jiménez, C.M. and Diaz, J.B.R. 2003. A statistical model to estimate potential yields in peach before bloom. *Journal*
1415 *of the American Society of Horticultural Science*128: 297-301.
- 1416 Jiménez, C.M. and Diaz, J.B.R. 2004. Statistical model estimates potential yields in ‘Golden Delicious’ and ‘Royal
1417 Gala’ apples before bloom. *Journal of the American Society of Horticultural Science* 129: 20-25.
- 1418 Jose, S., & Bardhan, S. (2012). Agroforestry for biomass production and carbon sequestration: An overview.
1419 *Agroforestry Systems*, 86(2), 105–111.
- 1420 Jose, S. (2009). Agroforestry for ecosystem services and environmental benefits: An overview. *Agroforestry Systems*,
1421 76(1), 1–10.
- 1422 Juhos K. és Tőkei, L. (2012). A hazai szőlőkben és gyümölcsösökben tárolt szén mennyisége. [Carbon stock of
1423 vineyard and orchards in Hungary]. Report based on a project supported by the National Food Chain Safety Office,
1424 Forestry Directorate. Corvinus University of Budapest Budapesti Corvinus Egyetem Kertészettudományi Kar
1425 Talajtan és Vízgazdálkodás Tanszék (in Hungarian).
- 1426 Kandler, G.; Bosch, B. 2013: Methodenentwicklung für die 3. Bundeswaldinventur: Modul 3 Überprüfung und
1427 Neukonzeption einer Biomassefunktion: Abschlussbericht, 69 S., Forstliche Versuchs-und Forschungsanstalt
1428 Baden-Württemberg, Abt. Biometrie und Informatik
- 1429 Kerckhoffs, L.H.J. and Reid, J.B. 2007. Carbon sequestration in the standing biomass of orchard crops in New Zealand.
1430 Report prepared for Horticulture New Zealand Ltd New Zealand Institute for Crop & Food Research Ltd, Hastings,
1431 New Zealand
- 1432 Krasuska, E., Rosenqvist, H., (2012). Economics of energy crops in Poland today and in the future, *Biomass and*
1433 *Bioenergy*, 38, 23-33.
- 1434 Kongsager, R., Napier, J., Mertz, O., (2013), The carbon sequestration potential of tree crop plantations, *Mitigation*
1435 *and Adaptation Strategies for Global Change*,18(8), 1197-1213
- 1436 Kort, J. and Turnock, R. 1999. Carbon reservoir and biomass in Canadian prairie shelterbelts. *Agroforestry Systems*
1437 44: 175-186.
- 1438 Kroodsma, D. A. and Field, C. B., 2006. Carbon sequestration in California agriculture, 1980–2000. *Ecological*
1439 *Applications*, 16(5): 1975-1985

First Order Draft

- 1440 Lakprasadi, H. G. R. K. and Navaratne, C. M., (2012). Estimation of carbon sequestration by cinnamon grown in
1441 WL2a agro ecological zone, Proceedings of 17th International Forestry and Environment Symposium 2012, 17,
- 1442 Lasco, R. D., Evangelista, R. S., & Pulhin, F. B. (2010). Potential of Community-Based Forest Management to
1443 Mitigate Climate Change in the Philippines. *Small-Scale Forestry*, 9(4), 429–443.
- 1444 Lovatt, C.J. 1996 Nitrogen allocation within the "Hass" avocado. *Californian Avocado Society 1996 Yearbook* 80:75-
1445 83.
- 1446 Makumba, W., Akinnifesi, F. K., Janssen, B., & Oenema, O. (2007). Long-term impact of a gliricidia-maize
1447 intercropping system on carbon sequestration in southern Malawi. *Agriculture, Ecosystems and Environment*,
1448 118(1–4), 237–243.
- 1449 McConkey, B., Angers, D., Bentham, M., Boehm, M., Brierley, T., Cerkowniak, D., Liang, B.C., Collas, P., de Gooijer,
1450 H., Desjardins, R., Gameda, S., Grant, B., Huffman, T., Hutchinson, J., Hill, L., Krug, P., Martin, T., Patterson,
1451 G., Rochette, P., Smith, W., VandenBygaart, B., Vergé, X., Worth, D. 2007a. CanAG-MARS Methodology and
1452 Greenhouse Gas Estimates for Agricultural Land in the LULUCF Sector for NIR 2006. Report submitted to the
1453 Greenhouse Gas Division, Environment Canada, by the Research Branch of Agriculture and Agri-Food Canada,
1454 April.
- 1455 Michele Karina Cotta Walter 2012. *Análise do Estoque de Carbono em Sistemas Agrícolas e Florestais em Passo*
1456 *Fundo e Frederico Westphalen, rs*
- 1457 Milne, R. and Brown, T. A. 1997. Carbon in the vegetation and soils of Great Britain. *Journal of Environmental*
1458 *Management*, 49, 413-433.
- 1459 Miria, A., Khan A.B., (2015). Growth and Carbon Storage Study in some Multipurpose Tree Species of Pondicherry
1460 Area, *The Indian Forester*, 141(6), 625-620
- 1461 Mohsin. F., Singh, R.P., Singh, K., 2005, Nutrient Uptake of Poplar Plantation at Various Ages of Growth in Isolated
1462 and Intercropped Stands under Agro-forestry System, *The Indian Forester*, 131(5), 681-693.
- 1463 Mokany, K., R.J. Raison & A.S.P. Rokushkin, 2006: Critical analysis of root:shoot ratios in terrestrial biomes. *Global*
1464 *Change Biology*, 12, 84 -96
- 1465 Montanaro, G., Xiloyannis, C., Nuzzo, V., Dichio, B., 2017. Orchard management, soil organic carbon and ecosystem
1466 services in Mediterranean fruit tree crops, *Scientia Horticulturae*, 217, 92-101
- 1467 Morandé, J.A., Stockert, C.M., Liles, G.C., Williams, J.N. Smart, D.R., and Viers, J.H., 2017, From berries to blocks:
1468 carbon stock quantification of a California vineyard, *Carbon Balance and Management*, 12:5
- 1469 Morgan, K.T., Scholberg, J.M.S., Obreza, T.A., Wheaton, T.A., 2006 Size, Biomass, and Nitrogen Relationships with
1470 Sweet Orange Tree Growth. *J. Amer. Soc. Hort. Sci* 131(1): 146-156
- 1471 Moxley, J., Angelopoulos, N., Buckingham, S., Laidlaw, S., Malcolm. H., Norton, L., Olave, R., Rees, R., Rowe, R.,
1472 Tomlinson, S., Thomson, A., and Topp., K. (2014b) Capturing the effect of Cropland and Grassland Management
1473 on biomass carbon stocks in the UK LULUCF inventory. Unpublished report by CEH for Department of Energy
1474 and Climate Change contract TRN265/09/2011
- 1475 Murphy, T., Jones, G., Vanclay, J., and Glencross, K., 2013. Preliminary carbon sequestration modelling for the
1476 Australian macadamia industry. *Agroforestry Systems*, 87, 689-698.
- 1477 Nendel, C. and Kersebaum, K.C. 2004. A simple model approach to simulate nitrogen dynamics in vineyard soils.
1478 *Ecological Modelling* 177: 1-5.
- 1479 Palmer, J.W., J.N. Wünsche, M. Meland and A. Hann. 2002. Annual dry-matter production by three apple cultivars at
1480 four within-row spacings in New Zealand (2002). *Journal of Horticultural Science Biot.* 77: 712-717.
- 1481 Pessler C, Carbon Storage in Orchards. Master / Diploma Thesis -Institut für Waldökologie (IFE), BOKU-Universität
1482 für Bodenkultur, pp 105, 2012.
- 1483 Popken, S., 2011: Obstanbau, Weinanbau und Weihnachtsbaumkulturen in Deutschland. Zwischenbericht des
1484 Forschungsprojekts „Methodenentwicklung zur Erfassung der Biomasse mehrjährig verholzter Pflanzen außerhalb
1485 von Waldflächen“; Johann Heinrich von Thünen-Institut, Institut für Weltforstwirtschaft

- 1486 Rajput, B.S., Bhardwaj, D.R. & Pala, N.A., (2015). Carbon dioxide mitigation potential and carbon density of different
1487 land use systems along an altitudinal gradient in north-western Himalayas, *Agroforestry Systems*, 89(3), 525-536
- 1488 Rizvi, R. H., Dhyani, S. K., Yadav, R. S., & Singh, R. (2011). Biomass production and carbon stock of poplar
1489 agroforestry systems in Yamunanagar and Saharanpur districts of northwestern India. *Current Science*, 100(5),
1490 736–742.
- 1491 Sanjeev K. C., Naveen Gupta, Ritu, Sudhir Yadav, Rajni Chauhan, (2009). Biomass and Carbon Allocation in
1492 Different Parts of Agroforestry Tree Species, *The Indian Forester*, 134(7), 981-993.
- 1493 Scandellari, F., Caruso, G., Liguori, G., Meggio, F., Palese, A.M., Zanotelli, D., Celano, G., Gucci, R., Inglese, P.,
1494 Pitacco A. Tagliavini M., 2016, A survey of carbon sequestration potential of orchards and vineyards in Italy,
1495 *European Journal of Horticultural Science* 81(2), 106-114
- 1496 Schmitt-Harsh, M., Evans, T. P., Castellanos, E., & Randolph, J. C. (2012). Carbon stocks in coffee agroforests and
1497 mixed dry tropical forests in the western highlands of Guatemala. *Agroforestry Systems*, 86(2), 141–157.
- 1498 Schroeder, P. 1994. Carbon storage benefits of agroforestry systems. *Agroforestry Systems* 27: pp. 89-97.
- 1499 Segura, M., Kanninen, M., & Suárez, D. (2006). Allometric models for estimating aboveground biomass of shade
1500 trees and coffee bushes grown together. *Agroforestry Systems*, 68(2), 143–150.
- 1501 Singh, B, Singh G., (2015). Biomass Production and Carbon Stock in a Silvi-Horti Based Agroforestry System in Arid
1502 Region of Rajasthan, *The Indian Forester*, 141(12), 1237-1243.
- 1503 Singh, K.C. (2005). Relative Growth and Biomass Production of some MPTS under Silvi-pastoral System on a Stony
1504 Rangeland of Arid Zone, *The Indian Forester*, 131(5), 719-723.
- 1505 Singh, N., Lodhiya, L. S., (2016), Fuelwood and Fodder Consumption Pattern an Altitudinal Gradient (1000 - 1200
1506 M) in Mountain Villages of Almora District, *The Indian Forester*, 142(12), 1199-1206
- 1507 Singh, G. (2017), Carbon Sequestration during Restoration of Degraded Hills by Rainwater Harvesting and
1508 Afforestation in Rajasthan, India, *The Indian Forester*, 143(3), 213-222
- 1509 Siregar, C.A. & Gintings, A.N. 2000. Research activities related on ground biomass measurement at forestry research
1510 and development agency. Paper presented at the Workshop on Improving LUC and Greenhouse Gas Emissions
1511 Biophysical Data. 16 December 2000. Institute Pertanian Bogor, Indonesia
- 1512 Somarriba E Cerda R Orozco L Cifuentes M Dávila H et. al. (2013). Carbon stocks and cocoa yields in agroforestry
1513 systems of Central America. *Agriculture, Ecosystems & Environment* 173:46-57.
- 1514 Soto-Pinto, L., Anzueto, M., Mendoza, J., Ferrer, G. J., & Jong, B. (2009). Carbon sequestration through agroforestry
1515 in indigenous communities of Chiapas, Mexico. *Agroforestry Systems*, 78(1), 39–51.
- 1516 Splechtna, B. & Glatzel, G. (2005): Optionen der Bereitstellung von Biomasse aus Wäldern und Energieholzplantagen
1517 für die energetische Nutzung. Berlin-Brandenburgische Akademie der Wissenschaften, Berlin, Materialien Nr. 1
- 1518 Swamy K. R. Vijayakumar P.K., Girish Sankri, Shivanna H., Inamati, S.S., (2012). Carbon Sequestration Potential of
1519 Selected Tree Species Planted in Shelterbelts. *My Forest*, 48(4), 275-280
- 1520 Swamy, S. L., & Puri, S. (2005). Biomass production and C-sequestration of *Gmelina arborea* in plantation and
1521 agroforestry system in India. *Agroforestry Systems*, 64(3), 181–195.
- 1522 Takimoto, A., Nair, P. K. R., & Nair, V. D. (2008). Carbon stock and sequestration potential of traditional and
1523 improved agroforestry systems in the West African Sahel. *Agriculture, Ecosystems and Environment*, 125(1–4),
1524 159–166.
- 1525 Umrao, R. Bijalwan, A., Naugraiya, M. N. (2010), Productivity Status of Ten Year Old Silviculture System in Red
1526 Lateritic Soil of Chhattisgarh Plains, *The Indian Forester*, 136(1), 107-116
- 1527 Villalobos, F.J., Testi, L., Hidalgo, J., Pastor, M., Orgaz, F., 2006. Modelling potential growth and yield of olive (*Olea*
1528 *europaea* L.) canopies. *European Journal of Agronomy*, 24(4): 296-303

First Order Draft

- Wirth, C., Schulze, E. D., Schwalbe, G., Tomczyk, I., Weber, G.-E., Weller, E. 2004: Dynamik der Kohlenstoffvorräte in den Wäldern Thüringens: Abschlussbericht zur 1. Phase des BMBF-Projektes "Modelluntersuchung zur Umsetzung des Kyoto-Protokolls". Mitteilungen der Thüringer Landesanstalt für Wald, Jagd und Fischerei 23.
- Wu, T., Wang, Y., Yu, C., Chiarawipa, R., Zhang, X., Han, Z. Wu, L. (2012) 'Carbon Sequestration by Fruit Trees – Chinese Apple Orchards as an Example', PLoS ONE, 7(6).
- Yashmita-ulman, Avudainayagam, S., (2012). Organic Carbon Storage by Ailanthus excelsa Plantations, The Indian Forester, 138(11), 1041-1051
- Yashmita-ulman, Avudainayagam, S., (2014). Carbon Storage Potential of Eucalyptus Tereticornis Plantations, The Indian Forester, 140(1), 53-58
- Zanotelli, D., Montagnani, L., Manca, G., Scandellari, F., Tagliavini M., 2015. Net ecosystem carbon balance of an apple orchard, European Journal of Agronomy, 63, 97-104
- SOILS**
- Armentano, T.V. and Menges, E.S. (1986). Patterns of change in the carbon balance of organic soil-wetlands of the temperate zone. *Journal of Ecology* **74**: 755-774.
- Augustin, J., Merbach, W., Schmidt, W. and Reining, E. (1996). Effect of changing temperature and water table on trace gas emission from minerotrophic mires. *Journal of Applied Botany-Angewandte Botanik* **70**, 45-51.
- Bruce, J.P., Frome, M., Haites, E., Janzen, H., Lal, R. and Paustian, K. (1999). Carbon sequestration in soils. *Journal of Soil and Water Conservation* **54**:382-389.
- Davidson, E.A. and Ackerman, I.L. (1993). Changes in soil carbon inventories following cultivation of previously untilled soils. *Biogeochemistry*, **20**:161-164.
- Falloon, P. and Smith, P. (2003). Accounting for changes in soil carbon under the Kyoto Protocol: need for improved long-term data sets to reduce uncertainty in model projections. *Soil Use and Management*, **19**, 265-269.
- Freibauer, A. (2003). Biogenic Emissions of Greenhouse Gases from European Agriculture. *European Journal of Agronomy* **19**(2): 135-160.
- Freibauer, A. and Kaltschmitt, M. (eds). (2001). Biogenic greenhouse gas emissions from agriculture in Europe. European Summary Report of the EU concerted action FAIR3-CT96-1877, Biogenic emissions of greenhouse gases caused by arable and animal agriculture, 220 p.
- Glenn, S.M., Hayes, A. and Moore, T.R. (1993). Methane and carbon dioxide fluxes from drained peatland soils, southern Quebec. *Global Biogeochemical Cycles* **7**:247-257
- Kasimir-Klemetsson, A., Klemetsson, L., Berglund, K., Martikainen, P., Silvola, J. and Oenema, O. (1997). Greenhouse gas emissions from farmed organic soils: a review. *Soil Use and Management* **13**:245-250.
- Leifeld, J., Bassin, S. and Fuhrer, J. (2005). Carbon stocks in Swiss agricultural soils predicted by land-use, soil characteristics, and altitude. *Agriculture Ecosystems & Environment* **105**, 255-266.
- Lohila, A., Aurela, M., Tuovinen, J.P. and Laurila, T. (2004). Annual CO₂ exchange of a peat field growing spring barley or perennial forage grass. *Journal of Geophysical Research* **109**, D18116
- Maljanen, M., Martikainen, P.J., Walden, J. and Silvola, J. (2001). CO₂ exchange in an organic field growing barley or grass in eastern Finland. *Global Change Biology* **7**, 679-692.
- Maljanen, M., Komulainen, V.M., Hytonen, J., Martikainen, P. and Laine, J. (2004). Carbon dioxide, nitrous oxide and methane dynamics in boreal organic agricultural soils with different soil characteristics. *Soil Biology and Biochemistry* **36**, 1801-1808.
- Mann, L.K. (1986). Changes in soil carbon storage after cultivation. *Soil Science* **142**:279-288.
- McGill, W. B. (1996). Review and classification of ten soil organic matter models. In: Powlson D.S., Smith P., and Smith J.U. (eds.). Evaluation of Soil Organic Matter Models Using Existing Long-Term Datasets. Springer-Verlag, Heidelberg: pp. 111-132.

- Monte, L., Hakanson, L., Bergstrom, U., Brittain, J. and Heling, R. (1996). Uncertainty analysis and validation of environmental models: the empirically based uncertainty analysis. *Ecological Modelling* **91**:139-152.
- Nusser, S.M. and Goebel, J.J. (1997). The National Resources Inventory: a long-term multi-resource monitoring programme. *Environmental and Ecological Statistics* **4**:181-204.
- Nykänen, H., Alm, J., Lang, K., Silvola, J. and Martikainen, P.J. (1995). Emissions of CH₄, N₂O and CO₂ from a virgin fen and a fen drained for grassland in Finland. *Journal of Biogeography* **22**, 351-357.
- Ogle, S.M., Breidt, F.J., Eve, M.D. and Paustian, K. (2003). Uncertainty in estimating land-use and management impacts on soil organic carbon storage for U.S. agricultural lands between 1982 and 1997. *Global Change Biology* **9**:1521-1542.
- Ogle, S.M., Breidt, F.J. and Paustian, K. (2005). Agricultural management impacts on soil organic carbon storage under moist and dry climatic conditions of temperate and tropical regions. *Biogeochemistry* **72**:87-121.
- Ogle, S.M., Breidt, F.J. and Paustian, K. (2006). Bias and variance in model results associated with spatial scaling of measurements for parameterization in regional assessments. *Global Change Biology* **12**:516-523.
- Paustian, K., Andren, O., Janzen, H.H., Lal, R., Smith, P., Tian, G., Tiessen, H., van, Noordwijk, M. and Woomer, P.L. (1997). Agricultural soils as a sink to mitigate CO₂ emissions. *Soil Use and Management* **13**:230-244.
- Pierce, F. J., Fortin, M.-C. and Staton, M.J. (1994). Periodic plowing effects on soil properties in a no-till farming system. *Soil Science Society of America Journal* **58**:1782-1787.
- Powers, J. S., Read, J. M., Denslow, J. S. and Guzman, S. M. (2004). Estimating soil carbon fluxes following land-cover change: a test of some critical assumptions for a region in Costa Rica. *Global Change Biology* **10**:170-181.
- Smith, J.E. and Heath, L.S. (2001). Identifying influences on model uncertainty: an application using a forest carbon budget model. *Environmental Management* **27**:253-267.
- Smith, P., Powlson, D.S., Smith, J.U. and Elliott, E.T. (eds) (1997). Evaluation and comparison of soil organic matter models. Special Issue, *Geoderma* **81**:1-225.
- Smith, P., Powlson, D.S., Glendining, M.J. and Smith, J.U. (1998) Preliminary estimates of the potential for carbon mitigation in European soils through no-till farming. *Global Change Biology* **4**: 679-685.
- VandenBygaart, A.J., Gregorich, E.G., Angers, D.A., et al. (2004). Uncertainty analysis of soil organic carbon stock change in Canadian cropland from 1991 to 2001. *Global Change Biology* **10**:983-994.
- RICE CULTIVATION**
- Sander, B.O. and Wassmann, R. (2014). Common practices for manual greenhouse gas sampling in rice production: a literature study on sampling modalities of the closed chamber method. *Greenhouse Gas Measurement and Management* **4**:1-13.
- Cai, Z.C., Tsuruta, H. and Minami, K. (2000). Methane emission from rice fields in China: measurements and influencing factors. *Journal of Geophysical Research* **105**D13: 17231-17242.
- Cai, Z.C., Tsuruta, H., Gao, M., Xu, H. and Wei, C.F. (2003a). Options for mitigating methane emission from a permanently flooded rice field. *Global Change Biology* **9**: 37-45.
- Cai, Z.C., Sawamoto, T., Li, C.S., Kang, G.D., Boonjawat, J., Mosier, A. and Wassmann, R. (2003b). Field validation of the DNDC model for greenhouse gas emissions in East Asian cropping systems, *Global Biogeochemical Cycles* **17**4: 1107 doi:10.1029/2003GB002046,2003.
- Cicerone, R.J. and Shetter, J.D. (1981). Sources of atmospheric methane: Measurements in rice paddies and a discussion. *Journal of Geophysical Research* **86**: 7203-7209.
- Conrad, R. (1989). "Control of methane production in terrestrial ecosystems". In: Exchange of Trace Gases between Terrestrial Ecosystems and the Atmosphere, M.O. Andreae and D.S. Schimel (eds.), 39-58.
- Denier van der Gon, H.A.C. and Neue, H.U. (1995). Influence of organic matter incorporation on the methane emission from a wetland rice field. *Global Biogeochemical Cycles* **9**: 11-22.
- Denier van der Gon, H.A.C. and Neue, H.U. (2002). Impact of gypsum application on the methane emission

First Order Draft

- from a wetland rice field. *Global Biogeochemical Cycles* **8**: 127-134.
- Fitzgerald, G.J., Scow, K.M. and Hill, J.E.)2000(. Fallow season straw and water management effects on methane emissions in California rice. *Global Biogeochem. Cycles*, **14**: 767-775.
- Huang, Y., Jiao, Y., Zong, L.G., Zheng, X.H., Sass, R.L. and Fisher, F.M.)2002(. Quantitative dependence of methane emission on soil properties, *Nutrient Cycling in Agroecosystems* **64**)1-2(: 157-167.
- Huang, Y, Zhang, W., Zheng, X.H., Li, J. and Yu, Y.Q.)2004(. Modeling methane emission from rice paddies with various agricultural practices. *Journal of Geophysical Research-Atmospheres* **109**)D8(: Art. No. D08113 APR 29 2004.
- IAEA)1992(. Manual on measurement of methane and nitrous oxide emissions from agriculture. IAEA- TECDOC-674, pp. 91.
- IGAC)1994(. Global measurements standards of methane emissions for irrigated rice cultivation. Sass, R.L. and H.-U. Neue)eds.(IGAC Core Project Office, Cambridge, Mass., USA, 10 pp.
- IPCC)International Panel on Climate Change()1997(. Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories Workbook)Volume 2(. Cambridge University Press, Cambridge.
- IPCC)International Panel on Climate Change()2000(. Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. Cambridge University Press, Cambridge.
- IRRI)1995(. World rice statistics 1993-94, International Rice Research Institute, Los Banos, pp. 260.
- Li, C.S., Mosier, A., Wassmann, R., Cai, Z.C., Zheng, X.H., Huang, Y., Tsuruta, H., Boonjawat, J. and Lantin, R.)2004(. Modeling greenhouse gas emissions from rice-based production systems: Sensitivity analysis and upscaling, *Global Biogeochemical Cycles* **18**, doi: 10.1029/2003GB00204, 2004.
- Lindau, C.W., Bollich, P.K., de Laune, R.D., Mosier, A.R. and Bronson, K.F.)1993(. Methane mitigation in flooded Louisiana rice fields. *Biology and Fertility of Soils* **15**: 174-178.
- Minami, K.)1995(. The effect of nitrogen fertilizer use and other practices on methane emission from flooded rice. *Fertilizer Research* **40**: 71-84.
- Minamikawa, K., Tokida, T., Sudo, S., Padre, A., Yagi, K. (2015). *Guidelines for measuring CH₄ and N₂O emissions from rice paddies by a manually operated closed chamber method*. National Institute for Agro-Environmental Sciences, Tsukuba, Japan.
- Neue, H.U. and Sass, R.)1994(. Trace gas emissions from rice fields. In: Prinn R.G.)ed.(Global Atmospheric-Biospheric Chemistry. Environmental Science Res. 48. Plenum Press, New York, pp. 119-148.
- Nouchi, I., Mariko, S. and Aoki, K.)1990(. Mechanism of methane transport from the rhizosphere to the atmosphere through rice plants. *Plant Physiology* **94**: 59-66.
- Sass, R.)2002(. CH₄ emissions from rice agriculture. In 'Background Papers, IPCC Expert Meetings on Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. IPCC-NGGIP, p. 399-417, available at <http://www.ipcc-nggip.iges.or.jp/>.
- Sass, R.L., Fisher, F.M., Harcombe, P.A. and Turner, F.T.)1991(. Mitigation of methane emission from rice fields: Possible adverse effects of incorporated rice straw. *Global Biogeochemical Cycles*, **5**: 275-287.
- Sass, R.L., Fisher, F.M., Wang, Y.B., Turner, F.T. and Jund, M.F.)1992(. Methane emission from rice fields: The effect of floodwater management. *Global Biogeochemical Cycles* **6**: 249-262
- Sass, R. I., Fisher, F. M., Lewis, S. T., Jund, M. F. and Turner, F. T.)1994(. Methane emissions from rice fields: Effect of soil properties. *Global Biogeochemical Cycles* **2**, 135-140, 1994.
- Schütz, H., Holzapfel-Pschorn, A., Conrad, R., Rennenberg, H. and Seiler, W.)1989(. A 3-year continuous record on the influence of daytime, season, and fertilizer treatment on methane emission rates from an Italian rice paddy. *Journal of Geophysical Research* **94**: 16405-16416.
- Takai, Y.)1970(. The mechanism of methane fermentation in flooded paddy soil. *Soil Science and Plant Nutrition* **16**: 238-244.
- Wassmann, R., and Aulakh, M.S.)2000(. The role of rice plants in regulating mechanisms of methane emissions.

Biology and Fertility of Soils **31**: 20-29.

Wassmann, R., Neue, H.U., Bueno, C., Lantin, R.S., Alberto, M.C.R., Buendia, L.V., Bronson, K., Papen, H. and Rennenberg, H.)1998(. Methane production capacities of different rice soils derived from inherent and exogenous substrates. *Plant and Soil* **203**: 227-237.

Wassmann, R., Buendia, L.V., Lantin, R.S., Makarim, K., Chareonsilp, N. and Rennenberg, H.)2000(. Characterization of methane emissions from rice fields in Asia. II. Differences among irrigated, rainfed, and deepwater rice. *Nutrient Cycling in Agroecosystems* **58**: 107–119.

Watanabe, A. and Kimura, M.)1998(. Factors affecting variation in CH₄ emission from paddy soils grown with different rice cultivars: A pot experiment. *Journal of Geophysical Research* **103**: 18947-18952.

Yagi, K. and Minami, K.)1990(. Effect of organic matter application on methane emission from some Japanese paddy fields. *Soil Science and Plant Nutrition* **36**: 599-610.

Yagi, K., Tsuruta, H., Kanda, K. and Minami, K.)1996(. Effect of water management on methane emission from a Japanese rice paddy field: Automated methane monitoring. *Global Biogeochemical Cycles* **10**: 255-267.

Yagi, K., Minami, K. and Ogawa, Y.)1998(. Effect of water percolation on methane emission from rice paddies: a lysimeter experiment. *Plant and Soil* **198**: 193-200.

Yan, X., Yagi, K., Akiyama, H. and Akimoto, H.)2005(. Statistical analysis of the major variables controlling methane emission from rice fields. *Global Change Biology* **11**, 1131-1141, doi: 10/1111/j.1365-2486.2005.00976.x.r

Annex 5A.1 Estimation of default stock change factors for mineral soil C emissions/removals for cropland

Default stock change factors will be updated in Table 5.5 based on an analysis of a global dataset of experimental results for tillage, input, set-aside, and land use to a 30cm depth. The land-use factor represents the loss of carbon that occurs after 20 years of continuous cultivation. Tillage and input factors represent the effect on C stocks after 20 years following the management change. Set-aside factors represent the effect of temporary removal of cultivated cropland from production and placing it into perennial cover for a period of time that may extend to 20 years.

Semi-parametric mixed effect models are being derived to estimate the new factors. Variables included depth, number of years since the management change, and the type of management change (e.g., reduced tillage vs. no-till). For depth, data are not aggregated to a standardized set of depths but rather each of the original depth increments are used in the analysis (e.g., 0-5 cm, 5-10 cm, and 10-30 cm) as separate observations of stock changes. Similarly, time series data are not aggregated, even though those measurements are taken from the same plots. Consequently, random effects are used to account for the dependencies in times series data and among data points representing different depths from the same study.

Special consideration is given to representing depth increments in order to avoid aggregating data across increments in the original dataset. Consider a field in which the soil has a characteristic $Y(s)$, such as soil organic C, that changes in value with depth, from $s = 0$ decimeters (surface) to $s = 8$ dm. For simplicity, suppose this field is perfectly uniform (no spatial variation) and we have no measurement error, so that a single soil core will be sufficient to describe the soil. We extract a 0-8dm core and measure the average value of Y in the core (averaging over decimeters), to obtain the true average value over $[0,8]$ as $\mu_{[0,8]} = 1/3$. Now suppose instead that we measured the core in increments: 0-1dm, 1-3dm, and 3-8dm. Again, we measure the average value of Y in each core increment, obtaining

$$\mu_{[0,1]} = 0.8802083, \mu_{[1,3]} = 0.5677083, \text{ and } \mu_{[3,8]} = 0.1302083.$$

The simple average of these three values is 0.5260417, much larger than the true value of 1/3. The simple average is wrong. Instead, we should take a weighted average that reflects the differently-sized increments:

$$\frac{(1-0)\mu_{[0,1]} + (3-1)\mu_{[1,3]} + (8-3)\mu_{[3,8]}}{8} = \frac{2.6666667}{8} = \frac{1}{3}.$$

Next, suppose that instead of one soil core, we took two sets of soil cores. The first core uses 0-1dm, 1-3dm, and 3-8dm as before, but the second uses only 0-4dm and 4-8dm. Using the same weighting scheme as before, we can compute weighted averages within each core. Then using the fact that both cores have the same amount of information about 0-8dm, we would take a simple average of the two weighted averages:

$$\frac{(1-0)\mu_{[0,1]} + (3-1)\mu_{[1,3]} + (8-3)\mu_{[3,8]}}{(2)(8)} + \frac{(4-0)\mu_{[0,4]} + (8-4)\mu_{[4,8]}}{(2)(8)} = \frac{2.6666667}{16} + \frac{2.6666667}{16} = \frac{1}{3}.$$

Now suppose we wanted to use those data to estimate the average characteristic in the increment 2-4dm, an increment which is not used in either core. Instead, it is partially contained in the first core's increments $[1, 3]$ and $[3, 8]$, and fully contained in the second core's increment $[0,4]$. We might try using a fraction of each increment, one-half of $[1, 3]$, one-fifth of $[3, 8]$, and one-half of $[0, 4]$:

$$\frac{(1/2)(3-1)\mu_{[1,3]} + (1/5)(8-3)\mu_{[3,8]}}{(2)(2)} + \frac{(1/2)(4-0)\mu_{[0,4]}}{(2)(2)} = \frac{0.6979167}{4} + \frac{1.1666667}{4} = 0.4661458.$$

In fact, the correct value is $\mu_{[2,4]} = 0.3958333$, so we are way off target with this ad hoc approach.

Suppose we took a different approach, by trying to reconstruct the true $Y(s)$. These points are highly suggestive of a quadratic relationship. In fact, unknown to the scientist, the true relationship is exactly quadratic,

$$Y(s) = \frac{1}{64}(s-8)^2 = 1 - 0.25s + 0.015625s^2.$$

We fit the following quadratic model to the data using the midpoints x of the increments.

$$\beta_0 + \beta_1 x + \beta_2 x^2$$

The problem is using the midpoints of the increment ranges and ignoring the fact that these are increment data, not point data. To do this properly, we need to create a custom set of covariates, which are functions of the increment endpoints. These functions come from integrating the underlying quadratic function over the increments. We have centered at the maximum depth, 8dm, for convenience. This means that the true quadratic relationship is

$$\alpha_0 + \alpha_1(s - 8) + \alpha_2(s - 8)^2 = 0 + 0(s - 8) + \frac{1}{64}(s - 8)^2.$$

Because we now have the quadratic model exactly right, we can go back to the problem of estimating $\mu_{[2,4]}$ and we are able to calculate exactly the right answer, 0.3958333. More generally, when combining increment data for statistical analysis, we have to take into account the increment nature of the data or we will have statistically invalid inferences, particularly for regression relationships. In general, we will not know the true relationship and will need to model it flexibly. Any model (like the quadratic in our example) needs to be converted into a set of custom covariates by integrating, just as we did here.

Using this customized approach, we are in the process of estimating management and input factors to a 30 cm depth over a 20 year time period. Variance will be calculated for each of the factor values, and can be used with simple error propagation methods or to construct probability distribution functions with a normal density.

Sources of Data for Land Use Factor

Aborisade KD & Aweto AO (1990) Effects of Exotic Tree Plantations of Teak (*Tectona-Grandis*) and Gmelina (*Gmelina-Arborea*) on a forest soil in South-Western Nigeria. *Soil Use and Management* 6:43-45.

Adachi M, Bekku YS, Rashidah W, Okuda T, Koizumi H (2006) Differences in soil respiration between different tropical ecosystems. *Applied Soil Ecology*, 34, 258-265.

Agbenin JO & JT Goladi (1997) Carbon, nitrogen and phosphorus dynamics under continuous cultivation as influenced by farmyard manure and inorganic fertilizers in the savanna of northern Nigeria. *Agric Ecosyst & Environ* 63:17-24.

Aina PO (1979) Soil Changes Resulting from Long-Term Management-Practices in Western Nigeria. *Soil Science Society of America Journal*, 43, 173-177.

Alcantara FAd, Buurman P, Furtini Neto AE, Curi N, Roscoe R (2004) Conversion of grassy cerrado into riparian forest and its impact on soil organic matter dynamics in an Oxisol from southeast Brazil. *Geoderma*, 123, 305-317.

Allen JC (1985) Soil Response to Forest Clearing in the United States and the Tropics: Geological and Biological factors. *Biotropica* 17:1:15-27.

An DD, He Y, Han AP, Wang J (2003) The effect of different utilization on soil property and microorganism in sub-alpine meadow. *Pratacultural Science* 20(6):1-6

Ashagrie Y, Zech W, Guggenberger G (2005) Transformation of a *Podocarpus falcatus* dominated natural forest into a monoculture *Eucalyptus globulus* plantation at Munesa, Ethiopia: soil organic C, N and S dynamics in primary particle and aggregate-size fractions. *Agriculture Ecosystems & Environment*, 106, 89-98.

Assad ED, Pinto HS, Martins SC, Groppo JD, Salgado PR, Evangelista B, Vasconcellos E, Sano EE, Pavao E, Luna R, Camargo PB, Martinelli LA (2013) Changes in soil carbon stocks in Brazil due to land use: paired site comparisons and a regional pasture soil survey. *Biogeosciences* 10:6141-6160.

Aweto AO (1981) Secondary Succession and Soil Fertility Restoration in Southwestern Nigeria .2. Soil Fertility Restoration. *Journal of Ecology*, 69, 609-614.

Aweto AO (1988) Effects of Shifting Cultivation on a Tropical Rain-Forest Soil in Southwestern Nigeria. *Turrialba*, 38, 19-22.

Aweto AO, Ayuba HK (1993) Effect of Continuous Cultivation with Animal Manuring on a Sub-Sahelian Soil near Maiduguri, North Eastern Nigeria. *Biological Agriculture & Horticulture*, 9, 343-352.

Aweto AO, Ishola MA (1994) The Impact of Cashew (*Anacardium-Occidentale*) on Forest Soil. *Experimental Agriculture*, 30, 337-341.

First Order Draft

- 1799 Ayanaba A, Tuckwell SB, Jenkinson DS (1976) Effects of clearing and cropping on organic reserves and biomass of
1800 tropical forest soils. *Soil Biology & Biochemistry*, 8, 519-525.
- 1801 Banaticla RN, Lasco R (2006) Carbon storage of land cover types in the wetsern margin of Mt. Maliling, Laguna,
1802 Philippines: a case study. *J Nature Studies*, 5, 77-89.
- 1803 Bashkin MA & Binkley D (1998) Changes in soil carbon following afforestation in Hawaii. *Ecology* 79:828-833.
- 1804 Batlle-Bayer L, Batjes NH & Bindraban PS (2010) Changes in Organic Carbon Stocks upon Land Use Conversion in
1805 the Brazilian Cerrado: A Review. *Agriculture, Ecosystems & Environment* 137:47-58.
- 1806 Bautista-Cruz A, del Castillo RF (2005) Soil changes during secondary succession in a tropical montane cloud forest
1807 area. *Soil Science Society of America Journal*, 69, 906-914.
- 1808 Bernhardreversat F (1988) Soil nitrogen mineralization under a Eucalyptus plantation and natural Acacia forest in
1809 Senegal. *Forest Ecology and Management*, 23, 233-244.
- 1810 Berhongaray G, Alvarez R, Paepe JD, Caride C & Cantet R (2013) Land use effects on soil carbon in the Argentine
1811 Pampas. *Geoderma* 192:97-110.
- 1812 Bernardi ACC, Machado PLO, Madari BE, Tavares SR, de Campos DVB & Crisostomo LA (2007) Carbon and
1813 nitrogen stocks of an Arenosol under irrigated fruit orchards in semiarid Brazil. *Science Agriculture (Piracicaba, Brazil)*
1814 64:2:169-175.
- 1815 Bertol I, Santos JCP (1995) Soil use and physical-hidric properties on the plateau of Santa Catarina. *Pesquisa*
1816 *Agropecuaria Brasileira*, 30, 263-267.
- 1817 Berthrong ST, Piñeiro G, Jobbágy EG, Jackson RB (2012) Soil C and N changes with afforestation of grasslands
1818 across gradients of precipitation and plantation age. *Ecological Applications* 22:76-86.
- 1819 Beyer L (1994) Effect of cultivation on physico-chemical, humus-chemical and biotic properties and fertility of two
1820 forest soils. *Agric Ecosyst & Environ* 48:179-188.
- 1821 Binkley D, Kaye J, Barry M & Ryan M (2004) First rotation changes in carbon and nitrogen in a Eucalyptus plantation
1822 in Hawaii. *Soil Sci Soc Am J* 68:1713-1719.
- 1823 Binkley D, Resh SC (1999) Rapid changes in soils following eucalyptus afforestation in Hawaii. *Soil Science Society*
1824 *of America Journal*, 63, 222-225.
- 1825 Bonde TA, Christensen BT, Cerri CC (1992) Dynamics of soil organic matter as reflected by natural C-13 abundance
1826 in particle size fractions of forested and cultivated Oxisols. *Soil Biology & Biochemistry*, 24, 275-277.
- 1827 Bowman RA & Anderson RL (2002) Conservation Reserve Program: Effects on soil organic carbon and preservation
1828 when converting back to cropland in northeastern Colorado. *J Soil & Water Cons* 57:121-126.
- 1829 Brand J, Pfund JL (1998) Site and watershed-level assessment of nutrient dynamics under shifting cultivation in
1830 eastern Madagascar. *Agriculture Ecosystems & Environment*, 71, 169-183.
- 1831 Brown S & Lugo AE (1990) Effects of forest clearing and succession on the carbon and nitrogen content of soils in
1832 Puerto Rico and US Virgin Islands. *Plant & Soil* 124:53-64.
- 1833 Bruun TB, Mertz O & Elberling B (2006) Linking yields of upland rice in shifting cultivation to fallow length and
1834 soil properties. *Agriculture, Ecosystems & Environment* 113:139-149.
- 1835 Burke IC, Elliott ET & Cole CV (1995) Influence of macroclimate, landscape position, and management on soil
1836 organic matter in agroecosystems. *Ecol App* 5:124-131.
- 1837 Burke IC, Lauenroth WK & Coffin DP (1995) Soil organic matter recovery in semiarid grasslands: Implications for
1838 the conservation reserve program. *Ecological Applications* 5:3:793-801.
- 1839 Buschbacher R, Uhl C, Serrao EAS (1988) Abandoned Pastures in Eastern Amazonia .2. Nutrient Stocks in the Soil
1840 and Vegetation. *Journal of Ecology*, 76, 682-699.
- 1841 Buschiazzi DE, Panigatti JL & Unger PW (1998) Tillage effects on soil properties and crop production in the
1842 subhumid and semiarid Argentinean Pampas. *Soil Tillage Res* 49:105-116.

- 1843 Buyanovksy GA, Kucera CL & Wagner GH (1987) Comparative analyses of carbon dynamics in native and cultivated
1844 ecosystems. *Ecology* 68:2023-2031.
- 1845 Cadisch G, Imhof H, Urquiaga S, Boddey RM, Giller KE (1996) Carbon turnover ($\delta^{13}\text{C}$) and nitrogen
1846 mineralization potential of particulate light soil organic matter after rainforest clearing. *Soil Biology & Biochemistry*,
1847 28, 1555-1567.
- 1848 Cai XB, Zhang Y, Shao W (2008) Characteristics of soil fertility in alpine steppes at different degradation grades. *Acta*
1849 *Ecologia Sinica* 28(3):1034-1044
- 1850 Cambardella CA & Elliott ET (1992) Particulate soil organic-matter changes across a grassland cultivation sequence.
1851 *Soil Sci Soc Am J* 56:777-783.
- 1852 Cambardella CA & Elliott ET (1994) Carbon and nitrogen dynamics of soil organic matter fractions from cultivated
1853 grassland soils. *Soil Sci Soc Am J* 58:123-130.
- 1854 Campos AC, Oleschko KL, Etchevers B & Hidalgo CM (2007) Exploring the effects of changes in land use on soil
1855 quality on the eastern slope of Cofre de Perote Volcano (Mexico). *For Ecol & Mgmt* 248:174-182.
- 1856 Cao C, Jiang D, Quan G, Geng L, Cui Z, Luo Y (2004) Soil Physical and Chemical Characters Changes of Caragana
1857 microphylla Plantation for Sand Fixation in Keerqin Sandy Land. *Journal of Soil and Water Conservation* 18(6):108-
1858 102
- 1859 Carvalho JLN, Cerri CEP, Feigl BJ, Piccolo MC, Godinho VP & Cerri CC (2009) Carbon sequestration in agricultural
1860 soils in the Cerrado region of the Brazilian Amazon. *Soil and Tillage Research* 103:342-349.
- 1861 Carvalho JLN, Cerri CEP, Feigl BJ, Piccolo MdC, Godinho VdP, Herpin U, Cerri CC (2009) Conversion of cerrado
1862 into agricultural land in the south-western Amazon: carbon stocks and soil fertility. *Scientia Agricola*, 66, 233-241.
- 1863 Cerri CEP, Coleman K, Jenkinson DS, Bernoux M, Victoria R, Cerri CC (2003) Modeling soil carbon from forest and
1864 pasture ecosystems of Amazon, Brazil. *Soil Science Society of America Journal*, 67, 1879-1887.
- 1865 Cerri CEP, Easter M, Paustian K, et al. (2007) Predicted soil organic carbon stocks and changes in the Brazilian
1866 Amazon between 2000 and 2030. *Agriculture Ecosystems & Environment*, 122, 58-72.
- 1867 Cerri CC, Volkoff B & Andreux F (1991) Nature and behaviour of organic matter in soils under natural forest, and
1868 after deforestation, burning and cultivation, near Manaus. *For Ecol Mgmt* 38:247-257.
- 1869 Chan KY (1997) Consequences of changes in particulate organic carbon in vertisols under pasture and cropping. *Soil*
1870 *Science Society of America Journal*, 61, 1376-1382.
- 1871 Chandran P, Ray SK, Durge SL, Raja P, Nimkar AM, Bhattacharyya T & Pal DK (2009) Scope of horticultural land-
1872 use system in enhancing carbon sequestration in ferruginous soils of the semi-arid tropics. *Current Science* 97:1039-
1873 1046.
- 1874 Chen L, Gong J, Fu B, Huang Z, Huang Y, Gui L (2007) Effects of land use conversion on soil organic carbon
1875 sequestration in the loess hilly area, loess Plateau of China. *Ecological Research* 22:641-648.
- 1876 Chen Y (2006) Study of Dynamic Soil Characteristics under Artificially Planted Caragana-Pearshrub. *Journal of*
1877 *Zhangzhou Teachers College (Nat. Sci.)* 3:83-88.
- 1878 Chia RW, Kim DG, Yimer F (2017) Can afforestation with *Cupressus lusitanica* restore soil C and N stocks depleted
1879 by crop cultivation to levels observed under native systems? *Agriculture, Ecosystems and Environment* 242:67-75.
- 1880 Chidumayo EN, Kwibisa L (2003) Effects of deforestation on grass biomass and soil nutrient status in miombo
1881 woodland, Zambia. *Agriculture Ecosystems & Environment*, 96, 97-105.
- 1882 Chiti T, Grieco E, Perugini L, Rey A & Valentini R (2014) Effect of the replacement of tropical forests with tree
1883 plantations on soil organic carbon levels in the Jomoro district, Ghana. *Plant Soil* 375:47-59.
- 1884 Chone T, Andreux F, Correa JC, Volkoff B, Cerri CC (1991) Changes in organic matter on an Oxisol from the Central
1885 Amazonian forest during eight years as pasture determined by ^{13}C isotope composition. In *Diversity of Environmental*
1886 *Biogeochemistry* Vol. 6 (ed Berthelin J), pp. 397 - 405. Elsevier, Amsterdam.

First Order Draft

- 1887 Cleveland CC, Townsend AR, Schmidt SK, Constance BC (2003) Soil microbial dynamics and biogeochemistry in
1888 tropical forests and pastures, southwestern Costa Rica. *Ecological Applications*, 13, 314-326.
- 1889 Collins HP, Blevins RL, Bundy LG, Christenson DR, Dick WA, Huggins DR & Paul EA (1999) Soil carbon dynamics
1890 in corn-based agroecosystems: results from carbon-13 natural abundance. *Soil Sci Soc Am J* 63:584-591.
- 1891 Conant RT, Paustian K, Elliott ET (2001) Grassland management and conversion into grassland: effects on soil carbon.
1892 *Ecological Applications* 11;2:343-355.
- 1893 Conti G, Perez-Harguindeguy N, Quetier F, Gorne LD, Jaureguiberry P, Bertone GA, Enrico L, Cuchietti A & Diaz
1894 S (2014) Large changes in carbon storage under different land-use regimes in subtropical seasonally dry forests of
1895 southern South America. *Agriculture, Ecosystems and Environment* 197:68-76.
- 1896 Cook RL, Binkley D, Mendes JCT & Stape JL (2014) Soil carbon stocks and forest biomass following conversion of
1897 pasture to broadleaf and conifer plantations in Southeastern Brazil. *Forest Ecology and Management* 324:37-45.
- 1898 Corazza EJ, Silva JE, Resck DVS, Gomes AC (1999) Behaviour of different management systems as a source or sink
1899 of carbon in relation to cerrado vegetation. *Revista Brasileira de Ciencia do Solo*, 23, 425-432.
- 1900 Dai Q, Liu G, Xue S, Yu N, Zhang C, Lan X (2008b) Active organic matter and carbon pool management index of
1901 soil at the abandoned cropland in erosion environment. *Journal of Northwest Forestry University* 23(6):24-28.
- 1902 Dai Q, Liu G, Xue S, Yu N, Zhang C, Lan X 2008b (2008d) Dynamic of Plant Population Characteristics on
1903 Abandoned Arable Land in Eroded Hilly Loess Plateau. *Acta Agriculturae Boreali-occidentalis Sinica* 17(4):320-328.
- 1904 da Silva Junior ML, Desjardins T, Sarrazin M, Silva de Melo V, da Silva Martins PF, Rodrigues Santos E & de
1905 Carvalho CJR (2009) Carbon content in Amazonian Oxisols after forest conversion to pasture. *R. Bras. Ci. Solo*,
1906 33:1603-1611.
- 1907 Dalal RC & Mayer RJ (1986) Long-term trends in fertility of soils under continuous cultivation and cereal cropping
1908 in Southern Queensland. I. Overall changes in soil properties and trends in winter cereal yields. *Aust J Soil Res* 24:265-
1909 279.
- 1910 Dalal RC, Harms BP, Krull E, Wang WJ (2005) Total soil organic matter and its labile pools following mulga (*Acacia*
1911 *aneura*) clearing for pasture development and cropping 1. Total and labile carbon. *Australian Journal of Soil Research*,
1912 43, 13-20.
- 1913 d'Annunzio R, Conche S, Landais D, Saint-Andre L, Joffre R, Barthes BG (2008) Pairwise comparison of soil organic
1914 particle-size distributions in native savannas and Eucalyptus plantations in Congo. *Forest Ecology and Management*,
1915 255, 1050-1056.
- 1916 Dawoe EK, Quashie-Sam JS & Oppong SK (2014) Effect of land-use conversion from forest to cocoa agroforest on
1917 soil characteristics and quality of a Ferric Lixisol in lowland humid Ghana. *Agroforest Systems* 88:87-99.
- 1918 de Blecourt M, Brumme R, Xu J, Corre MD & Veldkamp E (2013) Soil Carbon Stocks Decrease following Conversion
1919 of Secondary Forests to Rubber (*Hevea brasiliensis*) Plantations. *PLoS ONE* 8(7):1-9.
- 1920 de Camargo PB, Trumbore SE, Martinelli LA, Davidson EA, Nepstad DC, Victoria RL (1999) Soil carbon dynamics
1921 in regrowing forest of eastern Amazonia. *Global Change Biology*, 5, 693-702.
- 1922 Dechert G, Veldkamp E, Anas I (2004) Is soil degradation unrelated to deforestation? Examining soil parameters of
1923 land use systems in upland Central Sulawesi, Indonesia. *Plant & Soil* 265:197-209.
- 1924 de Koning GHJ, Veldkamp E & Lopez-Ulloa M (2003) Quantification of carbon sequestration soils following pasture
1925 to forest conversion in northwestern Ecuador. *Global Biogeochemical Cycles* 17:1-12.
- 1926 Delelegn YT, Purahong W, Blazevic A, Yitaferu B, Wubet T, Goransson H, Godbold DL (2017) Changes in land use
1927 alter soil quality and aggregate stability in the highlands of northern Ethiopia. *Nature Scientific Reports* 7:13602.
- 1928 de Neergaard A, Magid J & Mertz O (2008) Soil erosion from shifting cultivation and other smallholder land use in
1929 Sarawak, Malaysia. *Agriculture, Ecosystems, & Environment* 125:182-190.
- 1930 Denef, K, Zotarelli, L, Boddey RM & Six J (2007) Microaggregate-associated carbon as a diagnostic fraction for
1931 management-induced changes in soil organic carbon in two Oxisols. *Soil Biol & Biochem* 39:1165-1172.

- 1932 de Freitas PL, Blancaneaux P, Gavinelli E, Larre-Larrouy MC, Feller C (2000) Nature and level of organic stock in
1933 clayey oxisols under different land use and management systems. *Pesquisa Agropecuaria Brasileira*, 35, 157-170.
- 1934 de Koning GHJ, Veldkamp E, Lopez-Ulloa M (2003) Quantification of carbon sequestration in soils following pasture
1935 to forest conversion in northwestern Ecuador. *Global Biogeochemical Cycles*, 17.
- 1936 de Moraes JFL, Neill C, Volkoff B, Cerri CC, Melillo J, Lima VC, Steudler PA (2002) Soil carbon and nitrogen stocks
1937 following forest conversion to pasture in the Western Brazilian Amazon Basin. *Acta Scientiarum Universidade*
1938 *Estadual de Maringa*, 24, 1369-1376.
- 1939 de Moraes JFL, Volkoff B, Cerri CC, Bernoux M (1996) Soil properties under Amazon forest and changes due to
1940 pasture installation in Rondonia, Brazil. *Geoderma*, 70, 63-81.
- 1941 de Moraes JFL, Volkoff B, Cerri CC, Bernoux M (1996) Soil properties under Amazon forest and changes due to
1942 pasture installation in Rondonia, Brazil. *Geoderma*, 70, 63-81.
- 1943 Desjardins T, Andreux F, Volkoff B & Cerri CC (1994) Organic-carbon and C-13 contents in soils and soil size-
1944 fractions, and their changes due to deforestation and pasture installation in eastern Amazonia. *Geoderma* 61:103-118.
- 1945 Desjardins T, Barros E, Sarrazin M, Girardin C & Mariotti A (2004) Effects of forest conversion to pasture on soil
1946 carbon content and dynamics in Brazilian Amazonia. *Agri Ecosys & Environ* 103:365-373.
- 1947 Detwiler RP (1986) Land-Use Change and the Global Carbon-Cycle - the Role of Tropical Soils. *Biogeochemistry*, 2,
1948 67-93.
- 1949 Eaton JM, Lawrence D (2009) Loss of carbon sequestration potential after several decades of shifting cultivation in
1950 the Southern Yucatan. *Forest Ecology and Management*, 258, 949-958.
- 1951 Eclesia RP, Jobbagy EG, Jackson RB, Biganzoli F, & Piñeiro G (2012) Shifts in soil organic carbon for plantation
1952 and pasture establishment in native forests and grasslands of South America. *Global Change Biology* 18:3237-3251.
- 1953 Eden MJ, McGregor DFM, Vieira NAQ (1990) The Maraca Rain-Forest Project .3. Pasture Development on Cleared
1954 Forest Land in Northern Amazonia. *Geographical Journal*, 156, 283-296.
- 1955 Eden MJ, McGregor DFM & Vieira NAQ (1990) Pasture development on cleared forest land in northern Amazonia.
1956 *Geographical Journal* 156:283-296.
- 1957 Ekanade O (1991) The nature of soil properties under mature forest and plantations of fruiting and exotic trees in the
1958 tropical rain forest fringes of SW Nigeria. *Journal of World Forest Resource Management*, 5, 101-114.
- 1959 Elliott ET, Palm CA, Reuss DE, Monz CA (1991) Organic matter contained in soil aggregates from a tropical
1960 chronosequence – correction for sand and light fraction. *Agriculture Ecosystems & Environment*, 34, 443-451.
- 1961 Elmore AJ, Asner GP (2006) Effects of grazing intensity on soil carbon stocks following deforestation of a Hawaiian
1962 dry tropical forest. *Global Change Biology*, 12, 1761-1772.
- 1963 England JR, Paul KI, Cunningham SC, Madhavan DB, Baker TG, Read Z, Wilson BR, Cavagnaro TR, Lewis T,
1964 Perring MP, Herrmann T & Polglase PJ (2016) Previous land use and climate influence differences in soil organic
1965 carbon following reforestation of agricultural land with mixed-species plantings. *Agri Ecosys & Environ* 227:61-72.
- 1966 Epron D, Marsden C, M'Bou AT, Saint-Andre L, d'Annunzio R & Nouvellon Y (2009) Soil carbon dynamics
1967 following afforestation of a tropical savannah with Eucalyptus in Congo. *Plant & Soil* 323:309-322.
- 1968 Erickson H, Keller M & Davidson E (2001) Nitrogen oxide fluxes and nitrogen cycling during postagricultural
1969 succession and forest fertilization in the humid tropics. *Ecosystems* 4:67-84.
- 1970 Fabrizzi KP, Rice CW, Amado TJC, Fiorin J, Barbagelata P, Melchiori R (2009) Protection of soil organic C and N
1971 in temperate and tropical soils: effect of native and agroecosystems. *Biogeochemistry*, 92, 129-143.
- 1972 Farley KA, Kelly EF & Hofstede RGM (2004) Soil organic carbon and water retention after conversion of grasslands
1973 to pine plantations in the Ecuadorian Andes. *Ecosystems* 7:729-739.
- 1974 Feldpausch TR, Rondon MA, Fernandes ECM, Riha SJ & Wandelli E (2004) Carbon and nutrient accumulation in
1975 secondary forests regenerating on pastures in central Amazonia. *Ecological Applications* 14:S164-S176.

First Order Draft

- 1976 Feller C, Albrecht A, Blanchart E, Cabidoche YM, Chevallier T, Hartmann C, Eschenbrenner V, Larre-Larrouy MC
 1977 & Ndandou JF (2001) Soil organic carbon sequestration in tropical areas. General considerations and analysis of some
 1978 edaphic determinants for Lesser Antilles soils. *Nutrient Cycling in Agroecosystems* 61:19-31.
- 1979 Fernandes SAP, Bernoux M, Cerri CC, Feigl BJ & Piccolo MC (2002) Seasonal variation of soil chemical properties
 1980 and CO₂ and CH₄ fluxes in unfertilized and P-fertilized pastures in an Ultisol of the Brazilian Amazon. *Geoderma*
 1981 107:227-241.
- 1982 Fernandez I, Carrasco B & Cabaneiro A (2012) Evolution of soil organic matter composition and edaphic carbon
 1983 effluxes following oak forest clearing for pasture: climate change implications. *European Journal of Forest Research*
 1984 131:1681-1693.
- 1985 Fisher MJ, Rao IM, Ayarza MA, Lascano CE, Sanz JI, Thomas RJ & Vera RR (1994) Carbon storage by introduced
 1986 deep-rooted grasses in the South American savannas. *Nature* 371:236-238.
- 1987 Follett RF, Paul EA, Leavitt SW, Halvorson AD, Lyon D & Peterson GA (1997) Carbon isotope ratios of Great Plains
 1988 soils and in wheat-fallow systems. *Soil Sci Soc Am J* 61:1068-1077.
- 1989 Freibauer A (1996) Short Term Effects of Land Use on Aggregates, Soil Organic Matter, and P Status of a Clayey
 1990 Cerrado Oxisol, Brazil. University Bayreuth, Bayreuth, 55 pp.
- 1991 Freixo AA, Machado PLOA, Guimaraes CM, Silva CA, Fadigas FS (2002) Carbon and nitrogen storage and organic
 1992 fraction distribution of a Cerrado Latosol under different cultivation systems. *Revista Brasileira de Ciencia do Solo*,
 1993 26, 425-434.
- 1994 Fu BJ, Guo XD, Chen LD et al (2001) Soil nutrient changes due to land use changes in Northern China: a case study
 1995 in Zunhua County, Hebei Province. *Soil Use and Management* 17:294-296.
- 1996 Fu B, Chen L, Ma K, Zhou H, Wang J (2000) The relationships between land use and soil conditions in the hilly area
 1997 of the loess plateau in northern Shaanxi, China. *Catena* 39:69-78.
- 1998 Fu H, Chen Y, Zhou Z, Ai D, Zhou Z (2003) Change of vegetation and soil environment of desert grassland in the
 1999 early period of restoration in Alxa, Inner Mongolia. *Journal of Desert Research* 23(6):661-664.
- 2000 Fuhrmann S, Neufeldt H, Westerhof R, Ayarza MA, da Silva JE, Zech W (1999) Soil organic carbon, carbohydrates,
 2001 amino sugars, and potentially mineralisable nitrogen under different land-use systems in Oxisols of the Brazilian
 2002 Cerrados. In *Sustainable Land Management for the Oxisols of the Latin American Savannas* (eds Thomas R, Ayarza
 2003 MA), pp. 110-122. CIAT Publ., Cali, Columbia.
- 2004 Fujisaka S, Castilla C, Escobar G, Rodrigues V, Veneklaas EJ, Thomas R, Fisher M (1998) The effects of forest
 2005 conversion on annual crops and pastures: Estimates of carbon emissions and plant species loss in a Brazilian Amazon
 2006 colony. *Agriculture Ecosystems & Environment*, 69, 17-26.
- 2007 Garcia-Franco N, Wiesmeier M, Goberna M, Martinez-Mena M, Albaladejo J (2014) Carbon dynamics after
 2008 afforestation of semiarid shrublands: Implications of site preparation techniques. *Forest Ecology and Management*
 2009 319:107-115.
- 2010 Garcia-Oliva F, Casar I, Morales P & Maass JM (1994) Forest-to-pasture conversion influences on soil organic-carbon
 2011 dynamics in a tropical deciduous forest. *Oecologia* 99:392-396.
- 2012 Garcia-Oliva F, Lancho JFG, Montano NM, Islas P (2006) Soil Carbon and nitrogen dynamics followed by a forest-
 2013 to-pasture conversion in western Mexico. *Agroforestry Systems*, 66, 93-100.
- 2014 Garcia-Oliva F, Sanford RL, Kelly E (1999) Effects of slash-and-burn management on soil aggregate organic C and
 2015 N in a tropical deciduous forest. *Geoderma*, 88, 1-12.
- 2016 Gamboa AM & Galicia L (2011) Differential influence of land use/cover change on topsoil carbon and microbial
 2017 activity in low-latitude temperate forests. *Agri Ecosys & Environ* 142:280-290.
- 2018 Geissen V, Pena-Pena K, Huerta E (2009) Effects of different land use on soil chemical properties, decomposition
 2019 rate and earthworm communities in tropical Mexico. *Pedobiologia*, 53, 75-86.
- 2020 Ghuman BS, Lal R, Shearer W (1991) Land Clearing and Use in the Humid Nigerian Tropics .1. Soil Physical-
 2021 Properties. *Soil Science Society of America Journal*, 55, 178-183.

- 2022 Girma T (1998) Effect of cultivation on physical and chemical properties of a Vertisol in Middle Awash Valley,
2023 Ethiopia. *Comm Soil Sci Plant Anal* 29:587-598.
- 2024 Gong J, Chen L, Fu B, Li Y, Huang Z, Huang Y, Peng H (2004) Effects of land use and vegetation restoration on soil
2025 quality in a small catchment of the Loess Plateau. *Chinese Journal of Applied Ecology* 15(12):2292-2296.
- 2026 Gosling P, van der Gast C, Bending GD (2017) Converting highly productive arable cropland in Europe to grassland:
2027 a poor candidate for carbon sequestration. *Nature Scientific Reports* 7:10493.
- 2028 Gregorich EG, Ellert BH, Drury CF & Liang BC (1996) Fertilization effects on soil organic matter turnover and corn
2029 residue C storage. *Soil Sci Soc Am J* 60:472-476.
- 2030 Guggenberger G & Zech W (1999) Soil organic matter composition under primary forest, pasture, and secondary
2031 forest succession, Region Hueta Norte, Costa Rica. *Forest Ecology and Management* 124:93-104.
- 2032 Han J, Han Y, Sun T, Wang X (2004) Effects of returning cultivated land to herbage on soil organic matter and
2033 nitrogen in the agro-pastoral transitional zone of north China. *Acta Prataculturae Sinica* 13(4):21-28.
- 2034 Han Y, Han J, Wang K, Zhang Y (2005) Effects of utilization periods on cropland soil chemical properties in
2035 the farming2pastoral transitional zone after replaced with pasture. *Pratacultural Science* 22(3):50-53.
- 2036 Harms BP, Dalal RC, Cramp AP (2005) Changes in soil carbon and soil nitrogen after tree clearing in the semi-arid
2037 rangelands of Queensland. *Australian Journal of Botany*, 53, 639-650.
- 2038 Harden JW, Sharpe JM, Parton WJ, Ojima DS, Fries TL, Huntington TG & Dabney SM (1999) Dynamic replacement
2039 and loss of soil carbon on eroding cropland. *Global Biogeochem Cycles* 13:885-901.
- 2040 Hartemink AE (1997) Soil fertility decline in some major soil groupings under permanent cropping in Tanga region,
2041 Tanzania. *Geoderma*, 75, 215-229.
- 2042 He X, Chang Q, Wen Z, Jiao F, Li R (2006) Desertified soil fertility under different artificial vegetations in farming-
2043 pasturing interlock zone of northern Shaanxi Province. *Journal of Desert Research* 26(6):915-919.
- 2044 Hertl D, Harteveld MA, Leuschner C (2009) Conversion of a tropical forest into agroforest alters the fine root-related
2045 carbon flux to the soil. *Soil Biology & Biochemistry* 41:481-490.
- 2046 Hölscher D, Ludwig B, Moller RF, Folster H (1997) Dynamic of soil chemical parameters in shifting agriculture in
2047 the Eastern Amazon. *Agriculture Ecosystems & Environment*, 66, 153-163.
- 2048 Hou X, Han X, Wang S, Song C (2008) Different Land Uses and Management Effects on Soil Fertilities in Black
2049 Soil. *Journal of Soil and Water Conservation* 22(6):99-104.
- 2050 Hsieh YP (1996) Soil organic carbon pools of two tropical soils inferred by carbon signatures. *Soil Science Society
2051 of America Journal*, 60, 1117-1121.
- 2052 Hu Y, Zeng D, Fan Z, Ai GY (2007) Effects of degraded sandy grassland afforestation on soil quality in semi-arid
2053 area of Northern China. *Chinese Journal of Applied Ecology* 18(11):2391-2397.
- 2054 Huang D, Wang K, Wu WL (2007) Dynamics of soil physical and chemical properties and vegetation succession
2055 characteristics during grassland desertification under sheep grazing in an agro-pastoral transition zone in Northern
2056 China. *Journal of Arid Environments* 70:120-136.
- 2057 Hughes RF, Kauffman JB & Cummings DL (2002) Dynamics of aboveground and soil carbon and nitrogen stocks
2058 and cycling of available nitrogen along a land-use gradient in Rondonia, Brazil. *Ecosystems* 5:244-259.
- 2059 Hughes RF, Kauffman JB, Cummings DL (2000) Fire in the Brazilian Amazon 3. Dynamics of biomass, C, and
2060 nutrient pools in regenerating forests. *Oecologia*, 124, 574-588.
- 2061 Hughes RF, Kauffman JB, Jaramillo VJ (2000) Ecosystem-scale impacts of deforestation and land use in a humid
2062 tropical region of Mexico. *Ecological Applications* 10:515-527.
- 2063 Ihori T, Burke IC, Lauenroth WK & Coffin DP (1995) Effects of cultivation and abandonment on soil organic matter
2064 in northeastern Colorado. *Soil Sci Soc Am J* 59:1112-1119.

First Order Draft

- 2065 Ishizuka S, Iswandi A, Nakajima Y, Yonemura S, Sudo S, Tsuruta H, Murdiyaso D (2005) The variation of greenhouse
2066 gas emissions from soils of various land-use/cover types in Jambi province, Indonesia. *Nutrient Cycling in*
2067 *Agroecosystems* 71:17-32.
- 2068 Islam KR & Weil RR (2000) Land use effects on soil quality in a tropical forest ecosystem of Bangladesh. *Agriculture,*
2069 *Ecosystems & Environment* 79:9-16.
- 2070 Jakelaitis A, da Silva AA, dos Santos JB, Vivian R (2008) Quality of soil surface layer under forest, pastures and
2071 cropped areas. *Pesquisa Agropecuaria Tropical*, 38, 118-127.
- 2072 Janssen BH, Wienk JF (1990) From forest to arable land. In *Mechanized annual cropping on low fertility acid soils in*
2073 *the humid tropics - A case study of the Zanderij soils in Suriname*
- 2074 Jaramillo VJ, Kauffman JB, Renteria-Rodriguez L, Cummings DL & Ellingson LJ (2003) Biomass, carbon, and
2075 nitrogen pools in Mexican tropical dry forest landscapes. *Ecosystems* 6:609-629.
- 2076 Jia S, He X, Chen Y (2004) Effect of Land Abandonment on Soil Organic Carbon Sequestration in Loess Hilly Areas.
2077 *Journal of Soil and Water Conservation* 18(3):78-81.
- 2078 Jia X, Li X, Li Y (2007) Soil organic carbon and nitrogen dynamics during the re-vegetation progress in the arid desert
2079 region. *J. Plant Ecology (Chinese version)* 31:66-74.
- 2080 Jimenez JJ, Lal R, Leblanc HA & Russo RO (2007) Soil organic carbon pool under native tree plantations in the
2081 Caribbean lowlands of Costa Rica. *Forest Ecology & Management* 241:134-144.
- 2082 Jun W, Liqing S (2007) Effects of Land Use on Soil Nutrients in Tibetan Region, Northwest Yunnan, China. *Journal*
2083 *of Northeast Forestry University* 35(10):45-47.
- 2084 Juo ASR, Franzluebbers K, Dabiri A, Ikhile B (1995) Changes in soil properties during long-term fallow and
2085 continuous cultivation after forest clearing in Nigeria. *Agriculture Ecosystems & Environment*, 56, 9-18.
- 2086 Juo ASR, Lal R (1977) Effect of fallow and continuous cultivation on chemical and physical properties of an Alfisol
2087 in western Nigeria. *Plant and Soil*, 47, 567-584.
- 2088 Juo ASR, Lal R (1979) Nutrient Profile in a Tropical Alfisol under Conventional and No-Till Systems. *Soil Science*,
2089 127, 168-173.
- 2090 King JA, Campbell BM (1994) Soil organic matter relations in 5 land-cover types in Miombo Region (Zimbabwe).
2091 *Forest Ecology and Management*, 67, 225-239.
- 2092 Kainer KA, Duryea ML, de Macedo NC & Williams K (1998) Brazil nut seedling establishment and autecology in
2093 extractive reserves of Acre, Brazil. *Ecological Applications* 8:397-410.
- 2094 Karhu K, Wall A, Vanhala P, Liski J, Esala M & Regina K (2011) Effects of afforestation and deforestation on boreal
2095 soil carbon stocks-Comparison of measured C stocks with Yasso07 model results. *Geoderma* 164:33-45.
- 2096 Kawanabe S, Nan Y, Zhang S, Oshida T (2000) A Change of Vegetation and Soil of the Desertified Grasslands in
2097 the Process of Recovery. 1. At the sites of the sand dune and the flat sand land. *Soil and Water Conservation*
2098 *Technology Bulletin* 4:16–20.
- 2099 Keith, AM, Rowe RL, Parmar K, Perks MP, Mackie E, Dondini M & McNamara NP (2015) Implications of land-use
2100 change to short rotation forestry in Great Britain for soil and biomass carbon. *Global Change Biology Bioenergy*
2101 7:541-552.
- 2102 Kotto-Same J, Woomer PL, Appolinaire M, Louis Z (1997) Carbon dynamics in slash-and-burn agriculture and land
2103 use alternatives of the humid forest zone in Cameroon. *Agriculture Ecosystems & Environment*, 65, 245-256.
- 2104 Koutika LS, Bartoli F, Andreux F, Cerri CC, Burtin G, Chone T & Philippy R (1997) Organic matter dynamics and
2105 aggregation in soils under rain forest and pastures of increasing age in the eastern Amazon Basin. *Geoderma* 76:87-
2106 112.
- 2107 Manlay RLJ, Ickowicz A, Masse D, Floret C, Richard D, Feller C (2004) Spatial carbon, nitrogen and phosphorus
2108 budget of a village in the West African savanna - I. Element pools and structure of a mixed-farming system.
2109 *Agricultural Systems*, 79, 55-81.

- 2110 Krishnaswamy J & Richter DD (2002) Properties of advanced weathering-stage soils in tropical forests and pastures.
2111 Soil Sci Soc Am J 66:244-253.
- 2112 Lal R (1998) Land use and soil management effects on soil organic matter dynamics on alfisols in western Nigeria.
2113 Soil Processes and the Carbon Cycle, 109-126.
- 2114 Lemenih M, Karlun E, Olsson M (2005) Soil organic matter dynamics after deforestation along a farm field
2115 chronosequence in southern highlands of Ethiopia. Agriculture Ecosystems & Environment, 109, 9-19.
- 2116 Lemenih M, Karlun E, Olsson M (2005) Assessing soil chemical and physical property responses to deforestation
2117 and subsequent cultivation in smallholders farming system in Ethiopia. Agriculture, Ecosystems & Environment
2118 105:373-386.
- 2119 Lemma B, Kleja DB, Nilsson I & Olsson M (2006) Soil carbon sequestration under different exotic tree species in
2120 southwestern highlands of Ethiopia. Geoderma 136:886-898.
- 2121 Lepsch IF, Menk JRF, Oliveira JB (1994) Carbon Storage and Other Properties of Soils under Agriculture and Natural
2122 Vegetation in Sao-Paulo State, Brazil. Soil Use and Management, 10, 34-42.
- 2123 Li M, Dong Y, Qi Y, Geng Y (2005) Effect of Land-use Change on the Contents of C & N in Temperate Grassland
2124 Soils. Grassland of China 27(1):1-6.
- 2125 Li Y, Li X, Zhang P, Yin P (2007) Effects of land use on organic carbon and nutrient contents in desert soil. Journal
2126 of Gansu Agricultural University 42(2):103-107.
- 2127 Li YY, Shao MA, ZhenG JY, Li QF (2007) Impact of grassland recovery and reconstruction on soil organic carbon
2128 in the northern Loess Plateau. Acta Ecologica Sinica 27(6):2279-2287.
- 2129 Li XG, Wang ZF, Ma QF, Li FM (2007a) Crop cultivation and intensive grazing affect organic C pools and aggregate
2130 stability in arid grassland soil. Soil & Tillage Research 95:172-181.
- 2131 Lilienfein J, Wilcke W, Vilela L, Ayarza A, do Carmo Lima S, Zech W (2003) Soil fertility under native cerrado and
2132 pasture in the Brazilian savanna. Soil Sci Soc Am J 67:1195-1205.
- 2133 Lima AMN, Silva IR, Neves JCL, Novais RF, Barros NF, Mendonca ES, Smyth TJ, Moreira MS & Leite FP (2006)
2134 Soil organic carbon dynamics following afforestation of degraded pastures with Eucalyptus in southeastern Brazil.
2135 Forest Ecology & Management 235:219-231.
- 2136 Lisboa C, Conant RT, Haddix ML, Cerri CEP & Cerri CC, (2009) Soil carbon turnover measurement by physical
2137 fractionation at a forest-to-pasture chronosequence in the Brazilian Amazon. Ecosystems 12:1212-1221.
- 2138 Lugo AE, Sanchez MJ (1986) Land-Use and Organic-Carbon Content of Some Subtropical Soils. Plant and Soil, 96,
2139 185-196.
- 2140 Luizao RCC, Bonde TA & Rosswall T (1992) Seasonal variation of soil microbial biomass-the effects of clearfelling
2141 a tropical rainforest and establishment of pasture in the Central Amazon. Soil Biology & Biochemistry 24:805-813.
- 2142 Ma K, He X, Ma B, Luo D, Ma Y (2006) Effects of land use pattern on soil in the Loess Plateau of south Ningxia.
2143 Ecology and Environment 15(6):1231-1236.
- 2144 Macedo MO, Resende AS, Garcia PC, Boddey RM, Jantalia CP, Urquiaga S, Campello EFC & Franco AA (2008)
2145 Changes in soil C and N stocks and nutrient dynamics 13 years after recovery of degraded land using leguminous
2146 nitrogen-fixing trees. Forest Ecology & Management 255:1516-1524.
- 2147 Maia SMF, Ogle SM, Cerri CEP & Cerri CC (2009) Effects of grassland management on soil carbon sequestration in
2148 Rondonia and Mato Grosso states, Brazil. Geoderma 149:84-91.
- 2149 Makumba W, Akinnifesi FK, Janssen B, Oenema O (2007) Long-term impact of a gliricidia-maize intercropping
2150 system on carbon sequestration in southern Malawi. Agriculture Ecosystems & Environment, 118, 237-243.
- 2151 Manlay RJ, Kaire M, Masse D, Chotte JL, Ciornei G & Floret C (2002) Carbon, nitrogen and phosphorus allocation
2152 in agro-ecosystems of a West African savanna I. The plant component under semi-permanent cultivation. Agriculture,
2153 Ecosystems, & Environment 88:215-232.

First Order Draft

- 2154 Manlay RJ, Masse D, Chotte JL, Feller C, Kaire M, Fardoux J & Pontanier R (2002) Carbon, nitrogen and phosphorus
2155 allocation in agro-ecosystems of a West African savanna II. The soil component under semi-permanent cultivation.
2156 *Agriculture, Ecosystems, & Environment* 88:233-248.
- 2157 Maquere V, Laclau JP, Bernoux M, Saint-Andre L, Goncalves JLM, Cerri CC, Piccolo MC & Ranger J (2008)
2158 Influence of land use (savanna, pasture, Eucalyptus plantations) on soil carbon and nitrogen stocks in Brazil. *European*
2159 *Journal of Soil Science* 59:863-877.
- 2160 Marin-Spiotta E, Silver W, Swanston CW & Ostertag R (2009) Soil organic matter dynamics during 80 years of
2161 reforestation of tropical pastures. *Global Change Biology* 15:1584-1597.
- 2162 Markewitz D, Davidson E, Moutinho P & Nepstad D (2004) Nutrient loss and redistribution after forest clearing on a
2163 highly weathered soil in Amazonia. *Ecological Applications* 14:S177-S199.
- 2164 Martins EdL, Coringa JdES, Weber OLdS (2009) Organic carbon in granulometric fraction and in humic substances
2165 of a Brazilian Oxisol under different land use systems. *Acta Amazonica*, 39, 655-660.
- 2166 Masto RE, Chhonkar PK, Purakayastha TJ, Patra AK & Singh D (2008) Soil quality indices for evaluation of long-
2167 term land use and soil management practices in semi-arid sub-tropical India. *Land Degradation & Development*
2168 19:516-529.
- 2169 Materechera SA, Mkhabela TS (2001) Influence of land-use on properties of a ferrallitic soil under low external input
2170 farming in southeastern Swaziland. *Soil & Tillage Research*, 62, 15-25.
- 2171 McGrath DA, Smith CK, Gholz HL, Oliveira FD (2001) Effects of land-use change on soil nutrient dynamics in
2172 Amazonia. *Ecosystems*, 4, 625-645.
- 2173 Mendham DS, O'Connell AM & Grove TS (2003) Change in soil carbon after land-clearing or afforestation in highly
2174 weathered lateritic and sandy soils of southwestern Australia. *Agriculture, Ecosystems & Environment* 95:143-156.
- 2175 Mikhailova EA, Bryant RB, Vassenev II, Schwager SJ & Post CJ (2000) Cultivation effects on soil carbon and
2176 nitrogen contents at depth in the Russian Chernozem. *Soil Sci Soc Am J* 64:738-745.
- 2177 Morris AR (1984) A comparison of soil nutrient levels under grassland and two rotations of *Pinus patula* in the Usutu
2178 Forest, Swaziland. *Proceedings IUFRO Symposium on Site and Productivity of Fast Growing Plantations. Volume 2.*,
2179 881-892.
- 2180 Motavalli PP, Discekici H, Kuhn J (2000) The impact of land clearing and agricultural practices on soil organic C
2181 fractions and CO₂ efflux in the Northern Guam aquifer. *Agriculture Ecosystems & Environment*, 79, 17-27.
- 2182 Motavalli PP & McConnell J (1998) Land use and soil nitrogen status in a tropical pacific island environment. *Journal*
2183 *of Environmental Quality* 27:119-123.
- 2184 Muller MML, Guimaraes MD, Desjardins T, Martins PFD (2001) Pasture degradation in the Amazon region: soil
2185 physical properties and root growth. *Pesquisa Agropecuaria Brasileira*, 36, 1409-1418.
- 2186 Mutuo PK, Cadisch G, Albrecht A, Palm CA, Verchot L (2005) Potential of agroforestry for carbon sequestration and
2187 mitigation of greenhouse gas emissions from soils in the tropics. *Nutrient Cycling in Agroecosystems*, 71, 43-54.
- 2188 Nadal-Romero E, Cammeraat E, Perez-Cardiel E & Lasanta T (2016) How do soil organic carbon stocks change after
2189 cropland abandonment in Mediterranean humid mountain areas? *Science of Total Environment* 566-567:741-752.
- 2190 Navarrete IA, Tsutsuki K (2008) Land-use impact on soil carbon, nitrogen, neutral sugar composition and related
2191 chemical properties in a degraded Ultisol in Leyte, Philippines. *Soil Science and Plant Nutrition*, 54, 321-331.
- 2192 Navarrete D, Sitch S, Aragao LEOC, & Pedroni L (2016) Conversion from forests to pastures in the Colombian
2193 Amazon leads to contrasting soil carbon dynamics depending on land management practices. *Global Change Biology*
2194 22:3503-3517.
- 2195 Neill C, Melillo JM, Steudler PA, Cerri CC, de Moraes JFL, Piccolo MC & Brito M (1997) Soil carbon and nitrogen
2196 stocks following forest clearing for pasture in the southwestern Brazilian Amazon. *Ecological Applications* 7:1216-
2197 1225.
- 2198 Neill C, Cerri CC, Melillo JM, Feigl BJ, Steudler PA, Moraes JFL, Piccolo MC (1998) Stocks and dynamics of soil
2199 carbon following deforestation for pasture in Rondonia. *Soil Processes and the Carbon Cycle*, 9-28.

- 2200 Neufeldt H, Resck DVS, Ayarza MA (2002) Texture and land-use effects on soil organic matter in Cerrado Oxisols,
2201 Central Brazil. *Geoderma*, 107, 151-164.
- 2202 Ogle SM, Breidt FJ, Paustian K (2005) Agricultural management impacts on soil organic carbon storage under moist
2203 and dry climatic conditions of temperate and tropical regions/ *Biogeochemistry* 72:87-121.
- 2204 Ogunkunle AO, Eghaghara OO (1992) Influence of land-use on soil properties in a forest region of southern Nigeria.
2205 *Soil Use and Management*, 8, 121-125.
- 2206 Ohta S (1990) Initial soil changes associated with afforestation with *Acacia auriculiformis* and *Pinus kesiya* on
2207 denuded grasslands of the Pantabangan Area, Central Luzon, the Phillipines. *Soil Sci. Plant Nutr.* 36:633-643.
- 2208 Osher LJ, Matson PA & Amundson R (2003) Effect of land-use change on soil carbon in Hawaii. *Biogeochemistry*
2209 65:213-232.
- 2210 Parfitt RL, Theng BKG, Whitton JS & Shepherd TG (1997) Effects of clay minerals and land use on organic matter
2211 pools. *Geoderma* 75:1-12.
- 2212 Paul S, Flessa H, Veldkamp E & Lopez-Ulloa M (2008) Stabilization of recent soil carbon in the humid tropics
2213 following land use changes: evidence from aggregate fractionation and stable isotope analyses. *Biogeochemistry*
2214 87:247-263.
- 2215 Pennock DJ & van Kessel C (1997) Effect of agriculture and of clear-cut forest harvest on landscape-scale soil organic
2216 carbon storage in Saskatchewan. *Can J Soil* 77:211-218.
- 2217 Perrin AS, Fujisaki K, Petitjean C, Sarrazin M, Godet M, Garric B, Horth JC, Balbino LC, Filho AS, de Almeida
2218 Machado PLO & Brossard, M (2014) Conversion of forest to agriculture in Amazonia with the chop-and-mulch
2219 method: does it improve the soil carbon stock? *Agriculture, Ecosystems and Environment* 184:101-114.
- 2220 Piccolo GA, Andriulo AE & Mary B (2008) Changes in soil organic matter under different land management in
2221 Misiones Province (Argentina). *Scientia Agricola (Piracicaba, Braz.)* 65:290-297.
- 2222 Potter KN, Torbert HA, Johnson HB & Tischler CR (1999) Carbon storage after long-term grass establishment on
2223 degraded soils. *Soil Sci* 164:718-725.
- 2224 Potvin C, Whidden E & Moore T (2004) A case study of carbon pools under three different land-uses in Panama.
2225 *Climatic Change* 67:291-307.
- 2226 Powers JS & Veldkamp E (2005) Regional variation in soil carbon and $\delta^{13}\text{C}$ in forests and pastures of northeastern
2227 Costa Rica. *Biogeochemistry* 72:315-336.
- 2228 Powers JS (2004) Changes in soil carbon and nitrogen after contrasting land-use transitions in northeastern Costa Rica.
2229 *Ecosystems* 7:134:146.
- 2230 Rangel OJP, Silva CA & Guimarães PTG (2007) Stock and Fractions of the Organic Matter of Latosol Cultivated
2231 with Coffee in Different Planting Spacings. *Revista Brasileira de Ciência do Solo*, 31, 1341-1353.
- 2232 Rasiah V, Florentine SK, Williams BL, Westbrooke ME (2004) The impact of deforestation and pasture abandonment
2233 on soil properties in the wet tropics of Australia. *Geoderma*, 120, 35-45.
- 2234 Reeder JD, Schuman GE & Bowman RA (1998) Soil C and N changes on conservation reserve program lands in the
2235 central Great Plains. *Soil Tillage Res* 47:339-349.
- 2236 Reiners WA, Bouwman AF, Parsons WFJ & Keller M (1994) Tropical rainforest conversion to pasture: changes in
2237 vegetation and soil properties. *Ecological Applications* 4:363-377.
- 2238 Resh SC, Binkley D & Parrotta JA (2002) Greater soil carbon sequestration under nitrogen-fixing trees compared with
2239 *Eucalyptus* species. *Ecosystems* 5:217-231.
- 2240 Rhoades CC, Eckert GE & Coleman DC (2000) Soil carbon differences among forest, agriculture, and secondary
2241 vegetation in lower montane Ecuador. *Ecological Applications* 10:497-505.
- 2242 Richards AE, Dalal RC & Schmidt S (2007) Soil carbon turnover and sequestration in native subtropical tree
2243 plantations. *Soil Biology & Biochemistry* 39:2078-2090.

First Order Draft

- 2244 Riezebos HT, Loerts AC (1998) Influence of land use change and tillage practice on soil organic matter in southern
2245 Brazil and eastern Paraguay. *Soil & Tillage Research*, 49, 271-275.
- 2246 Rojas JM, Prause J, Sanzano GA, Arce OEA, Sanchez MC (2016) Soil quality indicators selection by mixed models
2247 and multivariate techniques in deforested areas for agricultural use in NW of Chaco, Argentina. *Soil & Tillage Research*
2248 155:250-262.
- 2249 Ronggui Wu, H. Tiessen (2002) Effect of Land Use on Soil Degradation in Alpine Grassland Soil, China. *Soil Science*
2250 *Society of America Journal* 66:1648-1655.
- 2251 Roscoe R & Buurman P (2003) Tillage effects on soil organic matter in density fractions of a Cerrado Oxisol. *Soil &*
2252 *Tillage Research* 70:107-119.
- 2253 Rossi J, Govaerts A, De Vos B, Verbist B, Vervoort A, Poesen J, Muys B & Deckers J (2009) Spatial structures of
2254 soil organic carbon in tropical forests--a case study of Southeastern Tanzania. *Catena* 77:19-27.
- 2255 Russell AE, Raich JW, Fisher RF & Valverde-Barrantes OJ (2007) Tree species effects on soil properties in
2256 experimental plantations in tropical moist forest. *Soil Sci Soc Am J* 71:1389-1397.
- 2257 Saggat S, Yeates GW & Shepherd TG (2001) Cultivation effects on soil biological properties, microfauna and organic
2258 matter dynamics in Eutric Gleysol and Gleyic Luvisol soils in New Zealand. *Soil Tillage Res* 58:55-68.
- 2259 Sa JCDM, Cerri CC, Dick WA, Lal R, Venske Filho SP, Piccolo MC, Feigl BE (2001) Organic matter dynamics and
2260 carbon sequestration rates for a tillage chronosequence in a Brazilian Oxisol. *Soil Science Society of America Journal*,
2261 65, 1486-1499.
- 2262 Saha SK, Nair PKR, Nair VD & Kumar BM (2009) Soil carbon stock in relation to plant diversity of homegardens in
2263 Kerala, India. *Agroforestry Systems* 76:53-65.
- 2264 Saha SK, Nair PKR, Nair VD & Kumar BM (2010) Carbon storage in relation to soil size-fractions under tropical
2265 tree-based land-use systems. *Plant & Soil* 328:433-446.
- 2266 Salimon CI, Davidson EA, Victoria RL, Melo AWF (2004) CO₂ flux from soil in pastures and forests in southwestern
2267 Amazonia. *Global Change Biology* 10:833-843.
- 2268 Sanchez PA, Villachica JH, Bandy DE (1983) Soil fertility dynamics after clearing a tropical rainforest in Peru. *Soil*
2269 *Science Society of America Journal*, 47, 1171-1178.
- 2270 Saynes V, Hidalgo C, Etchevers JD, Campo JE (2005) Soil C and N dynamics in primary and secondary seasonally
2271 dry tropical forests in Mexico. *Applied Soil Ecology*, 29, 282-289.
- 2272 Schedlbauer JL, Kavanagh KL (2008) Soil carbon dynamics in a chronosequence of secondary forests in northeastern
2273 Costa Rica. *Forest Ecology and Management*, 255, 1326-1335.
- 2274 Schiffman PM & Johnson WC (1989) Phytomass and detrital carbon storage during forest regrowth in the southeastern
2275 United States Piedmont. *Can J For Res* 19:69-78.
- 2276 Schwendenmann L & Pendall E (2006) Effects of forest conversion into grassland on soil aggregate structure and
2277 carbon storage in Panama: evidence from soil carbon fractionation and stable isotopes *Plant & Soil* 288: 217-232.
- 2278 Shang C, Tiessen H (1997) Organic matter lability in a tropical oxisol: Evidence from shifting cultivation, chemical
2279 oxidation, particle size, density, and magnetic fractionations. *Soil Science*, 162, 795-807.
- 2280 Sheng X, Liu Y, Sun J (2004) Effect of land-use change on soil habitat in north Hebei plateau during last 50 year.
2281 *Chinese Journal of Applied Ecology* 15(4):589-592.
- 2282 Siband P (1974) Evolution of soil characters and fertility in a red soil in Casamance. *Agronomie Tropicale*, 29, 1228-
2283 1248.
- 2284 Silva AM, Nogueira DP, Ikematsu P, et al. (2009) Carbon stocks and isotopic composition of the organic matter in
2285 soils covered by native vegetation and pasture in Sorocaba, SP, Brazil. *International Journal of Environmental*
2286 *Research*, 3, 435-440.
- 2287 Silver WL, Kueppers LM, Lugo AE, Ostertag R & Matzek V (2004) Carbon sequestration and plant community
2288 dynamics following reforestation of tropical pasture. *Ecological Applications* 14:1115-1127.

- 2289 Sitompul SM, Hairiah K, Cadisch G, Van Noordwijk M (2000) Dynamics of density fractions of macro-organic matter
2290 after forest conversion to sugarcane and woodlots, accounted for in a modified Century model. *Netherlands Journal*
2291 *of Agricultural Science*, 48, 61-73.
- 2292 Six J, Elliott ET, Paustian K & Doran JW (1998) Aggregation and soil organic matter accumulation in cultivated and
2293 native grassland soils. *Soil Sci Soc Am J* 62:1367-1377.
- 2294 Six J, Paustian K, Elliott ET & Combrink C (2000) Soil structure and organic matter: I. Distribution of aggregate-
2295 size classes and aggregate-associated carbon. *Soil Sci Soc Am J* 64:681-689.
- 2296 Slobodian N, Van Rees K & Pennock D (2002) Cultivation-induced effects on belowground biomass and organic
2297 carbon. *Soil Sci Soc Am J* 66:924-930.
- 2298 Smiley GL & Kroschel J (2008) Temporal changes in carbon stocks of cocoa-gliricidia agroforests in Central Sulawesi,
2299 Indonesia. *Agroforest Systems* 73:219-231.
- 2300 Smith CK, Oliveira FD, Gholz HL & Baima A (2002) Soil carbon stocks after forest conversion to tree plantations in
2301 lowland Amazonia, Brazil. *Forest Ecology and Management* 164:257-263.
- 2302 Sohng J, Singhakumara BMP & Ashton MS (2017) Effects on soil chemistry of tropical deforestation for agriculture
2303 and subsequent reforestation with special reference to changes in carbon and nitrogen. *Forest Ecology & Management*
2304 389:331-340.
- 2305 Solomon D, Fritzsche F, Lehmann J, Tekalign M & Zech W (2002) Soil organic matter dynamics in the subhumid
2306 agroecosystems of the Ethiopian Highlands: evidence from natural ¹³C abundance and particle-size fractionation. *Soil*
2307 *Sci Soc Am J* 66: 969-978.
- 2308 Solomon D, Lehmann J, Kinyangi J, et al. (2007) Long-term impacts of anthropogenic perturbations on dynamics and
2309 speciation of organic carbon in tropical forest and subtropical grassland ecosystems. *Global Change Biology*, 13, 511-
2310 530.
- 2311 Solomon D, Lehmann J, Zech W (2000) Land use effects on soil organic matter properties of chromic luvisols in semi-
2312 arid northern Tanzania: carbon, nitrogen, lignin and carbohydrates. *Agriculture Ecosystems & Environment*, 78, 203-
2313 213.
- 2314 Sommer R, Denich M & Vlek PLG (2000) Carbon storage and root penetration in deep soils under small-farmer land-
2315 use systems in the Eastern Amazon region, Brazil. *Plant and Soil* 219:231-241.
- 2316 Srivastava SC, Singh JS (1991) Microbial-C, microbial-N and microbial-P in dry tropical forests soils – effects of
2317 alternative land-uses and nutrient flux. *Soil Biology & Biochemistry*, 23, 117-124.
- 2318 Sparling GP, Schipper LA, Hewitt AE & Degens BP (2000) Resistance to cropping pressure of two New Zealand soils
2319 with contrasting mineralogy. *Aust J Soil Res* 38:85-100.
- 2320 Su Y (2007) Soil carbon and nitrogen sequestration following the conversion of cropland to alfalfa forage land in
2321 northwest China. *Soil & Tillage Research* 92:181-189.
- 2322 Su YZ, Zhao HL, Zhang TH, Zhao XY (2004a) Soil properties following cultivation and non-grazing of a semi-arid
2323 sandy grassland in northern China. *Soil & Tillage Research* 75:27-36.
- 2324 Su Y, Li Y, Zhao H (2006) Soil properties and their spatial pattern in a degraded sandy grassland under post-grazing
2325 restoration, Inner Mongolia, northern China. *Biogeochemistry* 79:297-314.
- 2326 Su Y, Zhao H, Li Y (2004b) Spatial pattern of soil chemical properties in degraded sandy grassland under post-grazing
2327 natural restoration in Horqin sandy land. *Acta Pedologica sinica* 41(3):369-374.
- 2328 Su Y, Zhao H, Zhang T, Cui J (2002a) Characteristics of Sandy Grassland Soils under Post-grazing Natural
2329 Restoration in Horqin Sandy Land. *Journal of Desert Research* 22(4):333-338.
- 2330 Szott LT, Palm CA (1996) Nutrient stocks in managed and natural humid tropical fallows. *Plant and Soil*, 186, 293-
2331 309.
- 2332 Templer PH, Groffman PM, Flecker AS & Power AG (2005) Land use change and soil nutrient transformations in the
2333 Los Haitises region of the Dominican Republic. *Soil Biology & Biochemistry* 37:215-225.

First Order Draft

- 2334 Tian H, Zhou D, Guo P (2001b) The change of soil and vegetation with different years of leaving uncultivated. *Journal*
2335 *of Northeast Normal University* 33(4):72-77.
- 2336 Tian J, Zhou Z, Bao B, Sun J (2008) Variations of soil particle size distribution with land-use types and influences on
2337 soil organic carbon and nitrogen. *Journal of Plant Ecology (Chinese version)* 32(3):601-610.
- 2338 Tiessen H, Salcedo IH, Sampaio E (1992) Nutrient and soil organic matter dynamics under shifting cultivation in
2339 semiarid northeastern Brazil. *Agriculture Ecosystems & Environment*, 38, 139-151.
- 2340 Tiessen H, Stewart JWB & Bettany JR (1982) Cultivation effects on the amounts and concentrations of carbon,
2341 nitrogen, and phosphorus in grassland soils. *Agron J* 74:831-835.
- 2342 Tornquist CG, Hons FM, Feagley SE & Haggard J (1999) Agroforestry system effects on soil characteristics of the
2343 Sarapiquí region of Costa Rica. *Agriculture Ecosystems & Environment* 73:19-28.
- 2344 Townsend AR, Vitousek PM & Trumbore SE (1995) Soil organic matter dynamics along gradients in temperature and
2345 land use on the island of Hawaii. *Ecology* 76:721-733.
- 2346 Trouve C, Mariotti A, Schwartz D, Guillet B (1994) Soil Organic-Carbon Dynamics under Eucalyptus and Pinus
2347 Planted on Savannas in the Congo. *Soil Biology & Biochemistry*, 26, 287-295.
- 2348 Trumbore SE, Davidson EA, Decamargo PB, Nepstad DC, Martinelli LA (1995) Belowground Cycling of Carbon in
2349 Forests and Pastures of Eastern Amazonia. *Global Biogeochemical Cycles*, 9, 515-528.
- 2350 Uhl C, Jordan CF (1984) Succession and nutrient dynamics following forest cutting and burning in Amazonia. *Ecology*,
2351 65, 1476-1490.
- 2352 Unger PW (2001) Total carbon, aggregation, bulk density, and penetration resistance of cropland and nearby grassland
2353 soils. In: R. Lal (Ed) *Soil carbon sequestration and the greenhouse effect* (pp. 77-92). SSSA Special Publication No.
2354 57, Madison, WI.
- 2355 Vagen TG, Walsh MG, Shepherd KD (2006) Stable isotopes for characterisation of trends in soil carbon following
2356 deforestation and land use change in the highlands of Madagascar. *Geoderma*, 135, 133-139.
- 2357 van Dam D, Veldkamp E & van Breemen N (1997) Soil organic carbon dynamics: variability with depth in forested
2358 and deforested soils under pasture in Costa Rica. *Biogeochemistry* 39:343-375.
- 2359 van Noordwijk M, Cerri C, Woomer PL, Nugroho K, Bernoux M (1997) Soil carbon dynamics in the humid tropical
2360 forest zone. *Geoderma*, 79, 187-225.
- 2361 van Straaten O, Corre MD, Wolf K, Tchienkoua M, Cuellar E, Matthews RB & Veldkamp E (2015) Conversion of
2362 lowland tropical forests to tree cash crop plantations loses up to one-half of stored soil organic carbon. *PNAS*
2363 112:9956-9960.
- 2364 Veldkamp E (1994) Organic-Carbon Turnover in 3 Tropical Soils under Pasture after Deforestation. *Soil Sci Soc Am*
2365 J 58:175-180.
- 2366 Veldkamp E, Becker A, Schwendenmann L, Clark DA & Schulte-Bisping H (2003) Substantial labile carbon stocks
2367 and microbial activity in deeply weathered soils below a tropical wet forest. *Global Change Biology* 9:1171-1184.
- 2368 Villarino SH, Studdert GA, Baldassini P, Cendoya MG, Ciuffoli L, Mastrangelo M & Pineiro G (2017) Deforestation
2369 impacts on soil organic carbon stocks in the Semiarid Chaco Region, Argentina. *SciTotal Env* 575:1056-1065.
- 2370 Voroney RP, Van Veen JA & Paul EA (1981) Organic C dynamics in grassland soils. 2. Model validation and
2371 simulation of the long-term effects of cultivation and rainfall erosion. *Can J Soil* 61:211-224.
- 2372 Wadsworth G, Southard RJ, Singer MJ (1988) Effects of Fallow Length on Organic-Carbon and Soil Fabric of Some
2373 Tropical Uduits. *Soil Science Society of America Journal*, 52, 1424-1430.
- 2374 Wairu M & Lal R (2003) Soil organic carbon in relation to cultivation and topsoil removal on sloping lands of
2375 Kolombangara, Solomon Islands. *Soil & Tillage Research* 70:19-27.
- 2376 Walker SM & Desanker PV (2004) The impact of land use on soil carbon in Miombo woodlands of Malawi. *Forest*
2377 *Ecology & Management* 203:345-360.

- 2378 Wang G, Haiyan M, Ju Q, Juan C (2004) Impact of land use changes on soil carbon, nitrogen and phosphorus and
2379 water pollution in an arid region of northwest China. *Soil Use and Management* 20:32-39.
- 2380 Wang JM, Zhang XC (2009) Changes of carbon storage in vegetation and soil during different successional stages of
2381 rehabilitated grassland. *Acta Prataculturae Sinica* 18(1):1-8.
- 2382 Wang S, Wilkes A, Zhang Z, Chang X, Lang R, Wang Y, Niu H (2011) Management and land use change effects on
2383 soil carbon in northern China's grasslands: a synthesis. *Agriculture, Ecosystems and Environment* 142:329-340.
- 2384 Wang WY, Wang QJ, Wang CY, Shi HL, Wang G (2005) The effect of land management on carbon and nitrogen
2385 status in plants and soils of alpine meadows on the Tibetan plateau. *Land Degradation & Development* 16:405-415.
- 2386 Wang WY Wang Qiji, Wang G (2006) Effects of land degradation and rehabilitation on soil carbon and nitrogen
2387 content on alpine Kobersia meadow. *Ecology and Environment* 15:362-366.
- 2388 Wang X, Liu J, Zhang X, Lei RX, Lai Y (2007) Effects of Landuse Change on Soil Nutrients and Enzyme Activities
2389 and Their Correlations in Semiarid Area of the Loess Plateau. *Bulletin of Soil and Water Conservation* 27(6):50-56.
- 2390 Wang Z, Gao B, Li X (2006) Effects of different land use types on carbonhydrate content and aggregate stability in
2391 arid grassland soil. *Journal of Gansu Agricultural University* 41(3):91-95.
- 2392 Wang Z, Han X, Li L (2008) Effects of grassland conversion to croplands on soil organic carbon in the temperate
2393 Inner Mongolia. *Journal of Environmental Management* 86:529-534.
- 2394 Weaver PL, Birdsey RA, Lugo AE (1987) Soil organic matter in secondary forests of Puerto Rico. *Biotropica*, 19, 17-
2395 23.
- 2396 Wick B, Tiessen H, Menezes RSC (2000) Land quality changes following the conversion of the natural vegetation
2397 into silvo-pastoral systems in semi-arid NE Brazil. *Plant and Soil*, 222, 59-70.
- 2398 Wick B, Veldkamp E, de Mello WZ, Keller M, Crill P (2005) Nitrous oxide fluxes and nitrogen cycling along a pasture
2399 chronosequence in Central Amazonia, Brazil. *Biogeosciences* 2:175-187.
- 2400 Wu X, Zhang L, Ding Y, Wang Q, Lu H, Wang X (2006) Effect of Land Use on Soil Properties in Inter-distributing
2401 Area of Farming and Pasturing of Keerqin Sandy Land. *Journal of Soil and Water Conservation* 20(4):116-119.
- 2402 Yan Y, Tang H, Chang R, Liu L (2008) Variation of Below-Ground Carbon Sequestration Under Long Term
2403 Cultivation and Grazing in the Typical Steppe of Nei Monggol in North China. *Environmental Science* 29(5):1388-
2404 1393.
- 2405 Yang X, Blagodatsky S, Lippe M, Liu F, Hammond J, Xu J & Cadisch G (2016) Land-use change impact on time-
2406 averaged carbon balances: rubber expansion and reforestation in a biosphere reserve, south-west China. *For Ecol &*
2407 *Mgmt* 372:149-163.
- 2408 Yang JC, Huang JH, Pan QM, Tang JW, Han XG (2004) Long-term impacts of land-use change on dynamics of
2409 tropical soil carbon and nitrogen pools. *Journal of Environmental Sciences-China*, 16, 256-261.
- 2410 Yemefack M, Rossiter DG, Jetten VG (2006) Empirical modelling of soil dynamics along a chronosequence of shifting
2411 cultivation systems in southern Cameroon. *Geoderma*, 133, 380-397.
- 2412 Yin P, Li X, Li Y, Zhang P (2008) Changes in soil organic C and N and aggregate stability along a chronosequence
2413 of Cultivation in sub-alpine grassland. *Journal of Gansu Agricultural University* 43(4):97-102.
- 2414 Yonekura Y, Ohta S, Kiyono Y, Aksa D, Morisada K, Tanaka N & Kanzaki M (2010) Changes in soil carbon stock
2415 after deforestation and subsequent establishment of "Imperata" grassland in the Asian humid tropics. *Plant & Soil*
2416 329:495-507.
- 2417 Yu W, Ma Q, Zhao X, Zhou H, Li J (2007) Changes of soil active organic carbon pool under different land use types.
2418 *Chinese Journal of Ecology* 26 (12):2013- 2016.
- 2419 Yue Q, Chang Q, Liu J, Liu M, Wang D (2007) Effect of different land utilization on soil nutrient and soil enzyme in
2420 Loess Plateau. *Journal of Northwest A & F University (Nat. Sci. Ed.)* 35(12):103-108.
- 2421 Zhan Z, Li X, Zhang D, Wang Z (2005) Effects of land use on organic C concentration and structural properties in
2422 alpine grassland soil. *Acta Pedologica Sinica* 42(5):777-782.

First Order Draft

- 2423 Zhang H, Thompson ML & Sandor JA (1988) Compositional differences in organic matter among cultivated and
2424 uncultivated Argiudolls and Hapludalfs derived from loess. *Soil Sci Soc Am J* 52:216-222.
- 2425 Zhao W, Xiao H, Liu Z, et al (2005) Soil degradation and restoration as affected by land use change in the semiarid
2426 Bashang area, northern China. *Catena* 59:173-186.
- 2427 Zhou Z, Sun OJ, Huang J (2007) Soil carbon and nitrogen stores and storage potential as affected by land-use in an
2428 agro-pastoral ecotone of northern China. *Biogeochemistry* 82:127-138.
- 2429 Zinn YL, Lal R, Resck DVS (2005) Changes in soil organic carbon stocks under agriculture in Brazil. *Soil &*
2430 *Tillage Research*, **84**, 28-40.
- 2431 Zinn YL, Resck DVS & da Silva JE (2002) Soil organic carbon as affected by afforestation with Eucalyptus and Pinus
2432 in the Cerrado region of Brazil. *Forest Ecology & Management* 166:285-294.
- 2433 Zingore S, Manyame C, Nyamugafata P, Giller KE (2005) Long-term changes in organic matter of
2434 woodland soils cleared for arable cropping in Zimbabwe. *European Journal of Soil Science*, **56**, 727-736.
- 2435 Zou X, Bashkin M (1998) Soil carbon accretion and earthworm recovery following revegetation in
2436 abandoned sugarcane fields. *Soil Biology & Biochemistry*, **30**, 825-830.
- 2437
- 2438 **Sources of Data for Tillage Management Factor**
- 2439 Ahl C, Joergensen RG, Kandeler E, Meyer B & Woehler V (1998) Microbial biomass and activity in silt and sand
2440 loams after long-term shallow tillage in central Germany. *Soil Tillage Res* 49:93-104.
- 2441 Al-Kaisi M, Yin X, Licht M (2005) Soil carbon and nitrogen changes as influenced by tillage and cropping systems
2442 in some Iowa soils. *Agriculture, Ecosystems & Environment* 105,4:635-647.
- 2443 Al-Kaisi M, Yin X, Licht M (2005) Soil carbon and nitrogen changes as affected by tillage system and crop biomass
2444 in a corn-soybean rotation. *Applied Soil Ecology* 30,3:174-191.
- 2445 Alvarez R, Russo ME, Prystupa P, Scheiner JD & Blotta L (1998) Soil carbon pools under conventional and no-
2446 tillage systems in the Argentine Rolling Pampa. *Agron J* 90:138-143.
- 2447 Alvaro-Fuentes J, Cantero-Martinez C, Lopez MV, Paustian K, Deneff K, Stewart CE, Arrue JL (2009) Soil
2448 aggregation and soil organic carbon stabilization: Effects of management in semiarid Mediterranean agroecosystems.
2449 *Soil Science Society of America Journal* 73,5:1519-1529.
- 2450 Alvaro-Fuentes J, Lopez MV, Cantero-Martinez C, Arrue JL (2008) Tillage effects on soil organic carbon fractions in
2451 Mediterranean dryland agroecosystems. *Soil Science Society of America Journal* 72,2:541-547.
- 2452 Alvaro-Fuentes J, Plaza-Bonilla D, Arrue JL, Lampurlanes J, Cantero-Martinez C (2014) Soil organic carbon storage
2453 in a no-tillage chronosequence under Mediterranean conditions. *Plant Soil* 376:31-41.
- 2454 Angers DA, Bolinder MA, Carter MR, Gregorich EG, Drury CF, Liang BC, Voroney RP, Simard RR, Donald RG,
2455 Beyaert RP & Martel J (1997) Impact of tillage practices on organic carbon and nitrogen storage in cool, humid soils
2456 of eastern Canada. *Soil Tillage Res* 41:191-201.
- 2457 Angers DA, Voroney RP & Côté D (1995) Dynamics of soil organic matter and corn residues affected by tillage
2458 practices. *Soil Sci Soc Am J* 59:1311-1315.
- 2459 Anken T, Weisskopf P, Zihlmann U, Forrer H, Jansa J, Perhacova K (2004) Long-term tillage system effects under
2460 moist cool conditions in Switzerland. *Soil and Tillage Research* 78,2: 171-183.
- 2461 Balesdent J, Mariotti A & Boissgonnier D (1990) Effect of tillage on soil organic carbon mineralization estimated from
2462 ^{13}C abundance in maize fields. *J Soil Sci* 41:587-596.
- 2463 Barber RG, Orellana M, Navarro F, Diaz O & Soruco MA (1996) Effects of conservation and conventional tillage
2464 systems after land clearing on soil properties and crop yield in Santa Cruz, Bolivia. *Soil Tillage Res* 38:133-152.
- 2465 Bayer C, Martin-Neto L, Mielniczuk J, Pavinato A, Dieckow J (2006) Carbon sequestration in two Brazilian Cerrado
2466 soils under no-till. *Soil and Tillage Research* 86,2:237-245.

- 2467 Bayer C, Mielniczuk J, Amado TJC, Martin-Neto L & Fernandes SV (2000) Organic matter storage in a sandy clay
2468 loam Acrisol affected by tillage and cropping systems in southern Brazil. *Soil Tillage Res* 54:101-109.
- 2469 Bayer C, Mielniczuk J, Martin-Neto L & Ernani PR (2002) Stocks and humification degree of organic matter fractions
2470 as affected by no-tillage on a subtropical soil. *Pl Soil* 238:133-140.
- 2471 Beare MH, Hendrix PF & Coleman DC (1994) Water-stable aggregates and organic matter fractions in conventional-
2472 and no-tillage soils. *Soil Sci Soc Am J* 58:777-786.
- 2473 Bhattacharyya R, Kundu S, Pandey SC, Singh KP, Gupta HS (2008) Tillage and irrigation effects on crop yields and
2474 soil properties under the rice-wheat system in the Indian Himalayas. *Agricultural Water Management* 95,9:993-1002.
- 2475 Bhattacharyya R, Pandey SC, Bisht JK, Bhatt JC, Gupta HS, Tuti MD, Mahanta D, Mina BL, Singh RD, Chandra S,
2476 Srivastva AK, Kundu S (2013) Tillage and irrigation effects on soil aggregation and carbon pools in the Indian Sub-
2477 Himalayas. *Agron J* 105:101-112.
- 2478 Bhattacharyya R, Prakash V, Kundu S, Srivastva AK, Gupta HS (2009) Soil aggregation and organic matter in a sandy
2479 clay loam soil of the Indian Himalayas under different tillage and crop regimes. *Agriculture, Ecosystems, and*
2480 *Environment* 132,1-2:126-134.
- 2481 Black AL & Tanaka DL (1997) A conservation tillage-cropping systems study in the northern Great Plains of the
2482 United States. In: Paul EA, Elliot ET, Paustian K, Cole CV (Eds) *Soil organic matter in temperate agroecosystems:*
2483 *Long-term experiments in North America* (pp. 335-342). CRC Press, Boca Raton, FL.
- 2484 Blanco-Canqui H, Gantzer CJ, Anderson SH, Alberts EE (2004) Tillage and crop influences on physical properties
2485 for an Epiaqualf. *Soil Sci Soc Am J* 68:567-576.
- 2486 Blanco-Canqui H, Schlegel AJ, Heer WF (2011) Soil-profile distribution of carbon and associated properties in no-
2487 till along a precipitation gradient in the central Great Plains. *Agriculture, Ecosystems and Environment* 144,1:107-
2488 116.
- 2489 Boddey R, Jantalia C, Conceicao P, Zanatta J, Bayer C, Mielniczuk J, Dieckow J, Dos Santos H, Denardins J, Aita
2490 C, Giacomini S, Alves B, Urquiaga S (2010) Carbon accumulation at depth in Ferralsols under zero-till subtropical
2491 agriculture. *Global Change Biology* 16:748-795.
- 2492 Bordovsky DG, Choudhary M, & Gerard CJ (1999) Effect of tillage, cropping, and residue management on soil
2493 properties in the Texas rolling plains. *Soil Sci* 164:331-340.
- 2494 Borin M, Menini C & Sartori L (1997) Effects of tillage systems on energy and carbon balance in north-eastern Italy.
2495 *Soil Tillage Res* 40:209-226.
- 2496 Borresen T & Njos A (1993) Ploughing and rotary cultivation for cereal production in a long-term experiment on a
2497 clay soil in southeastern Norway. 1. Soil properties. *Soil Tillage Res* 28:97-108.
- 2498 Bowman RA & Anderson RL (2002) Conservation Reserve Program: Effects on soil organic carbon and preservation
2499 when converting back to cropland in northeastern Colorado. *J Soil & Water Cons* 57:121-126.
- 2500 Burch GJ, Mason IB, Fischer RA, Moore ID (1986) *Australian Journal of Soil Research* 24:377-391.
- 2501 Buschiazzi DE, Panigatti JL & Unger PW (1998) Tillage effects on soil properties and crop production in the
2502 subhumid and semiarid Argentinean Pampas. *Soil Tillage Res* 49:105-116.
- 2503 Buyanovsky GA & Wagner GH (1998) Carbon cycling in cultivated land and its global significance. *Glob Change*
2504 *Biol* 4:131-141.
- 2505 Calegari A, Hargrove WL, Rheinheimer D, Ralisch R, Tessier D, de Tourdonnet S, Guimaraes MdF (2008) Impact of
2506 long-term no-tillage and cropping system management on soil organic carbon in an Oxisol: A model for sustainability.
2507 *Agron J* 100:1013-1019.
- 2508 Campbell CA, Biederbeck VO, McConkey BG, Curtin D & Zentner RP (1999) Soil quality-Effect of tillage and
2509 fallow frequency. Soil organic matter quality as influenced by tillage and fallow frequency in a silt loam in
2510 southwestern Saskatchewan. *Soil Biol and Biochem* 31:1-7.
- 2511 Campbell CA, McConkey BG, Zentner RP, Selles F & Curtin D (1996) Long-term effects of tillage and crop rotations
2512 on soil organic C and total N in a clay soil in southwestern Saskatchewan. *Can J Soil* 76:395-401.

First Order Draft

- 2513 Carter MR (1991) Evaluation of shallow tillage for spring cereals on a fine sandy loam. 2. Soil physical, chemical and
2514 biological properties. *Soil Tillage Res* 21:37-52.
- 2515 Carter MR, Johnston HW & Kimpinski J (1988) Direct drilling and soil loosening for spring cereals on a fine sandy
2516 loam in Atlantic Canada. *Soil Tillage Res* 12:365-384.
- 2517 Carter MR, Mele PM, Steed GR (1994) The effects of direct drilling and stubble retention on water and bromide
2518 movement and earthworm species in a duplex soil. *Soil Science* 157:224-231.
- 2519 Carter MR, Sanderson JB, Ivany JA & White RP (2002) Influence of rotation and tillage on forage maize productivity,
2520 weed species, and soil quality of a fine sandy loam in the cool-humid climate of Atlantic Canada. *Soil Tillage Res*
2521 67:85-98.
- 2522 Cavanagh PP, Koppi AJ, McBratney AB (1991) The effects of minimum cultivation after three years on some physical
2523 and chemical properties of a red-brown earth at Forbes, N.S.W. *Australian Journal of Soil Research* 29:263-270.
- 2524 Chan KY & Mead JA (1988) Surface physical properties of a sandy loam soil under different tillage practices. *Aust J*
2525 *Soil Res* 26:549-559.
- 2526 Chan KY, Roberts WP & Heenan DP (1992) Organic carbon and associated soil properties of a red Earth after 10
2527 years of rotation under different stubble and tillage practices. *Aust J Soil Res* 30:71-83.
- 2528 Chan KY, Heenan DP, Oates A (2002) Soil carbon fractions and relationships to soil quality under different tillage
2529 and stubble management. *Soil & Tillage Research* 63:133-139.
- 2530 Chan KY, Heenan DP, So HB (2003) Sequestration of carbon and changes in soil quality under conservation tillage
2531 on light-textured soils in Australia: a review. *Australian Journal of Experimental Agriculture* 43:325-334.
- 2532 Chaney BK, Hodson DR & Braim MA (1985) The effects of direct drilling, shallow cultivation and ploughing on
2533 some soil physical properties in a long-term experiment on spring barley. *J Agric Sci (Camb)* 104:125-133.
- 2534 Chen H, Hou R, Gong Y, Li H, Fan M, Kuzyakov Y (2009) Effects of 11 years of conservation tillage on soil organic
2535 matter fractions in wheat monoculture in Loess Plateau of China. *Soil and Tillage Research* 106,1:85-94.
- 2536 Chen H, Marhan S, Billen N, Stahr K (2009) Soil organic-carbon and total nitrogen stocks as affected by different
2537 land uses in Baden-Wurttemberg (southwest Germany). *Journal of Plant Nutrition and Soil Science* 172:32-42.
- 2538 Chen Z, Dikgwatlhe SB, Xue J, Zhang H, Chen F, Xiao X (2015) Tillage impacts on net carbon flux in paddy soil of
2539 the Southern China. *Journal of Cleaner Production* 103,15:70-76.
- 2540 Cheng-Fang L, Dan-Na Z, Zhi-Kui K, Zhi-Sheng Z, Jin-Ping W, Ming-Li C, Cou-Gui C (2012) Effects of tillage and
2541 nitrogen fertilizers on CH₄ and CO₂ emissions and soil organic carbon in paddy fields of Central China. *PLoS ONE*
2542 7,5:34642.
- 2543 Choudhary VK, Kumar PS, Bhagawati R (2013) Response of tillage and in situ moisture conservation on alteration
2544 of soil and morpho-physiological differences in maize under Eastern Himalayan region of India. *Soil and Tillage*
2545 *Research* 134:41-48.
- 2546 Clapp CE, Allmaras RR, Layese MF, Linden DR & Dowdy RH (2000) Soil organic carbon and ¹³C abundance as
2547 related to tillage, crop residue, and nitrogen fertilization under continuous corn management in Minnesota. *Soil Tillage*
2548 *Res* 55:127-142.
- 2549 Corazza EJ, Silva JE, Resck DVS, Gomes AC, (1999). Behavior of different management systems as a source or sink
2550 of C-CO₂ in relation to cerrado type vegetation. *R.Bras Ci.Solo* 23:425-432.
- 2551 Costantini A, Cosentino D & Segat A (1996) Influence of tillage systems on biological properties of a Typic Argiudoll
2552 soil under continuous maize in central Argentina. *Soil Tillage Res* 38:265-271.
- 2553 Dalal RC (1989) Long-term effects of no-tillage, crop residue, and nitrogen application on properties of a Vertisol.
2554 *Soil Sci Soc Am J* 53:1511-1515.

- 2555 Dalal RC, Henderson PA & Glasby JM (1991) Organic matter and microbial biomass in a vertisol after 20 yr of
2556 zero tillage. *Soil Biol and Biochem* 23:435-441.
- 2557 Denef K, Zotarelli L, Boddey R, Six J (2007) Microaggregate-associated carbon as a diagnostic fraction for
2558 management-induced changes in soil organic carbon in two Oxisols. *Soil Biology and Biochemistry* 39,5:1165-
2559 1172.
- 2560 Devine S, Markewitz D, Hendrix P, Coleman D (2014) Soil aggregates and associated organic matter under
2561 conventional tillage, no-tillage, and forest succession after three decades. *PLoS ONE* 9:1.
- 2562 Dick WA & Durkalski JT (1997) No-tillage production agriculture and carbon sequestration in a typic fragiudalf
2563 soil of northeastern Ohio. In: Lal R, Kimble JM, Follett RF, Stewart BA (Eds) *Advances in Soil Science: Management of Carbon Sequestration in Soil* (pp. 59-71). CRC Press, Boca Raton, FL.
- 2565 Dick WA & Durkalski JT (1997) No-tillage production agriculture and carbon sequestration in a typic fragiudalf
2566 soil of northeastern Ohio. In: Lal R, Kimble JM, Follett RF, Stewart BA (Eds) *Advances in Soil Science: Management of Carbon Sequestration in Soil* (pp. 59-71). CRC Press, Boca Raton, FL.
- 2568 Dikgwatlhe SB, Chen Z, Lal R, Zhang H, Chen R (2014) Changes in soil organic carbon and nitrogen as affected
2569 by tillage and residue management under wheat-maize cropping system in the North China Plain. *Soil and Tillage Research* 144:110-118.
- 2571 Dimassi B, Mary B, Wylleman R, Labreuche J, Couture D, Piraux F, Cohan J-P (2014) Long-term effect of
2572 contrasted tillage and crop management on soil carbon dynamics during 41 years. *Agriculture, Ecosystems and Environment* 188:134-146.
- 2574 Dolan MS, Clapp CE, Allmaras RR, Baker JM, Molina JAE (2006) Soil organic carbon and nitrogen in a
2575 Minnesota soil as related to tillage, residue and nitrogen management. *Soil and Tillage Research* 89,2: 221-231.
- 2576 Doran JW, Elliott ET & Paustian K (1998) Soil microbial activity, nitrogen cycling, and long-term changes in
2577 organic carbon pools as related to fallow tillage management. *Soil Tillage Res* 49:3-18.
- 2578 Dou F, Wright A, Hons F (2008) Sensitivity of labile soil organic carbon to tillage in wheat-based cropping systems. *Soil Sci. Soc. Am. J.* 72:1445-1453.
- 2580 Du Z, Ren T & Hu C (2010) Tillage and residue removal effects on soil carbon and nitrogen storage in the North
2581 China Plain. *Soil Sci. Soc. Am. J.* 74:197-202
- 2582 Du Z, Ren T, Hu C, Zhang Q (2015) Transition from intensive tillage to no-till enhances carbon sequestration in
2583 microaggregates of surface soil in the North China Plain. *Soil & Tillage Research* 146:26-31.
- 2584 Duiker SW & Lal R (1999) Crop residue and tillage effects on carbon sequestration in a Luvisol in central Ohio. *Soil Tillage Res* 52:73-81.
- 2586 Edwards JH, Woods CW, Thurlow DL & Ruf ME (1992) Tillage and crop rotation effects on fertility status of a
2587 Hapludult soil. *Soil Sci Soc Am J* 56:1577-1582.
- 2588 Eghball B, Mielke LN, McCallister DL & Doran JW (1994) Distribution of organic carbon and inorganic nitrogen
2589 in a soil under various tillage and crop sequences. *J Soil & Water Cons* 49:201-205.
- 2590 Fabrizzi KP, Moron A, Garcia FO (2003) Soil carbon and nitrogen organic fractions in degraded vs. non-degraded
2591 Mollisols in Argentina. *Soil Sci Soc Am J* 67:1831-1841.
- 2592 Fabrizzi KP, Rice CW, Amado TJC, Fiorin J, Barbagelata P, Melchiori T (2009) Protection of soil organic C and
2593 N in temperate and tropical soils: effect of native and agroecosystems. *Biogeochemistry* 92,1-2:129-143.
- 2594 Fan RQ, Yang XM, Drury CF, Reynolds WD, Zhang XP (2014) Spatial distributions of soil chemical and physical
2595 properties prior to planting soybean in soil under ridge-, no- and conventional-tillage in a maize-soybean rotation. *Soil Use and Management* 30:414-422.
- 2597 Feiziene D, Feiza V, Slepetiene A, Liaudanskiene I, Kadziene G, Deveikyte I, Vaideliene A (2011) Long-term
2598 influence of tillage and fertilization on net carbon dioxide exchange rate on two soils with different textures. *Journal of Environmental Quality* 40:1787-1796.
- 2600 Fettell N, Gill H (1985) Long-term effects of tillage, stubble and nitrogen management on properties of a red-
2601 brown earth. *Australian Journal of Experimental Agriculture* 35:923-928.
- 2602 Fleige H & Baeumer K (1974) Effect of zero-tillage on organic carbon and total nitrogen content, and their
2603 distribution in different N-fractions in loessial soils. *Agro-Ecosyst* 1:19-29.
- 2604 Follett RF & Peterson GA (1988) Surface soil nutrient distribution as affected by wheat-fallow tillage systems. *Soil Sci Soc Am J* 52:141-147.

Ver. 20170605

- 2606 Franzluebbers AJ & Arshad MA (1996) Water-stable aggregation and organic matter in four soils under
2607 conventional and zero tillage. *Can J Soil* 76:387-393. Franzluebbers AJ, Hons FM & Zuberer DA (1995) Soil
2608 organic carbon, microbial biomass, and mineralizable carbon and nitrogen in sorghum. *Soil Sci Soc Am J* 59:460-
2609 466.
- 2610 Franzluebbers AJ, Langdale GW & Schomberg HH (1999) Soil carbon, nitrogen, and aggregation in response to
2611 type and frequency of tillage. *Soil Sci Soc Am J* 63:349-355.
- 2612 Franzluebbers AJ, Stuedemann JA (2002) Particulate and non-particulate fractions of soil Organic carbon under
2613 pastures in the Southern Piedmont USA. *Environmental Pollution* 116,1:53-62.
- 2614 Freitas PL, Blancaneaux P, Gavinelly E, Larre-Larrouy M-C & Feller C (2000) Nivel e natureza do estoque
2615 organico de latossols sob diferentes sistemas de uso e manejo, *Pesq.agropec.bras. Brasilia* 35: 157-170.
- 2616 Freixo AA, Machado P, dos Santos HP, Silva CA & Fadigas F (2002) Soil organic carbon and fractions of a Rhodic
2617 Ferralsol under the influence of tillage and crop rotation systems in southern Brazil. *Soil Tillage Res* 64:221-230.
- 2618 Gál A, Vyn TJ, Micheli E, Kladvik EJ, McFee WW (2007) Soil carbon and nitrogen accumulation with long-
2619 term no-till versus moldboard plowing overestimated with tilled-zone sampling depths. *Soil and Tillage Research*
2620 96,1-2:42-51.
- 2621 Garcia-Prechac F, Ernst O, Siri-Prieto G, Terra JA (2004) Integrating no-till into crop-pasture rotations in Uruguay.
2622 *Soil and Tillage Research* 77,1:1-13.
- 2623 Ghimire R, Adhikari K, Chen Z, Shah S, Dahal K (2012) Soil organic carbon sequestration as affected by tillage,
2624 crop residue, and nitrogen application in rice-wheat rotation system. *Paddy and Water Environment* 10,2:95-102.
- 2625 Ghuman BS & Sur HS (2001) Tillage and residue management effects on soil properties and yields of rainfed
2626 maize and wheat in a subhumid subtropical climate. *Soil Tillage Res* 58:1-10.
- 2627 Grabski AS, So HB, Schafer BM, Desborough PJ (1997) The effects of tillage on the properties and crop yields of
2628 a loam soil at Grafton, NSW, Australia. In 'Proceedings of the 14th ISTRO conference'. 27 July- 1 August 1997,
2629 Pulawy, Poland. Pp. 247-254.
- 2630 Green VS, Stott DE, Cruz JC, Curi N (2007) Tillage impacts on soil biological activity and aggregation in a
2631 Brazilian Cerrado Oxisol. *Soil and Tillage Research* 92,1-2:114-121.
- 2632 Gwenzi W, Gotosa J, Chakanetsa S, Mutema Z (2009) Effects of tillage systems on soil organic carbon dynamics,
2633 structural stability and crop yields in irrigated wheat (*Triticum aestivum* L.)-cotton (*Gossypium hirsutum* L.)
2634 rotation in semi-arid Zimbabwe. *Nutrient Cycling in Agroecosystems* 83:211.
- 2635 Halvorson AD, Vigil MF, Peterson GA & Elliott ET (1997) Long-term tillage and crop residue management study
2636 at Akron, Colorado. In: Paul EA, Elliot ET, Paustian K, Cole CV (Eds) *Soil organic matter in temperate*
2637 *agroecosystems: Long-term experiments in North America* (pp. 361-370). CRC Press, Boca Raton, FL.
- 2638 Halvorson AD, Wienhold BJ & Black AL (2002) Tillage, nitrogen, and cropping system effects on soil carbon
2639 sequestration. *Soil Sci Soc Am J* 66:906-912.
- 2640 Hansmeyer TL, Linden DR, Allan DL & Huggins DR (1997) Determining carbon dynamics under no-till, ridge-
2641 till, chisel, and moldboard tillage systems within a corn and soybean cropping sequence. In: Lal R, Kimble JM,
2642 Follett RF, Stewart BA (Eds) *Advances in Soil Science: Management of Carbon Sequestration in Soil* (pp. 93-97).
2643 CRC Press, Boca Raton, FL.
- 2644 Hao X, Chang C & Lindwall CW (2001) Tillage and crop sequence effects on organic carbon and total nitrogen
2645 content in an irrigated Alberta soil. *Soil Tillage Res* 62:167-169.
- 2646 Havlin JL & Kissel DE (1997) Management effects on soil organic carbon and nitrogen in the east-central Great
2647 Plains of Kansas. In: Paul EA, Elliot ET, Paustian K, Cole CV (Eds) *Soil organic matter in temperate*
2648 *agroecosystems: Long-term experiments in North America* (pp. 381-386). CRC Press, Boca Raton, FL.
- 2649 Heenan DP, McGhies WJ, Thomson FM, Chan KY (1995) Decline in soil organic carbon and total nitrogen in
2650 relation to tillage stubble management and rotation. *Australian Journal of Experimental Agriculture* 35:877-884.
- 2651 Heinze S, Rauber R, Joergensen RG (2010) Influence of mouldboard plough and rotary harrow tillage on microbial
2652 biomass and nutrient stocks in two long-term experiments on loess derived Luvisols. *Applied Soil Ecology* 46:405-
2653 412.
- 2654 Hendrix PF (1997) Long-term patterns of plant production and soil carbon dynamics in a Georgia piedmont
2655 agroecosystem. In: Paul EA, Elliot ET, Paustian K, Cole CV (Eds) *Soil organic matter in temperate*
2656 *agroecosystems: Long-term experiments in North America* (pp. 235-245). CRC Press, Boca Raton, FL.

- 2657 Hermle S, Anken T, Leifeld J, Weiskopf P (2008) The effect of the tillage system on soil organic carbon content
2658 under moist, cold-temperate conditions. *Soil and Tillage Research* 98,1:94-105.
- 2659 Hernanz JL, Lopez R, Navarrete L & Sanchez-Giron V (2002) Long-term effects of tillage systems and rotations
2660 on soil structural stability and organic carbon stratification in semiarid central Spain. *Soil Tillage Res* 66:129-141.
- 2661 Hernanz JL, Sanchez-Giron V, Navarrete L (2009) Soil carbon sequestration and stratification in a
2662 cereal/leguminous crop rotation with three tillage systems in semiarid conditions. *Agriculture, Ecosystems, and*
2663 *Environment* 133,1-2:114-122.
- 2664 Higashi T, Yunghui M, Komatsuzaki M, Miura S, Hirata T, Araki H, Kaneko N, Ohta H (2014) Tillage and cover
2665 crop species affect soil organic carbon in Andosol, Kanto, Japan. *Soil and Tillage Research* 138:64-72.
- 2666 Hou R, Ouyang Z, Li Y, Tyler D, Li F, Wilson G (2011) Effects of tillage and residue management on soil organic
2667 carbon and total nitrogen in the North China Plain. *Soil Science Society of America Journal* 76,1:230-240.
- 2668 Huggins DR, Allmaras RR, Clapp CE, Lamb JA, Randall GW, (2007) Corn-Soybean sequence and tillage effects
2669 on soil carbon dynamics and storage. *Soil Science Society of America Journal* 71,1:145-154.
- 2670 Hulugalle NR (2000) Carbon sequestration in irrigated vertisols under cotton-based farming systems. *Comm Soil*
2671 *Sci Plant Anal* 31:645-654.
- 2672 Hussain I, Olson KR, Wander MM & Karlen DL (1999) Adaption of soil quality indices and application to three
2673 tillage systems in southern Illinois. *Soil Tillage Res* 50:237-249.
- 2674 Ismail I, Blevins RL, Frye WW (1994) Long-term no-tillage effects on soil properties and continuous corn yields.
2675 *Soil Science Society of America Journal* 58,1:193-198.
- 2676 Jagadamma S, Lal R (2010) Distribution of organic carbon in physical fractions of soils as affected by agricultural
2677 management. *Biology and Fertility of Soils* 46,6:543-554.
- 2678 Jarecki MK, Lal R (2010) Crop management for soil carbon sequestration. *Critical Reviews in Plant Sciences*
2679 22,6:471-502.
- 2680 Jarvis R (1996) Nineteen years of no-till – the effects on soil properties and crop yield. In ‘Proceedings of no-till
2681 conference’. Katanning, WA. Pp. 20-25. Department of Agriculture Western Australia: Katanning, WA.
- 2682 Jemai I, Aissa NB, Guirat SB, Ben-Hammouda M, Gallali T (2012) On-farm assessment of tillage impact on the
2683 vertical distribution of soil organic carbon and structural soil properties in a semiarid region in Tunisia. *Journal of*
2684 *Environmental Management* 113:488-494.
- 2685 Jemai I, Aissa NB, Guirat SB, Ben-Hammouda M, Gallali T (2013) Impact of three and seven years of no-tillage
2686 on the soil water storage, in the plant root zone, under a dry subhumid Tunisian climate. *Soil and Tillage Research*
2687 126:26-33.
- 2688 Karlen DL, Kumar A, Kanwar RS, Cambardella CA & Colvin TS (1998) Tillage system effects on 15-year carbon-
2689 based and simulated N budgets in a tile-drained Iowa field. *Soil Tillage Res* 48:155-165.
- 2690 Karlen DL, Wollenhaupt NC, Erbach DC, Berry EC, Swan JB, Eash NS & Jordahl JL (1994) Long-term tillage
2691 effects on soil quality. *Soil Tillage Res* 32:313-327.
- 2692 Kumar S, Kadono A, Lal R, Dick W (2012) Long-term no-till impacts on organic carbon and properties of two
2693 contrasting soils and corn yields in Ohio. *Soil Science Society of America Journal* 76,5:1798-1809.
- 2694 Kumar S, Nakajima T, Mbonimpa EG, Gautam S, Somireddy UR, Kadono A, Lal R, Chintala R, Rafique R, Fausey
2695 N (2014) Long-term tillage and drainage influences on soil organic carbon dynamics, aggregate stability and corn
2696 yield. *Soil Science and Plant Nutrition* 60,1:108-118.
- 2697 Kushwaha CP, Tripathi SK & Singh KP (2000) Variations in soil microbial biomass and n availability due to
2698 residue and tillage management in a dryland rice agroecosystem. *Soil Tillage Res* 56:153-166.
- 2699 Küstermann B, Munch JC, Hülsbergen K-J (2013) Effects of soil tillage and fertilization on resource efficiency
2700 and greenhouse gas emissions in a long-term field experiment in Southern Germany. *European Journal of*
2701 *Agronomy* 49:61-73.
- 2702 Lal R (1998) Soil quality changes under continuous cropping for seventeen seasons of an alfisol in western Nigeria.
2703 *Land Degrad & Develop* 9:259-274.
- 2704 Lal R, Mahboubi AA & Fausey NR (1994) Long-term tillage and rotation effects on properties of a central Ohio
2705 soil. *Soil Sci Soc Am J* 58:517-522.
- 2706 Lammerding D, Hontoria C, Tenorio J, Walter I (2010). Mediterranean dryland farming: effect of tillage practices
2707 on selected soil properties. *Agronomy Journal* 103,2:382-389.

Ver. 20170605

- 2708 Larney FJ, Bremer E, Janzen HH, Johnston AM & Lindwall CW (1997) Changes in total, mineralizable and light
2709 fraction soil organic matter with cropping and tillage intensities in semiarid southern Alberta, Canada. *Soil Tillage*
2710 *Res* 42:229-240.
- 2711 Laudicina VA, Novara A, Gristina L, Badalucco L (2014) Soil carbon dynamics as affected by long-term
2712 contrasting cropping systems and tillages under semiarid Mediterranean climate. *Applied Soil Ecology* 73:140-
2713 147.
- 2714 Liang A, McLaughlin N, Zhang X, Shen Y, Shi X, Fan R (2011) Short-term effects of tillage practices on soil
2715 aggregation fractions in a Chinese Mollisol. *Acta Agriculturae Scandinavica, Section B - Soil and Plant Science*
2716 61,6:535-542.
- 2717 Liang AZ, Zhang XP, Fang HJ, Yang XM, Drury CF (2007) Short-term effects of tillage practices on organic
2718 carbon in clay loam soil of Northeast China. *Pedosphere* 17,5:619-623.
- 2719 Lilienfein J, Wilcke W, Vilela L, do Carmo Lima S, Thomas R & Zech W (2000) Effect of no-tillage and
2720 conventional tillage systems on the chemical composition of soil solid phase and soil solution of Brazilian savanna.
2721 *J Plant Nutr Soil Sci* 163: 411-419.
- 2722 Liu E, Teclerian S, Yan C, Yu J, Gu R, Liu S, He W, Liu Q (2014) Long-term effects of no-tillage management
2723 practice on soil organic carbon and its fractions in the northern China. *Geoderma* 213:379-384.
- 2724 Lopez-Bellido R, Fontan J, Lopez-Bellido F, Lopez-Bellido L (2009) Carbon sequestration by tillage, rotations,
2725 and nitrogen fertilization in a Mediterranean Vertisol. *Agronomy Journal* 101,1:310-318.
- 2726 Lopez-Fando C, Dorado J, Pardo MT (2007) Effects of zone-tillage in rotation with no-tillage on soil properties
2727 and crop yields in a semi-arid soil from central Spain. *Soil and Tillage Research* 95,1-2:266-276.
- 2728 Lopez-Bellido RJ, Munoz-Romero V, Fuentes-Guerra R, Fernandez-Garcia P, Lopez-Bellido L (2017) No-till: A
2729 key tool for sequestering C and N in microaggregates on a Mediterranean Vertisol. *Soil & Tillage Research*
2730 166:131-137.
- 2731 Lopez-Fando C, Pardo MT (2009) Changes in soil chemical characteristics with different tillage practices in a
2732 semi-arid environment. *Soil and Tillage Research* 104,2:278-284.
- 2733 Lou Y, Xu M, Chen X, He X, Zhao K (2012) Stratification of soil organic C, N and C:N ratio as affected by
2734 conservation tillage in two maize fields of China. *Catena* 95:124-130.
- 2735 Martinez E, Fuentes J, Pino V, Silva P, Acevedo E (2013) Chemical and biological properties as affected by no-
2736 tillage and conventional tillage systems in an irrigated Haploxerol of Central Chile. *Soil and Tillage Research*
2737 126:238-245.
- 2738 Martin-Lammerding D, Tenorio JL, Albarran MM, Zambrana E, Walter E (2013) Influence of tillage practices on
2739 soil biologically active organic matter content over a growing season under semiarid Mediterranean climate.
2740 *Spanish Journal of Agricultural Research* 11,1
- 2741 Martin-Rueda I, Munoz-Guerra LM, Yunta F, Estaban E, Tenorio JL, Lucena JJ (2007) Tillage and crop rotation
2742 effects on barely yield and soil nutrients on a Calcicertidic Haploxeralf. *Soil and Tillage Research* 92,1-2:1-9.
- 2743 McCarty GW, Lyssenko NN & Starr JL (1998) Short-term changes in soil carbon and nitrogen pools during tillage
2744 management transition. *Soil Sci Soc Am J* 62: 1564-1571.
- 2745 McLeod MK, Schwenke GD, Cowie AL, Harden S (2013) Soil carbon is only higher in the surface soil under
2746 minimum tillage in Vertosols and Chromosols of New South Wales North-West Slopes and plains, Australia. *Soil*
2747 *Research* 51:680-694.
- 2748 Melero S, Lopez-Bellido R, Lopez-Bellido L, Munoz-Romero V, Moreno F, Murillo J (2011) Long-term effect of
2749 tillage, rotation and nitrogen fertilizer on soil quality in a Mediterranean Vertisol. *Soil and Tillage Research*
2750 114,2:97-107.
- 2751 Mielke LN, Doran JW & Richards KA (1986) Physical environment near the surface of plowed and no-tilled soils.
2752 *Soil Tillage Res* 7:355-366.
- 2753 Mikha M, Benjamin J, Vigil M, Nielson D (2010) Cropping intensity impacts on soil aggregation and carbon
2754 sequestration in the Central Great Plains. *Soil Science Society of America Journal* 74,5:1712-1719.
- 2755 Mikha M, Vigil M, Benjamin J (2013) Long-term tillage impacts on soil aggregation and carbon dynamics under
2756 wheat-fallow in the Central Great Plains. *Soil Science Society of America Journal* 77,2:594-605.
- 2757 Mrabet R, Saber N, El-brahli A, Lahlou S & Bessam F (2001) Total, particulate organic matter and structural
2758 stability of a Calcixeroll soil under different wheat rotations and tillage systems in a semiarid area of Morocco.
2759 *Soil & Tillage Research* 57: 225-235.

- 2760 Munoz-Romero V, Lopez-Bellido R, Fernandez-Garcia P, Redondo R, Murillo S, Lopez-Bellido L (2017) Effects
2761 of tillage, crop rotation and N application rate on labile and recalcitrant soil carbon in a Mediterranean Vertisol.
2762 *Soil & Tillage Research* 169:118-123.
- 2763 Murage E, Voroney P, Kay B, Deen B, Beyaert R (2006) Dynamics and turnover of soil organic matter as affected
2764 by tillage. *Soil Science Society of America Journal* 71,4:1363-1370.
- 2765 Nyamadzawo G, Chikowo R, Nyamugafata P, Nyamangara J, Giller K (2008) Soil organic carbon dynamics of
2766 improved fallow-maize rotation systems under conventional and no-tillage in Central Zimbabwe. *Nutrient Cycling*
2767 *in Agroecosystems* 81,1:85-93.
- 2768 Nyborg, M., Solberg ED, Malhi SS & Izaurrealde RC (1995) Fertilizer N, crop residue, and tillage alter soil C and
2769 N content in a decade. In: Lal R, Kimble JM, Follett RF, Stewart BA (Eds) *Advances in Soil Science: Management*
2770 *of Carbon Sequestration in Soil* (pp. 93-99). CRC Press, Boca Raton, FL.
- 2771 Olson KR, Lang JM, Ebelhar SA (2005) Soil organic carbon changes after 12 years of no-tillage and tillage of
2772 Grantsburg soils in Southern Illinois. *Soil and Tillage Research* 81,2:217-225.
- 2773 Packer IJ, Hamilton GJ, Koen TB (1992) Runoff, soil loss and soil physical property changes of light textured
2774 surface soils from long-term tillage treatments. *Australian Journal of Soil Research* 30:789-806.
- 2775 Page KL, Dalal RC, Pringle MJ, Bell M, Dang YP, Radford B, Bailey K (2013) Organic carbon stocks in cropping
2776 soils of Queensland, Australia, as affected by tillage management, climate, and soil characteristics. *Soil Research*
2777 51:596-607.
- 2778 Pierce FJ & Fortin MC (1997) Long-term tillage and periodic plowing of a no-tilled soil in Michigan: Impacts,
2779 yield, and soil organic matter. In: Paul EA, Elliot ET, Paustian K, Cole CV (Eds) *Soil organic matter in temperate*
2780 *agroecosystems: Long-term experiments in North America* (pp. 141-149). CRC Press, Boca Raton, FL.
- 2781 Plaza-Bonilla D, Cantero-Martinez C, Alvaro-Fuentes J (2011) Soil carbon dioxide flux and organic carbon
2782 content: effects of tillage and nitrogen fertilization. *Soil Science Society of America Journal* 75,5:1874-1884.
- 2783 Powlson DS & Jenkinson DS (1982) A comparison of the organic matter, biomass, adenosine triphosphate and
2784 mineralizable nitrogen contents of ploughed and direct-drilled soils, *J Agric Sci (Camb)* 97:713-721.
- 2785 Prasad JVNS, Srinivasa Rao CH, Srinivas K, Naga Jyothi CH, Venkateswarlu B, Ramachandrappa BK, Dhanapal
2786 GN, Ravichandra K, Mishra PK (2016) Effect of ten years of reduced tillage and recycling of organic matter on
2787 crop yields, soil organic carbon and its fractions in Alfisols of semi arid tropics of southern India. *Soil & Tillage*
2788 *Research* 156:131-139.
- 2789 Presley D, Sindelar A, Buckley M, Mengel D (2011) Long-term nitrogen and tillage effects on soil physical
2790 properties under continuous grain sorghum. *Agronomy Journal* 104,3:749-755.
- 2791 Puget P, Lal R (2005) Soil organic carbon and nitrogen in a Mollisol in Central Ohio as affected y tillage and land
2792 use. *Soil and Tillage Research* 81,1-2:201-213.
- 2793 Quincke J.A., Wortmann C.S., Mamo M., Franti T., Drijber RA (2006) Occasional tillage of no-till systems.
2794 *Agronomy Journal* 99,4:1158-1168.
- 2795 Rasmussen PE & Albrecht SL (1997) Crop management effects on organic arbon in semi-arid Pacific northwest
2796 soils. In: Lal R, Kimble JM, Follett RF, Stewart BA (Eds) *Advances in Soil Science: Management of Carbon*
2797 *Sequestration in Soil* (pp. 209-219). CRC Press, Boca Raton, FL.
- 2798 Rhoton FE, Bruce RR, Buehring NW, Elkins GB, Langdale CW & Tyler DD (1993) Chemical and physical
2799 characteristics of four soil types under conventional and no-tillage systems. *Soil Tillage Res* 28:51-61.
- 2800 Robertson F, Armstrong R, Partington D, Perris R, Oliver I, Aumann C, Crawford D, Rees D (2015) Effect of
2801 cropping practices on soil organic carbon: evidence from long-term field experiments in Victoria, Australia. *Soil*
2802 *Research* 53:636-646.
- 2803 Ross CW & Hughes KA (1985) Maize/oats forage rotation under 3 cultivation systems, 1978-83 2. Soil properties.
2804 *New Zealand J Agric Res* 28:209-219.
- 2805 Sa JCM, Cerri CC, Dick WA, Lal R, Filho SPV, Piccolo MC & Feigl BE (2001) Organic matter dynamics and
2806 carbon sequestration rates for a tillage chronosequence in a Brazilian Oxisol. *Soil Sci Soc Am J* 65:1486-1499.
- 2807 Sa JCM, Tivet F, Lal R, Briedis C, Hartman DC, dos Santos JZ, dos Santos JB (2014) Long-term tillage systems
2808 impacts on soil C dynamics, soil resilience and agronomic productivity of a Brazilian Oxisol. *Soil and Tillage*
2809 *Research* 136:38-50.
- 2810 Saffigna, P.G., D.S. Powlson, P.C. Brookes, and G.A. Thomas. 1989. Influence of sorghum residues and tillage
2811 on soil organic matter and soil microbial biomass in an Australian vertisol. *Soil Biol and Biochem* 21: 759-765.

Ver. 20170605

- 2812 Sainju U, Caesar T, Lenssen A, Evans R, Kolberg R (2009) Tillage and cropping sequence impacts on nitrogen
2813 cycling in dryland farming in Eastern Montana, USA. *Soil and Tillage Research* 103,2:332-341.
- 2814 Sainju U, Lenssen A, Caesar-Thonthat T, Waddell J (2005) Carbon sequestration in dryland soils and plant residue
2815 as influenced by tillage and crop rotation. *Journal of Environmental Quality* 35,4:1341-1347.
- 2816 Sainju U, Lenssen A, Caesar-TonThat R, Jabro J, Lartey R, Evans R, Allen B (2011) Dryland residue and soil
2817 organic matter as influenced by tillage, crop rotation, and cultural practice. *Plant and Soi.* 338,1-2:27-41.
- 2818 Sainju U, Singh B, Whitehead W, Wang S (2005) Carbon supply and storage in tilled and nontilled soils as
2819 influenced by cover crop and nitrogen fertilization. *Journal of Environmental Quality* 35,4:1507-1517.
- 2820 Sainju UM, Senwo ZN, Nyakatawa EZ, Tazisong IA, Reddy KC (2008) Soil carbon and nitrogen sequestration as
2821 affected by long-term tillage, cropping systems, and nitrogen fertilizer sources. *Agriculture, Ecosystems and*
2822 *Environment* 127,3-4:234-240.
- 2823 Sainju UM, Singh BP & Whitehead WF (2002) Long-term effects of tillage, cover crops, and nitrogen fertilization
2824 on organic carbon and nitrogen concentrations in sandy loam soils in Georgia, USA. *Soil Tillage Res* 63:167-179.
- 2825 Salinas-Garcia JR, Hons FM & Matocha JE (1997) Long-term effects of tillage and fertilization on soil organic
2826 matter dynamics. *Soil Sci Soc Am J* 61:152-159.
- 2827 Salinas-Garcia JR, Velazquez-Garcia JdJ, Gallardo-Valdez M, Diaz-Mederos P, Caballero-Hernandez F, Tapia-
2828 Vargas LM, Rosales-Robles E (2002) Tillage effects on microbial biomass and nutrient distribution in soils under
2829 rain-fed corn production in central-western Mexico. *Soil and Tillage Research* 66,2:143-152.
- 2830 Salvo L, Hernandez J, Ernst O (2010) Distribution of soil organic carbon in different size fractions, under pasture
2831 and crop rotations with conventional tillage and no-till systems. *Soil & Tillage Research* 109:116-122.
- 2832 Schomberg H, Jones O (1998) Carbon and nitrogen conservation in dryland tillage and cropping systems. *Soil*
2833 *Science Society of America Journal* 63,5: 1359-1366.
- 2834 Sheehy J, Six J, Alakukku L, Regina K (2013) Fluxes of nitrous oxide in tilled and no-tilled boreal arable soils.
2835 *Agriculture, Ecosystems and Environment* 164:190-199.
- 2836 Shi X, Yang X, Drury C, Reynolds W, McLaughlin N, Welacky T, Zhang X (2011) Zone tillage impacts on organic
2837 carbon of a clay loam in Southwestern Ontario. *Soil Science Society of America Journal* 75,3:1083-1089.
- 2838 Shrestha BM, Singh BR, Forte C, Certini G (2015) Long-term effects of tillage, nutrient application and crop
2839 rotation on soil organic matter quality assessed by NMR spectroscopy. *Soil Use and Management* 31:358-366.
- 2840 Shukla MK, Lal R, Ebinger M (2006) Determining soil quality indicators by factor analysis. *Soil and Tillage*
2841 *Research* 87,2: 194-204.
- 2842 Singh P, Heikkinen J, Ketoja E, Nuutinen V, Palojärvi A, Sheehy J, Esala M, Mitra S, Alakukku L, Regina K
2843 (2015) Tillage and crop residue management methods had minor effects on the stock and stabilization of topsoil
2844 carbon in a 30-year field experiment. *Science of the Total Environment* 518-519:337-344.
- 2845 Six J, Paustian K, Elliott ET & Combrink C (2000) Soil structure and organic matter: I. Distribution of aggregate-
2846 size classes and aggregate-associated carbon. *Soil Sci Soc Am J* 64:681-689.
- 2847 Sombrero A, de Benito A (2010) Carbon accumulation in soil. Ten-year study of conservation tillage and crop
2848 rotation in a semi-arid area of Castile-Leon, Spain. *Soil and Tillage Research* 107,2:64-70.
- 2849 Stenberg M, Stenberg B & Rydberg T (2000) Effects of reduced tillage and liming on microbial activity and soil
2850 properties in a weakly-structured soil. *Applied Soil Ecology* 14:135-145.
- 2851 Sun B, Hallett P, Caul S, Daniell T, Hopkins D (2011) Distribution of soil carbon and microbial biomass in arable
2852 soils under different tillage regimes. *Plant and Soil* 338,1-2:17-25.
- 2853 Taboada MA, Micucci FG, Cosentino DJ & Lavado RS (1998) Comparison of compaction induced by
2854 conventional and zero tillage in two soils of the Rolling Pampa of Argentina. *Soil Tillage Res* 49:57-63.
- 2855 Thomas GA, Dalal RC, Standley J (2007) No-till effects on organic matter, pH, cation exchange capacity and
2856 nutrient distribution in a Luvisol in the semi-arid subtropics. *Soil & Tillage Research* 94:295-304.
- 2857 Tian S, Wang Y, Ning T, Li N, Zhao H, Wang B, Li Z, Chi S (2013) Continued no-till and subsoiling improved
2858 soil organic carbon and soil aggregation levels. *Agronomy Journal* 106,1:212-218.
- 2859 Tivet F, Sa JDM, Lal R, Borszowski P, Briedis C, dos Santos J, Sa M, Hartman DDC, Eurich G, Farias A,
2860 Bousinac S, Seguy L (2013) Soil organic carbon fraction losses upon continuous plow-based tillage and its
2861 restoration by diverse biomass-C inputs under no-till in sub-tropical and tropical regions of Brazil. *Geoderma* 209-
2862 210:214-225.

- 2863 Ussiri D, Lal R (2009) Long-term tillage effects on soil carbon storage and carbon dioxide emissions in continuous
2864 corn cropping system from an alfisol in Ohio. *Soil and Tillage Research* 104,1:39-47.
- 2865 VandenBygaart AJ, Yang XM, Kay BD, Aspinall JD (2002) Variability in carbon sequestration potential in no-till
2866 soil landscapes of southern Ontario. *Soil & Tillage Research* 65:231-241.
- 2867 van Groenigen K, Hastings A, Forristal D, Roth B, Jones M, Smith P (2011) Soil C storage as affected by tillage
2868 and straw management: An assessment using field measurements and model predictions. *Agriculture, Ecosystems,*
2869 *and Environment* 140,1-2:218-225.
- 2870 Varvel GE, Wilhelm WW (2011) No-tillage increases soil profile carbon and nitrogen under long-term rainfed
2871 cropping systems. *Soil and Tillage Research* 114,1:28-36.
- 2872 Venterea RT, Baker JM, Dolan MS, Spokas KA (2006) Carbon and nitrogen storage are greater under biennial
2873 tillage in a Minnesota corn-soybean rotation. *Soil Science Society of America Journal* 70,5:1752-1762.
- 2874 Viaud V, Angers DA, Parnaudeau V, Morvan T, Menasseri Aubry S (2010) Response of organic matter to reduced
2875 tillage and animal manure in a temperate loamy soil. *Soil Use and Management* 27,1:84-93.
- 2876 Wander MM, Bidart MG & Aref S (1998) Tillage impacts on depth distribution of total and particulate organic
2877 matter in three Illinois soils. *Soil Sci Soc Am J*, 62:1704-1711.
- 2878 Wang WJ, Dalal RC (2006) Carbon inventory for a cereal cropping system under contrasting tillage, nitrogen
2879 fertilization and stubble management practices. *Soil and Tillage Research* 91,1-2:68-74.
- 2880 Wanniarachchi SD, Voroney RP, Vyn TJ, Beyaert RP & MacKenzie AF (1999) Tillage effects on the dynamics
2881 of total and corn-residue-derived soil organic matter in two southern Ontario soils. *Can J Soil* 79: 473-480.
- 2882 Wright A, Hons F (2004) Soil carbon and nitrogen storage in aggregates from different tillage and crop regimes.
2883 *Soil Science Society of America Journal* 69,1:141-147.
- 2884 Xu, S. Q., Zhang, M. Y., Zhang, H. L., Chen, F., Yang, G. L. and Xiao, X. P. 2013. Soil organic carbon stocks as
2885 affected by tillage systems in a double-cropped rice field. *Pedosphere* 23 (5): 696–704
- 2886 Yang XM & Kay BD (2001) Impacts of tillage practices on total, loose- and occluded-particulate, and humified
2887 organic carbon fractions in soils within a field in southern Ontario. *Can J Soil* 81: 149-156.
- 2888 Yang XM & Wander MM (1999) Tillage effects on soil organic carbon distribution and storage in a silt loam soil
2889 in Illinois. *Soil Tillage Res* 52:1-9.
- 2890 Zhang M, Sparrow S, Lewis C, Knight C (2007) Soil properties and barley yield under a twenty-years experiment
2891 of tillage, straw management and nitrogen application rates in the sub-arctic area of Alaska. *Acta Agriculturae*
2892 *Scandinavica Section B-Soil and Plant Science* 57:374-382.
- 2893 Zhang X, Xin X, Zhu A, Zhang J, Yang W (2017) Effects of tillage and residue managements on organic C
2894 accumulation and soil aggregation in a sandy loam soil of the North China plain. *Catena* 156: 176-183.
- 2895
- 2896 **Sources of Data for Input Factor (Literature review is not completed and additional reference will be added**
2897 **in the second order draft)**
- 2898 Agbenin JO & JT Goladi (1997) Carbon, nitrogen and phosphorus dynamics under continuous cultivation as
2899 influenced by farmyard manure and inorganic fertilizers in the savanna of northern Nigeria. *Agric Ecosyst &*
2900 *Environ* 63:17-24.
- 2901 Bayer C, Mielniczuk J, Amado TJC, Martin-Neto L & Fernandes SV (2000) Organic matter storage in a sandy
2902 clay loam Acrisol affected by tillage and cropping systems in southern Brazil. *Soil Tillage Res* 54:101-109.
- 2903 Black AL & Tanaka DL (1997) A conservation tillage-cropping systems study in the northern Great Plains of the
2904 United States. In: Paul EA, Elliot ET, Paustian K, Cole CV (Eds) *Soil organic matter in temperate agroecosystems:*
2905 *Long-term experiments in North America* (pp. 335-342). CRC Press, Boca Raton, FL.
- 2906 Bordovsky DG, Choudhary M, & Gerard CJ (1999) Effect of tillage, cropping, and residue management on soil
2907 properties in the Texas rolling plains. *Soil Sci* 164:331-340.
- 2908 Bremer E, Janzen HH & Johnston AM (1994) Sensitivity of total, light fraction and mineralizable organic matter
2909 to management practices in a Lethbridge soil. *Soil Sci Soc Am J* 74:131-138.
- 2910 Buyanovsky GA & Wagner GH (1998) Carbon cycling in cultivated land and its global significance. *Glob Change*
2911 *Biol* 4:131-141.

Ver. 20170605

- 2912 Campbell CA & Zentner RP (1993) Soil organic matter as influenced by crop rotations and fertilization. *Soil Sci*
2913 *Soc Am J* 57:1034-1040.
- 2914 Campbell CA, Biederbeck VO, McConkey BG, Curtin D & Zentner RP (1999) Soil quality-Effect of tillage and
2915 fallow frequency. Soil organic matter quality as influenced by tillage and fallow frequency in a silt loam in
2916 southwestern Saskatchewan. *Soil Biol and Biochem* 31:1-7.
- 2917 Campbell CA, Bowren KE, Schnitzer M, Zentner RP & Townley-Smith L (1991) Effect of crop rotations and
2918 fertilization on soil organic matter and some biochemical properties of a thick black Chernozem. *Can J Soil* 71:377-
2919 387.
- 2920 Campbell CA, McConkey BG, Zentner RP, Selles F & Curtin D (1996) Long-term effects of tillage and crop
2921 rotations on soil organic C and total N in a clay soil in southwestern Saskatchewan. *Can J Soil* 76:395-401.
- 2922 Campbell CA, Zentner RP, Selles F, Biederbeck VO, McConkey BG, Blomert B & Jefferson PG (2000)
2923 Quantifying short-term effects of crop rotations on soil organic carbon in southwestern Saskatchewan. *Can J Soil*
2924 80:193-202.
- 2925 Campbell CA, Lafond GP, Moulin AP, Townley-Smith L & Zentner RP (1997) Crop production and soil organic
2926 matter in long-term crop rotations in the sub-humid northern Great Plains of Canada. In: Paul EA, Elliot ET,
2927 Paustian K, Cole CV (Eds) *Soil organic matter in temperate agroecosystems: Long-term experiments in North*
2928 *America* (pp. 297-315). CRC Press, Boca Raton, FL.
- 2929 Dalal RC, Henderson PA & Glasby JM (1991) Organic matter and microbial biomass in a vertisol after 20 yr of
2930 zero tillage. *Soil Biol and Biochem* 23:435-441.
- 2931 Dick WA & Durkalski JT (1997) No-tillage production agriculture and carbon sequestration in a typic fragiudalf
2932 soil of northeastern Ohio. In: Lal R, Kimble JM, Follett RF, Stewart BA (Eds) *Advances in Soil Science:*
2933 *Management of Carbon Sequestration in Soil* (pp. 59-71). CRC Press, Boca Raton, FL.
- 2934 Dick WA, Edwards WM & McCoy EL (1997) Continuous application of no-tillage to Ohio soils: Changes in crop
2935 yields and organic matter-related soil properties. In: Paul EA, Elliot ET, Paustian K, Cole CV (Eds) *Soil organic*
2936 *matter in temperate agroecosystems: Long-term experiments in North America* (pp. 171-182). CRC Press, Boca
2937 Raton, FL.
- 2938 Freixo AA, Machado P, dos Santos HP, Silva CA & Fadigas F (2002) Soil organic carbon and fractions of a Rhodic
2939 Ferralsol under the influence of tillage and crop rotation systems in southern Brazil. *Soil Tillage Res* 64:221-230.
- 2940 Ghuman BS & Sur HS (2001) Tillage and residue management effects on soil properties and yields of rainfed
2941 maize and wheat in a subhumid subtropical climate. *Soil Tillage Res* 58:1-10.
- 2942 Graham MH, Haynes RJ & Meyer JH (2002) Soil organic matter content and quality: effects of fertilizer
2943 applications, burning and trash retention on a long-term sugarcane experiment in South Africa. *Soil Biol and*
2944 *Biochem* 34:93-102.
- 2945 Grandy AS, Porter GA & Erich MS (2002) Organic amendment and rotation crop effects on the recovery of soil
2946 organic matter and aggregation in potato cropping systems. *Soil Sci Soc Am J* 66:1311-1319.
- 2947 Halvorson AD, Wienhold BJ & Black AL (2002) Tillage, nitrogen, and cropping system effects on soil carbon
2948 sequestration. *Soil Sci Soc Am J* 66:906-912.
- 2949 Hernanz JL, Lopez R, Navarrete L & Sanchez-Giron V (2002) Long-term effects of tillage systems and rotations
2950 on soil structural stability and organic carbon stratification in semiarid central Spain. *Soil Tillage Res* 66:129-141.
- 2951 Janzen HH (1987) Soil organic matter characteristics after long-term cropping to various spring wheat rotations.
2952 *Can J Soil* 67:845-856.
- 2953 Lal R (1998) Soil quality changes under continuous cropping for seventeen seasons of an alfisol in western Nigeria.
2954 *Land Degrad & Develop* 9:259-274.
- 2955 Larney FJ, Bremer E, Janzen HH, Johnston AM & Lindwall CW (1997) Changes in total, mineralizable and light
2956 fraction soil organic matter with cropping and tillage intensities in semiarid southern Alberta, Canada. *Soil Tillage*
2957 *Res* 42:229-240.
- 2958 Potter KN, Jones OR, Torbert HA & Unger PW (1997) Crop rotation and tillage effects on organic carbon
2959 sequestration in the semiarid southern Great Plains. *Soil Sci* 162: 140-147.
- 2960 Rasmussen PE & Albrecht SL (1997) Crop management effects on organic carbon in semi-arid Pacific northwest
2961 soils. In: Lal R, Kimble JM, Follett RF, Stewart BA (Eds) *Advances in Soil Science: Management of Carbon*
2962 *Sequestration in Soil* (pp. 209-219). CRC Press, Boca Raton, FL.

- 2963 Sainju UM, Singh BP & Whitehead WF (2002) Long-term effects of tillage, cover crops, and nitrogen fertilization
2964 on organic carbon and nitrogen concentrations in sandy loam soils in Georgia, USA. *Soil Tillage Res* 63:167-179.
- 2965 Sherrod, L.A., G.A. Peterson, D.G. Westfall & L.R. Ahuja (2003). Cropping intensity enhances soil organic carbon
2966 and nitrogen in a no-till agroecosystem. *Soil Sci Soc Am J.* 67:1553-1543
- 2967 Sidhu AS & Sur HS (1993) Effect of incorporation of legume straw on soil properties and crop yield in a maize-
2968 wheat sequence. *Trop Agric (Trinidad)* 70:226-229.
- 2969 Westerhof R, Vilela L, Azarza M & Zech W (1998) Land use effects on labile N extracted with permanganate and
2970 the nitrogen management index in the Cerrado region of Brazil. *Biol Fertil Soils* 27:353-357.
- 2971

Annex 5A.2 Estimation of Default Emission Factors and Scaling Factors for CH₄ Emission from Rice Cultivation

The default emission factors and scaling factors were estimated using a statistical model (Yan et al., 2003) after updating the databases used in developing the IPCC 2006 Guidelines.

We conducted a literature review and collected the latest measurement data of CH₄ emission from rice fields that are available in peer-reviewed journals. For each measurement, documented information included average flux in the rice-growing seasons; integrated seasonal emission; water regimes in the rice-growing season; water conditions in the season before rice planting (preseason water status); timing, type, and amount of organic amendment; nitrogen fertilization; soil properties; location; climate; year; and season of measurement.

The effects of controlling variables on CH₄ flux were estimated by using a linear mixed effect model, which is suitable for analyzing unbalanced data, that is, data having unequal numbers of observations in the subclasses. CH₄ flux data do not meet model assumptions of a normal distribution of the error, and so they were transformed into a log-normal distribution. Thus, flux data were first log-transformed and then analyzed by the following linear model:

$$\ln(\text{flux}) = \text{constant} + a \times \ln(\text{SOC}) + pH_m + PW_i + WT_j + CL_k + OM_l \times \ln(1 + AOM_l),$$

where flux is the average CH₄ flux during the rice-growing season; SOC is soil organic carbon content; *a* is the effects of SOC; *PW_i* is the effect of preseason water status (*i* is flooded, long drainage, short drainage, double drainage, or unknown); *WT_j* is the effect of water regime in the rice-growing season (*j* is continuous flooding, single drainage, multiple drainage, wet season rainfed, dry season rainfed, deepwater, or unknown); *CL_k* is the effect of climate; *OM_l* is the effect of added organic materials (*l* is compost, farmyard manure, green manure, rice straw used on-season, or rice straw used off-season); *AOM_l* is the amount of organic amendment in t ha⁻¹; *pH_m* is the effect of soil pH (*m* is one of the pH classes). Since optimum soil pH has often been reported for CH₄ emission, soil pH was classified as a category variable and grouped into <4.5, 4.5–5.0, 5.0–5.5, 5.5–6.0, 6.0–6.5, 6.5–7.0, 7.0–7.5, 7.5–8.0 and ≥8.0.

The effects of the controlling variables on CH₄ flux were computed by fitting the model to observations using the SAS/STAT procedure MIXED (Release 8.01, SAS Institute Inc., Cary, NC, USA). Global and regional default emission factors and scaling factors (water regimes during the cultivation period, water regimes before the cultivation period, organic amendment) were estimated using the resulting model.

References (Additional References will be added in the Second Order Draft after completing the analysis)

Yan, X. Y., K. Yagi, H. Akiyama, and H. Akimoto (2005), Statistical analysis of the major variables controlling methane emission from rice fields, *Global Change Biol.*, 11(7), 1131 – 1141, doi:10.1111/j.1365-2486.2005.00976.x.

3006 **Annex 5A.3 Estimation of Coefficients and Factors for Biomass**
3007 **C Emissions and Removals in Cropland**

3008

3009 To be completed for the second order draft.

3010