

# CHAPTER 2

---

## DRAINED INLAND ORGANIC SOILS

### **Coordinating Lead Authors**

Matthias Drösler (Germany), Louis V. Verchot (USA), Annette Freibauer (Germany) and Genxing Pan (China)

### **Lead Authors**

Christopher David Evans (UK), Richard A. Bourbonniere (Canada), Jukka P. Alm (Finland), Susan Page (UK), Fahmuddin Agus (Indonesia), Kristell Hergoualc'h (France), John Couwenberg (EC/WI/Germany/Netherlands), Jyrki Jauhiainen (Finland), Supiandi Sabiham (Indonesia) and Changke Wang (China)

### **Contributing Authors**

Nalin Srivastava (TFI TSU), Laura Borgeau-Chavez (USA), Aljosja Hooijer (The Netherlands), Kari Minkkinen (Finland), Nancy French (USA), Tara Strand (USA/New Zealand), Andrey Sirin (Russian Federation), Robert Mickler (USA), Kevin Tansey (UK) and Narasimhan Larkin (USA)

### **Review Editors**

David Pare (Canada), Bernard Siska (Slovakia) and Iman Rusmana (Indonesia)

18

## Contents

19	2.1 Introduction.....	4
20	2.2 Land Remaining in a Land-use Category.....	5
21	2.2.1 CO <sub>2</sub> emissions and removals from drained inland organic soils .....	5
22	2.2.2 Non-CO <sub>2</sub> emissions and removals from drained inland organic soils .....	17
23	2.3 Land Converted to a New Land-use Category .....	39
24	2.2.3 CO <sub>2</sub> emissions and removals from drained inland organic soils .....	39
25	2.2.4 Non-CO <sub>2</sub> emissions and removals from drained inland organic soils .....	41
26	Annex 2A.1 Scientific background for developing CO <sub>2</sub> -C emission/removal factors for organic soil from the	
27	scientific literature in Table 2.1.....	43
28	Annex 2A.2 Derivation of ditch CH <sub>4</sub> emission factors.....	45
29	Annex 2A.3 Derivation of DOC emission factors.....	47
30	Annex 2A.4 Derivation of CO <sub>2</sub> -C and non-CO <sub>2</sub> emission factors for emissions from burning of drained	
31	organic soils from scientific literature in Tables 2.6 and 2.7.....	52
32	Appendix 2a.1 Estimation for Particulate Organic Carbon (POC) and Dissolved Inorganic Carbon	
33	(DIC) loss from peatlands and drained organic soils: Basis for future methodological development	54

34

35

## Equations

36

37	Equation 2.1 Annual carbon stock changes for a stratum of a land-use category as a sum of changes in all pools	
38	(Equation 2.3 in the Chapter 2, Volume 4, 2006 IPCC Guidelines).....	5
39	Equation 2.1 CO <sub>2</sub> -C emissions/removals by drained inland organic soils .....	6
40	Equation 2.3 Annual on-site CO <sub>2</sub> -C emissions/removals from drained inland organic soils excluding emissions	
41	from fires.....	7
42	Equation 2.4 Annual off-site CO <sub>2</sub> emissions due to DOC loss from drained organic soils (CO <sub>2</sub> ).....	15
43	Equation 2.5 Emission factor for annual CO <sub>2</sub> emissions due to DOC export from drained organic soils.....	15
44	Equation 2.6 Annual CH <sub>4</sub> emission from drained organic soils.....	18
45	Equation 2.7 Direct N <sub>2</sub> O emissions from managed organic soils .....	26
46	Equation 2.8 Annual CO <sub>2</sub> -C and non-CO <sub>2</sub> emissions from organic soil fire.....	32
47	Equation 2a.1 Calculation of POC export from drained organic soils .....	54

48

## Figures

49	Figure 2.1 Summary of fluxes from drained inland organic soils.....	5
50	Figure 2.2 Generic decision tree for identification of the appropriate tier to estimate greenhouse gas	
51	emissions from fires on organic soils .....	33

52

53

## Tables

54	Table 2.1 Tier 1 CO <sub>2</sub> emission/removal factors for drained organic soils in all land-use categories** .....	9
55	Table 2.2 Default DOC emission factors for drained organic soils.....	16
56	Table 2.3 Tier 1 CH <sub>4</sub> emission/removal factors for drained organic soils (EF <sub>CH<sub>4</sub>_land</sub> ) in all land-use categories ..	21
57	Table 2.4 Default CH <sub>4</sub> emission factors for drainage ditches .....	25
58	Table 2.5 Tier 1 Direct N <sub>2</sub> O emission/removal factors for drained organic soils in all land-use categories.....	28
59	Table 2.6 Organic soil fuel consumption values (mass of dry matter for a range of organic soil and fire types, to	
60	be used in conjunction with equation 2.8, to estimate the product of quantities M <sub>B</sub> and C <sub>f</sub> ) .....	35
61	Table 2.7 Emission factors (g kg <sup>-1</sup> dry matter burned) for organic soil fires. Values are means ± 95% CI (To be	
62	used as quantity G <sub>ef</sub> in Equation 2.8) .....	36
63	Table 2A.3 DOC concentration (above) or flux (below) comparisons between drained and undrained organic	
64	soils, used to derive default value for ΔDOC <sub>DRAINAGE</sub> .....	50

65

## Boxes

67	Box 2.1 Recent advances in satellite-derived fire products.....	38
----	---	----

68

## 69 2.1 INTRODUCTION

70 Organic soils are defined in Chapter 3 Annex 3A.5 of the *2006 IPCC Guidelines* and Section 5, Chapter 1,  
71 section 5 of this *Wetlands Supplement*. The guidance in this Chapter applies to all organic soils that have been  
72 drained, i.e., drainage of lands that started in the past and that still persists, or newly drained lands within the  
73 reporting period. This means that the water table level is at least temporarily below natural levels. Within each  
74 land-use category water table level is manipulated to varying degrees depending on land-use purpose, e.g., for  
75 cultivating cereals, rice, or for aquaculture, which can be reflected by different drainage classes.

76 This Chapter deals with inland organic soils, which do not meet the definition of “coastal” defined in Chapter 4  
77 of this *Wetlands Supplement*.

78 This Chapter provides supplementary guidance on estimating greenhouse gas emissions and removals from  
79 drained inland organic soils in the following land-use categories as defined in the *2006 IPCC Guidelines* Volume  
80 4: Chapter 4 (Forest Land), Chapter 5 (Cropland), Chapter 6 (Grassland), Chapter 7 (Wetlands), Chapter 8  
81 (Settlements) and Chapter 9 (Other Land). Managed coastal organic soils are covered in Chapter 4 of this  
82 Supplement. Rewetted organic soils are considered in Chapter 3 of the *Wetlands Supplement*.

83 This Chapter clarifies Volume 4 of the *2006 IPCC Guidelines* by summarizing all emission factors and  
84 harmonizing the methods for organic soils in all land-use types. On the basis of recent advances in scientific  
85 information, this Chapter also updates, improves, and completes methodologies and emission factors for  
86 greenhouse gas emissions and removals of the *2006 IPCC Guidelines* and fills gaps where new scientific  
87 knowledge allows implementation of robust methodologies and use of better emission factors at the Tier 1 level.

88 This Chapter updates the *2006 IPCC Guidelines* for:

- 89 • CO<sub>2</sub> emissions and removals from drained organic soils (referring to Chapters 4 to 9, Volume 4, *2006 IPCC*  
90 *Guidelines*);
- 91 • CH<sub>4</sub> emissions from drained organic soils (referring to Chapter 7, Volume 4, *2006 IPCC Guidelines*);
- 92 • N<sub>2</sub>O emissions from drained organic soils (referring to Chapter 11, Volume 4, *2006 IPCC Guidelines*).

93 This Chapter gives new guidance not contained in the *2006 IPCC Guidelines* by:


- 94 • providing methodologies and emission factors for CH<sub>4</sub> emissions from drainage ditches (referring to  
95 Chapters 4 to 9, Volume 4, *2006 IPCC Guidelines*);
- 96 • providing methodologies and emission factors for off-site CO<sub>2</sub> emissions associated with dissolved organic  
97 carbon (DOC) release from organic soils to drainage waters (referring to Chapters 4 to 9, Volume 4, *2006*  
98 *IPCC Guidelines*);
- 99 • providing methodologies and emission factors for CO<sub>2</sub>, CH<sub>4</sub> and CO emissions from peat fires

100 The chapter also contains an appendix that provides the basis for future methodological development for  
101 estimating CO<sub>2</sub> emissions associated with other forms of waterborne carbon loss, specifically particulate organic  
102 carbon (POC) and dissolved inorganic carbon (referring to Chapters 4 to 9), Volume 4, *2006 IPCC Guidelines*.  
103 All fluxes are summarized in Figure 2.1.

104

105 **Figure 2.1** Summary of fluxes from drained inland organic soils

106



107

## 108 2.2 LAND REMAINING IN A LAND-USE

## 109 CATEGORY

110 The 2006 IPCC Guidelines provide guidance for carbon stock changes in the carbon pools in above-ground and  
 111 below-ground biomass, dead wood and litter as well as soil for managed land on organic soils. This Chapter  
 112 updates the 2006 IPCC Guidelines for the soil organic carbon pool in organic soils.

### 113 2.2.1 CO<sub>2</sub> emissions and removals from drained inland

### 114 organic soils

115 This section deals with the impacts of drainage and management on CO<sub>2</sub> emissions and removals from organic  
 116 soils due to organic matter decomposition and loss of dissolved organic carbon (DOC) in drainage waters. DOC  
 117 losses lead to off-site CO<sub>2</sub> emissions. There are also erosion losses of particulate organic carbon (POC) and  
 118 waterborne transport of dissolved inorganic carbon (primarily dissolved CO<sub>2</sub>) derived from autotrophic and  
 119 heterotrophic respiration within the organic soil. At present the science and available data are not sufficient to  
 120 provide guidance on CO<sub>2</sub> emissions or removals associated with these waterborne carbon fluxes; Appendix 2a.1  
 121 provides a basis for future methodological development in this area. General information and guidance for  
 122 estimating changes in soil carbon stocks are provided in in Section 2.3.3, Chapter 2, Volume 4 in the 2006 IPCC  
 123 Guidelines which needs to be read before proceeding with the guidance provided here. This guidance is based on  
 124 the observation that in drained inland organic soils, emissions persist as long as the soil remains drained or as  
 125 long as organic matter remains (Wösten *et al.*, 1997; Deverel and Leighton, 2010).

126 Equation 2.3 in Chapter 2, Volume 4, 2006 IPCC Guidelines refers to annual carbon stock changes for a stratum  
 127 of a land-use category as a sum of changes in all pools. This section addresses the stratum of a land-use category  
 128 on drained inland organic soils. The Equation is repeated here as Equation 2.1 to demonstrate how the guidance  
 129 in this *Wetlands Supplement* links to the 2006 IPCC Guidelines.

130

131 **EQUATION 2.1**  
 132 **ANNUAL CARBON STOCK CHANGES FOR A STRATUM OF A LAND-USE CATEGORY AS A SUM OF**  
 133 **CHANGES IN ALL POOLS**  
 134 **(EQUATION 2.3 IN THE CHAPTER 2, VOLUME 4, 2006 IPCC GUIDELINES)**

$$135 \Delta C_{LU_i} = \Delta C_{AB} + \Delta C_{BB} + \Delta C_{DW} + \Delta C_{LI} + \Delta C_{SO} + \Delta C_{HWP}$$

136

137 Where:

138  $\Delta C_{LU_i}$  = carbon stock changes for a stratum of a land-use category

Final Draft

139 Subscripts denote the following carbon pools:

140 AB = above-ground biomass

141 BB = below-ground biomass

142 DW = dead wood

143 LI = litter

144 SO = soils

145 HWP = harvested wood products

146

147 The guidance for the carbon pools above-ground biomass, below-ground biomass, deadwood, litter and  
148 harvested wood products in the *2006 IPCC Guidelines* is not further dealt with in these guidelines.

149 This section of the *Wetlands Supplement* updates and complements the guidance on drained inland organic soils  
150 component of  $\Delta C_{SO}$ , which was called  $L_{organic}$  in Equation 2.24, Chapter 2, Volume 4 of the *2006 IPCC*  
151 *Guidelines*. For transparent distinction between drained and rewetted organic soils, the term is further specified  
152 as  $CO_2-C_{organic, drained}$  in Equation 2.2.  $CO_2-C_{organic, drained}$  consists of on-site  $CO_2$  emissions/removals of the  
153 organic soil from mineralization and sequestration processes ( $CO_2-C_{soil-on-site}$ ), off-site  $CO_2$  emissions from  
154 leached carbon from the organic soil ( $CO_2-C_{DOC}$ ) and anthropogenic peat fires ( $L_{fire}$ ). Countries are encouraged to  
155 consider particulate organic carbon (POC) when using higher tier methodologies (see Appendix 2a.1).  $CO_2$   
156 emissions from peat fires have not been explicitly addressed in Equation 2.3, Chapter 2, Volume 4 of the *2006*  
157 *IPCC Guidelines*, but can be important on drained inland organic soils. Therefore,  $CO_2$  emissions from peat fires  
158 are included in Equation 2.2 as  $L_{fire}$  (Section 2.2.2.3).

159

160

161

162

**EQUATION 2.1**

**CO<sub>2</sub>-C EMISSIONS/REMOVALS BY DRAINED INLAND ORGANIC SOILS**

$$CO_2 - C_{organic, drained} = CO_2 - C_{on-site} + CO_2 - C_{DOC} + L_{fire}$$

163 Where:

164  $CO_2-C_{organic, drained}$  =  $CO_2$ -C emissions/removals by drained organic soils, tonnes C yr<sup>-1</sup>

165  $CO_2-C_{on-site}$  =  $CO_2$ -C emissions/removals by drained organic soils, tonnes C yr<sup>-1</sup>

166  $CO_2-C_{DOC}$  =  $CO_2$ -C emissions from dissolved organic carbon exported from drained organic soils, tonnes  
167 C yr<sup>-1</sup>

168  $L_{fire}$  =  $CO_2$ -C emissions from burning of drained organic soils, tonnes C yr<sup>-1</sup>

### 169 **2.2.1.1 ON-SITE CO<sub>2</sub> EMISSIONS/REMOVALS FROM DRAINED** 170 **INLAND ORGANIC SOILS (CO<sub>2</sub>-C<sub>ON-SITE</sub>)**

171 This section gives supplementary guidance for  $CO_2$  emissions and removals from drained inland organic soils in  
172 all land-use categories as defined in Section 2.3.3, Chapter 2, Volume 4, of the *2006 IPCC Guidelines*. The IPCC  
173 land-use categories are discussed in Chapter 4 (Forest Land), Chapter 5 (Cropland), Chapter 6 (Grassland),  
174 Chapter 7 (Wetlands), Chapter 8 (Settlements) and Chapter 9 (Other Land). Flooded Lands (Chapter 7) are not  
175 included in this *Wetlands Supplement*.

176

177 Guidance is given for  $CO_2$  emissions from the soil carbon pool in drained organic soils in line with the Section  
178 3.3., Chapter 2, Volume 4 in the *2006 IPCC Guidelines*. Guidance for changes in the carbon pools in above-  
179 ground and below-ground biomass, dead wood, and litter on these lands is provided in the *2006 IPCC Guidelines*  
180 and remains unchanged.

### 181 **CHOICE OF METHOD**

182 The most important factors considered for estimating on-site  $CO_2$  emissions and removals from drained inland  
183 organic soils are land-use and climate. Other factors such as nutrient status (or fertility) of the soil and drainage  
184 level affect emissions and can be considered where appropriate and with higher Tier methods. It is *good practice*

185 to stratify land-use categories by climate domain (Table 4.1, Chapter 4, Volume 4 of the *2006 IPCC Guidelines*),  
 186 nutrient status (*GPG-LULUCF* and Section 7.2.1.1, Chapter 7, Volume 4 of the *2006 IPCC Guidelines*) and  
 187 drainage class (shallow or deep) according to the stratification in Table 2.1.

188 Nutrient status is defined in *GPG-LULUCF* and *2006 IPCC Guidelines* (Section 7.2.1.1, Chapter 7, Volume).  
 189 Generally, ombrogenic organic soils are characterized as nutrient poor, while minerogenic organic soils are  
 190 characterized as nutrient rich. This broad characterization may vary by peatland type or national circumstances.

191 Drainage class is defined as the mean annual water table averaged over a period of several years; the shallow-  
 192 drained class is defined as the mean annual water table depth of less than 30 cm below the surface; the deep-  
 193 drained class is defined as the mean annual water table depth of 30 cm and deeper below the surface.

194 For Tier 1 methods, if the typical range of mean annual water table levels of drained inland organic soils for each  
 195 land-use category is unknown, the default is that the organic soil is deep-drained (water-table depth is specific  
 196 for land-use categories and climate domains) because deep-drained conditions are the most widespread and  
 197 suitable for a wide range of management intensities. Higher Tier methods could further differentiate the drainage  
 198 intensity within land-use categories if there are significant areas which differ from the default deep-drained  
 199 conditions.

200

201 Figure 2.5 in, Section 2.3.3, Chapter 2, Volume 4 of the *2006 IPCC Guidelines* provides the decision tree for  
 202 identification of the appropriate tier to estimate CO<sub>2</sub> emissions from drained inland organic soils by land-use  
 203 category.

### 204 **Tier 1**

205 The basic methodology for estimating annual carbon loss from drained inland organic soils was presented in  
 206 Section 2.3.3 and Equation 2.26 in Volume 4 of the *2006 IPCC Guidelines* as further specified in Equation 2.2.  
 207 Equation 2.3 refers to  $CO_2-C_{on-site}$  in Equation 2.2 with stratification of land-use categories by climate domain  
 208 and nutrient status. Nutrient status and drainage classes only need to be differentiated for those land-use  
 209 categories and climate domains for which emission factors are differentiated in Table 2.1.

210

211

212

**EQUATION 2.3**  
**ANNUAL ON-SITE CO<sub>2</sub>-C EMISSIONS/REMOVALS FROM DRAINED INLAND ORGANIC SOILS**  
**EXCLUDING EMISSIONS FROM FIRES**

$$CO_2-C_{on-site} = \sum_{c,n,d} (A \bullet EF)_{c,n,d}$$

213

214 Where:

215  $CO_2-C_{on-site}$  = Annual on-site CO<sub>2</sub>-C emissions/removals from drained inland organic soils in a land-use  
 216 category, tonnes C yr<sup>-1</sup>

217 A = Land area of drained inland organic soils in a land-use category in climate domain c, nutrient status n,  
 218 and drainage class d, ha

219 EF = Emission factors for drained inland organic soils, by climate domain c, nutrient status n, and  
 220 drainage class d, tonnes C ha<sup>-1</sup> yr<sup>-1</sup>

### 221 **Tier 2**

222 The Tier 2 approach for CO<sub>2</sub> emissions/removals from drained inland organic soils incorporates country-specific  
 223 information in Equations 2.2 and 2.3 to estimate the CO<sub>2</sub> emissions/removals. Tier 2 uses the same procedural  
 224 steps for calculations as provided for Tier 1. Improvements to the Tier 1 approach may include: 1) a derivation of  
 225 country-specific emission factors; 2) specification of climate sub-domains considered suitable for refinement of  
 226 emission factors; 3) a finer, more detailed classification of management systems with a differentiation of land-  
 227 use intensity classes; 4) a differentiation by drainage classes; or 5) a finer, more detailed classification of nutrient  
 228 status, e.g., by nitrogen, phosphorus or pH.

229 It is *good practice* to derive country-specific emission factors if measurements representing the national  
 230 circumstances are available. Methodologies and measurement techniques shall be compatible with the scientific  
 231 background for the Tier 1 emission factors in Annex 2A.1. Moreover, it is *good practice* for countries to use a  
 232 finer classification for climate and management systems, in particular drainage classes, if there are significant  
 233 differences in measured carbon loss rates among these classes. Note that any country-specific emission factor  
 234 must be accompanied by sufficient national or regional land-use/management activity and environmental data to  
 235 represent the appropriate climate sub-domains and management systems for the spatial domain for which the  
 236 country-specific emission factor is applied.

237 The general guidance of the *2006 IPCC Guidelines*, Section 2.3.3, Chapter 2, Volume 4 also applies here.

---

Final Draft**238 Tier 3**

239 CO<sub>2</sub> emissions/removals from drained inland organic soils can be estimated with model and/or measurement  
240 approaches. Dynamic, mechanistic models will typically be used to simulate underlying processes while  
241 capturing the influence of land-use and management, particularly the effect of seasonally variable levels of  
242 drainage on decomposition (van Huissteden *et al.*, 2006). The general considerations for organic soils in the  
243 Section 2.3.3, Chapter 2, Volume 4 of the *2006 IPCC Guidelines* also apply here. It is *good practice* to describe  
244 the methodologies and models transparently, document the considerations for choosing and applying the model  
245 in the inventory and provide evidence that it represents the national circumstances according to the guidance in  
246 Section 5, Chapter 2, Volume 4 of the *2006 IPCC Guidelines*.

**247 CHOICE OF EMISSION/REMOVAL FACTORS****248 Tier 1**

249 All Tier 1 emission factors have been updated from the *2006 IPCC Guidelines* based on a large number of new  
250 measurement data in all land-use categories and climate zones. The new evidence allows for stratification of  
251 more land-use categories and climate domains by nutrient status than in the *2006 IPCC Guidelines*. In addition,  
252 temperate, nutrient-rich Grassland is further stratified into shallow-drained (less than approximately 30 cm below  
253 surface) and deep-drained. Within each land-use category, drained inland organic soils can experience a wide  
254 range of mean annual water table levels that depend upon regional climatic characteristics and specific land-use  
255 activity or intensity. For temperate Grassland EFs are given for shallow-drained and deep-drained soils. The  
256 shallow-drained and deep-drained Grassland emission factors differ significantly. Without additional national  
257 information about mean annual water table and/or land-use intensity as proxy, countries should choose deep-  
258 drained as default.

259 The *GPG-LULUCF* and *2006 IPCC Guidelines* (Section 7.2.1.1, Chapter 7, Volume 4) distinguish between  
260 nutrient-rich and nutrient-poor organic soils in some land-use categories and climate zones. This approach is  
261 maintained here, in line with guidance in the *2006 IPCC Guidelines*.

262 Default Tier 1 emission/removal factors for drained inland organic soils (Table 2.1) were generated using a  
263 combination of subsidence and flux data found in the literature as described in Annex 2A.1. CO<sub>2</sub>-C losses occur  
264 predominantly in the drained, oxic soil layer and thus reflect human-induced CO<sub>2</sub>-C fluxes. The part of the soil  
265 profile affected by drainage can be deeper or shallower than the default 0 to 30 cm layer considered in the Tier 1  
266 default methodology for SOC pools in mineral soils.

267 .

**TABLE 2.1**  
**TIER 1 CO<sub>2</sub> EMISSION/REMOVAL FACTORS FOR DRAINED ORGANIC SOILS IN ALL LAND-USE CATEGORIES\*\***

Land-use category		Climate / vegetation zone	Emission Factor** (tonnes CO <sub>2</sub> -C ha <sup>-1</sup> yr <sup>-1</sup> )	95% Confidence Interval		No. of sites	Citations/comments
Forest Land, drained	Nutrient-poor	Boreal	0.39	0.0	0.86	64	Komulainen <i>et al.</i> , 1999; Lohila <i>et al.</i> , 2011; Minkkinen & Laine, 1998; Minkkinen <i>et al.</i> , 1999; Ojanen <i>et al.</i> , 2010, 2013; Simola <i>et al.</i> , 2012
	Nutrient-rich	Boreal	1.0	0.68	1.3	78	Komulainen <i>et al.</i> , 1999; Laurila <i>et al.</i> , 2007; Lohila <i>et al.</i> , 2007; Mäkiranta <i>et al.</i> , 2007; Minkkinen & Laine, 1998; Minkkinen <i>et al.</i> , 1999, 2007b; Ojanen <i>et al.</i> , 2010, 2013; Simola <i>et al.</i> , 2012
Forest Land, drained		Temperate	2.6	2.0	3.3	8	Glenn <i>et al.</i> , 1993; Minkkinen <i>et al.</i> , 2007b; Von Arnold <i>et al.</i> , 2005a,b, Yamulki <i>et al.</i> , 2013
Forest Land and cleared Forest Land (shrubland***), drained		Tropical	5.3	-0.7	9.5	n/a	Ali <i>et al.</i> , 2006; Brady, 1997; Chimner & Ewel, 2005; Comeau <i>et al.</i> , 2013; Dariah <i>et al.</i> , 2013; Darung <i>et al.</i> , 2005; Furukawa <i>et al.</i> , 2005; Hadi <i>et al.</i> , 2005; Harisson <i>et al.</i> , 2007; Hergoualc'h & Verchot, 2011; Hertel <i>et al.</i> , 2009; Hirano <i>et al.</i> , 2009, 2012; Inubushi <i>et al.</i> , 2003; Ishida <i>et al.</i> , 2001; Jauhiainen <i>et al.</i> , 2008, 2012a; Melling <i>et al.</i> , 2005a, 2007a; Rahaoje <i>et al.</i> , 2000; Shimamura & Momose, 2005; Sulistiyanto, 2004; Sundari <i>et al.</i> , 2012
Forest plantations, drained****		Tropical	20	16	24	n/a	Basuki <i>et al.</i> , 2012; Hooijer <i>et al.</i> , 2012; Jauhiainen <i>et al.</i> , 2012a; Nouvellon <i>et al.</i> , 2012; Warren <i>et al.</i> , 2012
Grassland, drained		Boreal	5.7	2.9	8.6	8	Grönlund <i>et al.</i> , 2006; Kreshtapova & Maslov, 2004; Lohila <i>et al.</i> , 2004; Maljanen <i>et al.</i> , 2001a, 2004; Nykänen <i>et al.</i> , 1995; Shurpali <i>et al.</i> , 2009
Grassland, drained, nutrient-poor		Temperate	5.3	3.7	6.9	7	Kuntze, 1992; Drösler <i>et al.</i> , 2013

Final Draft

Grassland, deep-drained, nutrient- rich	Temperate	6.1	5.0	7.3	39	Augustin, 2003; Augustin <i>et al.</i> , 1996; Czaplak & Dembek, 2000; Drösler <i>et al.</i> , 2013; Elsgaard <i>et al.</i> , 2012; Jacobs <i>et al.</i> , 2003; Kasimir-Klemedtsson <i>et al.</i> , 1997; Langeveld <i>et al.</i> , 1997; Leifeld <i>et al.</i> , 2011; Lorenz <i>et al.</i> , 2002; Meyer <i>et al.</i> , 2001; Nieveen <i>et al.</i> , 2005; Okruszko 1989 (Höper, 2002); Schothorst, 1977 (Höper, 2002), Schrier-Uijl, 2010a, c; Veenendaal <i>et al.</i> , 2007; Weinzierl, 1997 (Höper, 2002)
Grassland, shallow drained, nutrient-rich	Temperate	3.6	1.8	5.4	13	Drösler <i>et al.</i> , 2013; Jacobs <i>et al.</i> , 2003 ; Lloyd, 2006
Grassland, drained	Tropical	9.6	4.5	17	n/a.	Updated from Table 6.3, Chapter 6, Volume 4, 2006 IPCC Guidelines *****
Cropland, drained	Boreal & Temperate	7.9	6.5	9.4	39	Drösler <i>et al.</i> , 2013; Elsgaard <i>et al.</i> , 2012, Grønlund <i>et al.</i> , 2008, Kasimir-Klemedtsson <i>et al.</i> , 1997; Leifeld <i>et al.</i> , 2011; Maljanen <i>et al.</i> , 2001a, 2003a, 2004, 2007a; Morrison <i>et al.</i> , 2013b, Petersen <i>et al.</i> 2012
Cropland and fallow, drained	Tropical	14	6.6	26	n/a	Ali <i>et al.</i> , 2006; Chimner, 2004; Chimner & Ewel, 2004; Dariah <i>et al.</i> , 2013; Darung <i>et al.</i> , 2005; Furukawa <i>et al.</i> , 2005; Gill and Jackson, 2000; Hairiah <i>et al.</i> , 2000; Hirano <i>et al.</i> , 2009; Ishida <i>et al.</i> , 2001; Jauhainen <i>et al.</i> , 2012; Melling <i>et al.</i> , 2007a;
Cropland, drained – paddy rice	Tropical	9.4	-0.2	20	n/a	Dariah <i>et al.</i> , 2013; Furukawa <i>et al.</i> , 2005; Hadi <i>et al.</i> , 2005; Hairiah <i>et al.</i> , 1999; Inubushi <i>et al.</i> , 2003; Ishida <i>et al.</i> , 2001; Matthews <i>et al.</i> , 2000; Melling <i>et al.</i> , 2007a
Agriculture – oil palm	Tropical	11	5.6	17	n/a	Comeau <i>et al.</i> , 2013; Dariah <i>et al.</i> , 2013; DID and LAWOO, 1996; Henson and Dolmat, 2003; Couwenberg, and Hooijer 2013; Lamade and Bouillet, 2005; Marwanto and Agus, 2013; Melling <i>et al.</i> , 2005a, 2007a, 2013; Warren <i>et al.</i> , 2012
Agriculture – sago palm (shallow drained)	Tropical	1.5	-2.3	5.4	n/a	Dariah <i>et al.</i> , 2013; Hairiah <i>et al.</i> , 1999; Ishida <i>et al.</i> , 2001; Lamade and Bouillet, 2005; Matthews <i>et al.</i> , 2000; Melling <i>et al.</i> , 2005a, 2007a; Watanabe <i>et al.</i> , 2009

Peatland Managed for Extraction	Boreal & Temperate	2.8	1.1	4.2	21	Ahlholm and Silvola 1990; Glatzel <i>et al.</i> , 2003, Hargreaves <i>et al.</i> , 2003; McNeil and Waddington 2003; Shurpali <i>et al.</i> , 2008; Strack and Zuback 2013; Sundh <i>et al.</i> , 2000; Tuittila and Komulainen, 1995; Tuittila <i>et al.</i> , 1999; 2004, Waddington <i>et al.</i> , 2010
Peat Extraction	Tropical	2.0	0.06	7.0	n/a.	Table 7.4, Chapter 7, Volume 4, 2006 IPCC Guidelines
Settlements	All climate zones	There is no fixed default emission/removal factor for Settlement. It is <i>good practice</i> to take the default emission/removal factor in Table 2.1 of the land-use category that is closest to the national conditions of drained organic soils under Settlements. Information about national conditions could include drainage level, vegetation cover, or other management activities. For example, drained organic soils in urban green areas, parks or gardens could use the default Tier 1 emission/removal factor for Grassland, deep-drained in Table 2.1.				
Other Land	All climate zones	<i>Other Land Remaining Other Land: 0</i> <i>Land Converted to Other Land: Maintain emission factor of previous land-use category</i>				
<p>** Mean</p> <p>*** Shrubland refers to any type of land sparsely or fully covered with shrubs or trees, which may or may not fulfil the national forest definition. It extends to degraded lands, which cannot be clearly classified as forest or non-forest.</p> <p>**** Number derived solely from Acacia plantation data.</p> <p>***** The emission factor for Cropland in the tropical zone was multiplied with the ratio between the emission factors for Grassland, drained, nutrient-poor and Cropland for the temperate zone; same for confidence interval. This new ratio updates the ratio applied to derive the emission factor for Grassland in the tropical zone in Table 6.3, Chapter 6, Volume 4, 2006 IPCC Guidelines.</p>						

269

270

---

 Final Draft

271 Common tropical plantations include oil palm, sago and *Acacia crassiparva*. In Table 2.1, plantations for food  
 272 and oil crops like sago and oil palm are classified under Cropland, while fibre plantations like Acacia are  
 273 classified as Forest Land. It is *good practice* to report plantations in the appropriate national land-use category  
 274 according to the national forest definition.

## 275 **Tier 2**

276 The Tier 2 approach for carbon loss from drained inland organic soils incorporates country-specific information  
 277 in Equation 2.2 to estimate the emissions. Also, Tier 2 uses the same procedural steps for calculations as  
 278 provided for Tier 1. Tier 2 emission factors by land-use category can, in general, be developed depending on a)  
 279 climate, b) drainage layout and intensity, c) nutrient status and d) land-use intensity and practices.

280 Tier 2 emission factors could include the following refinements:

- 281
- 282 • Use of country specific emission factors measured or calculated locally taking into account climatic factors  
 283 that provide for wetter or drier drainage classes than those defined here;
- 284 • Use of country specific emission factors measured or calculated locally taking into account slope factors  
 285 (e.g., blanket bogs) that may promote wetter or drier drainage classes than those defined here;
- 286 • Derivation of emission factors for boreal Forest Land by nutrient status (rich/poor) if the two EFs are  
 287 significantly different (See Table 2.1);
- 288 • Development of boreal and temperate Grassland emission factors according to land-use intensity, for  
 289 example to distinguish high-intensity (fertilized, ploughed and reseeded) Grassland from low-intensity  
 290 permanent Grassland, or moorland rough grazing (grazing by hardier breeds of sheep) on drained blanket  
 291 bogs.

292 CO<sub>2</sub> measurements by methods described in Annex 2A.1, disaggregated by management practices, should be  
 293 used to develop more precise, locally appropriate emission factors.

## 294 **Tier 3**

295 A Tier 3 approach allows for a variety of methods and might use measurements or process-based models or other  
 296 more elaborate approaches, adequately validated using observation data that take into account temporal and  
 297 spatial variations. Tier 3 should involve a comprehensive understanding and representation of the dynamics of  
 298 CO<sub>2</sub> emissions and removals on managed organic soils, including the effect of management practices, site  
 299 characteristics, peat type and depth, drainage depth, etc. Tier 3 approaches could start by developing  
 300 relationships between drainage or nutrient status and heterotrophic CO<sub>2</sub> emissions, which can be further refined  
 301 by land-use category and fertilization. Furthermore, forested organic soils undergo a cycle related to rotation of  
 302 the tree cohorts and carbon losses associated with harvesting and site preparation should be accounted. Models  
 303 could describe the rotational variation in water tables.

304 When peat is extracted, the peatland surface is disturbed by machinery and may be fertilized afterwards or  
 305 otherwise amended for regeneration. Moreover, drainage systems may be renewed and dredging of ditches may  
 306 cause disturbances that alter the greenhouse gas emissions and removals. These measures result in  
 307 emission/removal rates that vary predictably over time, which may in Tier 3 methods be captured by models  
 308 used. Emissions from stockpiles of drying peat are much more uncertain. Higher temperatures may cause  
 309 stockpiles to release more CO<sub>2</sub> than the excavation field, but data are not at present sufficient to provide  
 310 guidance. Methods for estimating this emission may be developed at Tier 3.

## 311 **CHOICE OF ACTIVITY DATA**

312 All management practices for land remaining in a land-use category are assumed to result in persistent emissions  
 313 from soils as long as the management system remains in place or as long as the land falls under the definition of  
 314 organic soils. Activity data consist of areas of land remaining in a land-use category on organic soils stratified by  
 315 climate domains, soil nutrient status, drainage class or additional criteria such as management practices. Total  
 316 areas should be determined according to approaches laid out in Chapter 3, Volume 4 of the *2006 IPCC*  
 317 *Guidelines* and should be consistent with those reported under other sections of the inventory. The estimation of  
 318 CO<sub>2</sub> emissions/removals from drained inland organic soils will be greatly facilitated if this information can be  
 319 used in conjunction with national soils and climate data, vegetation inventories, and other biophysical data.  
 320 Stratification of land-use categories according to climate domains, based on default or country-specific  
 321 classifications can be accomplished with overlays of land-use on suitable climate and soil maps.

322 Under most circumstances, the area of organic soils will remain constant over time. However, the area of  
 323 organic soils may change as organic soil disappears following drainage.

**Tier 1**

324 **Tier 1**  
325 The Tier 1 approach requires area data of managed land with organic soils for each land-use category,  
326 disaggregated by appropriate climate domains, nutrient status and drainage class as applicable. Classification  
327 systems for activity data that form the basis for a Tier 1 inventory are provided in the respective land-use  
328 chapters of the *2006 IPCC Guidelines*.

329 Several institutions, including ISRIC and FAO have country-specific and global maps that include organic soils  
330 (<http://www.fao.org/geonetwork/srv/en/main.home> or <http://www.isric.org/>). A global consortium has been  
331 formed to make a new digital soil map of the world at fine resolution (<http://www.globalsoilmap.net/>).

332 The *GPG-LULUCF* and *2006 IPCC Guidelines* (Section 7.2.1.1, Chapter 7, Volume 4) distinguish between  
333 nutrient-rich and nutrient-poor organic soils in some land-use categories and climate zones. This approach is  
334 maintained here, in line with guidance in the *2006 IPCC Guidelines*. Nutrient-poor organic soils predominate in  
335 boreal regions, while in temperate regions nutrient-rich organic soils are more common. It is *good practice* that  
336 boreal countries that do not have information on areas of nutrient-rich and nutrient-poor organic soils should use  
337 the emission factor for nutrient-poor organic soils. It is *good practice* that temperate countries that do not have  
338 such data use the emission factor for nutrient-rich organic soils. Only one default factor is provided for tropical  
339 regions, so disaggregating by soil fertility is not necessary in the tropical climate zone using the Tier 1 method.  
340 Due to lack of data, rice fields on tropical organic soils are not disaggregated by water management regimes.

341 The areas of shallow-drained and deep-drained inland organic soils with Grasslands need to be derived from  
342 national data. Data from water management plans, such as target water table levels can serve as a source of  
343 information. Land-use intensity, e.g., the time of the first cut of Grassland, grazing intensity or animal production  
344 levels can serve as a proxy as well as restrictions imposed by water management or biodiversity management  
345 (e.g. riparian zones, buffer zones, nature conservation for species or habitats with typical water regime).

346 Without additional national information about mean annual water table and/or land-use intensity as proxy,  
347 countries should choose deep-drained as the default.

**Tier 2 and 3**

348 **Tier 2 and 3**  
349 Activity data for higher Tier estimates are generally derived following the methods presented in Chapter 3 of the  
350 *2006 IPCC Guidelines*. Activity data may be spatially explicit and could be disaggregated by type of  
351 management, drainage depth, and/or nutrient status to improve the accuracy of the inventory if different land  
352 management systems use different drainage depths and/or nutrient levels, and if appropriate emissions factors are  
353 available. In general, practices that increase carbon stocks in mineral soils by increased organic material input  
354 (fertilization, liming, etc.) do not have a sequestration effect in drained organic soils.

355 The combination of land-use databases and soil maps or spatially explicit data allow delineation of combinations  
356 of land-use categories, climate domains, drainage classes and management systems and their changes over time  
357 on organic soils. Data and their documentation could combine information from a land-use transition matrix  
358 specifically made for organic soils. Stratification needs to be consistently applied across the entire time series.

359 Information sources about drainage with adequate disaggregation may include:

- 360 • National land-use statistics, land-use maps and soil maps, maps of water and nature conservation zones with  
361 restrictions for water management, wetlands.
- 362 • National water management statistics: in most countries, the agricultural land base including Cropland is  
363 usually surveyed regularly, providing data on distribution of different land-uses, crops, tillage practices and  
364 other aspects of management, often at sub-national regional level. These statistics may originate, in part,  
365 from remote sensing methods, from which additional information about wetness or periods with seasonal  
366 flooding could be extracted.
- 367 • Inventory data from a statistically-based, plot-sampling system of water table wells, ditches and surface  
368 waters on organic soils: water table is monitored at specific permanent sample plots either continuously or  
369 on plots that are revisited on a regular basis. It has to be documented that the water data represent the water  
370 table in the organic soil and for what land-use and drainage stratum and that the data cover a representative  
371 period, which represents a multi-year mean annual water table.
- 372 • Water management plans and documentation from water management installations.
- 373 • Drainage maps.
- 374 • Maps of rewetting projects including remote sensing.

Final Draft

## 375 **CALCULATION STEPS FOR TIER 1**

376 The steps for estimating the direct loss of soil carbon from drained inland organic soils are as follows:

377 **Step 1:** Determine areas with drained inland organic soils under each land-use category for land remaining in a  
378 land use category, disaggregated by climate domain and other appropriate factors as outlined above. Where  
379 needed for Tier 1 emission factors, land areas are further stratified by nutrient-rich and nutrient-poor organic  
380 soils. Temperate nutrient-rich Grassland is further stratified into shallow-drained and deep-drained classes.

381 **Step 2:** Assign the appropriate emission factor (EF) from Table 2.1 for annual losses of CO<sub>2</sub> to each land-use  
382 category, climate domain, nutrient status and drainage class stratum.

383 **Step 3:** Multiply each area with the appropriate emission factor using Equations 2.3.

## 384 **UNCERTAINTY ASSESSMENT**

385 Three broad sources of uncertainty exist in estimating emissions and removals in organic soils: 1) uncertainties  
386 in land-use and management activity and environmental data; 2) uncertainties in the emission/removal factors for  
387 Tier 1 or 2 approaches; and 3) model structure/parameter error for Tier 3 model-based approaches, or  
388 measurement error/sampling variability associated with Tier 3 measurement-based inventories. In general,  
389 precision of an inventory is increased and confidence ranges are smaller with more sampling to estimate values  
390 for land-use categories, while accuracy is more likely to be increased through implementation of higher Tier  
391 methods that incorporate country-specific information.

392 For Tier 1, the default uncertainty level of emissions/removal factors is the 95% confidence interval in Table 2.1.  
393 Countries developing specific emission factors for their inventories at higher tiers should assess the uncertainty  
394 of these factors.

395 If using aggregate land-use area statistics for activity data (e.g., FAO data), the inventory agency may have to  
396 apply a default level of uncertainty for the land area estimates on organic soils ( $\pm 20\%$ ; twice the uncertainty  
397 estimate in Table 3.7 for mineral soils in the 2006 Guidelines). It is *good practice* for the inventory compiler to  
398 derive uncertainties from country-specific activity data instead of using a default level of uncertainty.  
399 Uncertainties in activity data may be reduced through a better monitoring system, such as developing or  
400 extending a ground-based survey with additional sample locations and/or incorporating remote sensing to  
401 provide additional coverage. Uncertainties in activity data and emission/removal factors need to be combined  
402 using an appropriate method, such as simple error propagation equations. Details are given in Chapter 3, Volume  
403 1 of the 2006 IPCC Guidelines and Chapter 5 of GPG-LULUCF.

404 Accuracy can be increased by deriving country-specific factors using a Tier 2 method or by developing a Tier 3  
405 country-specific estimation system. The underlying basis for higher tier approaches will be measurements in the  
406 country or neighbouring regions that address the effect of land-use and management on CO<sub>2</sub> emissions/removals  
407 from drained organic soils. In addition, uncertainties can be reduced through stratification by significant factors  
408 responsible for within-country differences in land-use and management impacts, such as variation among climate  
409 domains and/or organic soil types.

### 410 **2.2.1.2 OFF-SITE CO<sub>2</sub> EMISSIONS VIA WATERBORNE CARBON** 411 **LOSSES FROM DRAINED ORGANIC SOILS**

412 Waterborne carbon comprises dissolved organic carbon (DOC), particulate organic carbon (POC), the dissolved  
413 gases CO<sub>2</sub> and CH<sub>4</sub>, and the dissolved carbonate species HCO<sub>3</sub><sup>-</sup> and CO<sub>3</sub><sup>2-</sup>. Particulate inorganic carbon (PIC)  
414 losses are negligible from organic soils. Collectively, waterborne carbon export can represent a major part of the  
415 overall carbon budget of an organic soil, and in some cases can exceed the net land-atmosphere CO<sub>2</sub> exchange  
416 (e.g., Billett *et al.*, 2004; Rowson *et al.*, 2010). It is therefore important that waterborne carbon is included in  
417 flux-based (i.e., gain-loss) approaches for soil carbon estimation, to avoid systematic under-estimation of soil  
418 carbon losses. Airborne (erosional) POC loss may also be significant where land-use leads to bare soil exposure,  
419 but few data exist to quantify this (see Appendix 2a.1).

420 Different forms of waterborne carbon have different sources, behaviour and fate, and different approaches are  
421 therefore required to quantify the off-site CO<sub>2</sub> emissions associated with each form. In most peatlands and  
422 organic soils, DOC forms the largest component of waterborne carbon export (e.g., Urban *et al.*, 1989; Dawson  
423 *et al.*, 2004; Jonsson *et al.*, 2007; Dinsmore *et al.*, 2010). DOC export can be affected by land-use, in particular  
424 drainage (Wallage *et al.*, 2006; Strack *et al.*, 2008; Urbanová *et al.*, 2011; Moore *et al.*, 2013). It is reactive  
425 within aquatic ecosystems and most DOC is thought to be ultimately converted to CO<sub>2</sub> and emitted to the

426 atmosphere (see Annex 2A.2 for supporting discussion). Therefore, it is *good practice* to include DOC in flux-  
427 based carbon estimation methods, and a Tier 1 methodology is described below.

428 Of the other forms of waterborne carbon, POC fluxes are typically very low from vegetated peatlands and  
429 organic soils, but can become very large where bare organic soil becomes exposed, e.g., due to erosion, peat  
430 extraction, burning and conversion to Cropland. Although it may be possible to estimate POC loss fluxes as a  
431 function of bare soil exposure, high uncertainty remains regarding the reactivity and fate of POC exported from  
432 organic soils. Some POC is likely to be converted to CO<sub>2</sub>, but POC that is simply translocated from the soil  
433 profile to other stable carbon stores, such as freshwater or marine sediments, may not lead to CO<sub>2</sub> emissions. Due  
434 to the uncertain fate of POC export, an estimation method is not presented at this time; current knowledge and  
435 data needs to support POC estimation in future are described in Appendix 2a.1.

436 Gaseous CO<sub>2</sub> and CH<sub>4</sub> dissolved in water transported laterally from the organic soil matrix represent indirectly  
437 emitted components of the total emission of these gases from the land surface. Dissolved CO<sub>2</sub> in excess of  
438 atmospheric pressure will also be degassed from drainage waters, whilst some dissolved inorganic carbon (DIC)  
439 may be transported downstream. At present, available data are insufficient (particularly from drained organic  
440 soils) to permit default emission factors to be derived. Additional information and future methodological  
441 requirements to support full accounting of emissions associated with waterborne inorganic carbon are included in  
442 Appendix 2a.1.

#### 443 CHOICE OF METHOD

444 The basic methodology for estimating annual off-site CO<sub>2</sub> emissions associated with waterborne carbon loss  
445 from drained organic soils is presented in Equation 2.4:

446

447 **EQUATION 2.4**  
448 **ANNUAL OFF-SITE CO<sub>2</sub> EMISSIONS DUE TO DOC LOSS FROM DRAINED ORGANIC SOILS (CO<sub>2</sub>)**

$$449 \quad CO_2-C_{DOC} = \sum_{c,n} (A \cdot EF_{DOC})_{c,n}$$

450 Where:

451 CO<sub>2</sub>-C<sub>DOC</sub> = Annual off-site CO<sub>2</sub>-C emissions due to DOC loss from drained organic soils, tonnes C yr<sup>-1</sup>

452 A<sub>c,n</sub> = Land area of drained organic soils in a land-use category in climate zone c and nutrient status n, ha

453 EF<sub>DOCc,n</sub> = Emission factors for annual CO<sub>2</sub> emissions due to DOC loss from drained organic soils, by  
454 climate zone c and nutrient status n, tonnes C ha<sup>-1</sup> yr<sup>-1</sup>

455 EF<sub>DOC</sub> can be calculated from Equation 2.5:

456

457 **EQUATION 2.5**  
458 **EMISSION FACTOR FOR ANNUAL CO<sub>2</sub> EMISSIONS DUE TO DOC EXPORT FROM DRAINED ORGANIC**  
459 **SOILS**

$$460 \quad EF_{DOC} = DOC_{FLUX\_NATURAL} \cdot (1 + \Delta DOC_{DRAINAGE}) \cdot Frac_{DOC-CO_2}$$

461 Where:

462 EF<sub>DOC</sub> = Emission factor for DOC from a drained site, tonnes C ha<sup>-1</sup> yr<sup>-1</sup>

463 DOC<sub>FLUX\_NATURAL</sub> = Flux of DOC from natural (undrained) organic soil, tonnes C ha<sup>-1</sup> yr<sup>-1</sup>

464 ΔDOC<sub>DRAINAGE</sub> = Proportional increase in DOC flux from drained sites relative to un-drained sites

465 Frac<sub>DOC-CO<sub>2</sub></sub> = Conversion factor for proportion of DOC converted to CO<sub>2</sub> following export from site

466 Because of the lack of data for other components of waterborne carbon fluxes and uncertainty about their sources  
467 and/or fate, off-site CO<sub>2</sub> emissions associated with waterborne carbon are only represented by DOC losses at this  
468 stage. However, if in the future adequate data become available or if adequate data are available for higher tiers,  
469 inventory compilers can expand Equation 2.4 to include POC and/or DIC (See section on methodological  
470 requirements in Appendix 2a.1).  
471  
472

Final Draft

473 **CHOICE OF EMISSION FACTOR**474 **Tier 1**

475 A detailed description of the derivation of default values for Tier 1 is provided in Annex 2A.2. In summary,  
 476 measurements show clear differentiation of natural DOC fluxes between boreal, temperate and tropical organic  
 477 soils, and Tier 1 emission factors therefore follow a broad classification based on climate zones. Annex 2A.2  
 478 provides details and data sources for the derivation of parameter values. Note that a single default value for  
 479  $\Delta\text{DOC}_{\text{DRAINAGE}}$  is currently proposed for all organic soil/land-use types, based on data from a range of studies  
 480 undertaken in different climate zones. A substantial body of scientific evidence indicates a high conversion of  
 481 organic soil-derived DOC to  $\text{CO}_2$  in aquatic systems, on which basis a default  $\text{Frac}_{\text{DOC-CO}_2}$  value of 0.9 ( $\pm 0.1$ ) is  
 482 proposed (see Annex 2A.2).

**TABLE 2.2**  
**DEFAULT DOC EMISSION FACTORS FOR DRAINED ORGANIC SOILS**

Climate zone	$\text{DOC}_{\text{FLUX\_NATURAL}}$ (t C ha <sup>-1</sup> yr <sup>-1</sup> )	$\Delta\text{DOC}_{\text{DRAINAGE}}$ <sup>a</sup>	$\text{EF}_{\text{DOC\_DRAINED}}$ (t C ha <sup>-1</sup> yr <sup>-1</sup> )
Boreal	0.08 (0.06-0.11)	0.60 (0.43-0.78)	0.12 (0.07-0.19)
Temperate	0.21 (0.17-0.26)		0.31 (0.19-0.46)
Tropical	0.57 (0.49-0.64)		0.82 (0.56-1.14)

Values shown in parentheses represent 95% confidence intervals. For data sources and supporting references see Tables 2A.2 and 2A.3.

<sup>a</sup> Due to the limited number of available studies, a single Tier 1 value for  $\Delta\text{DOC}_{\text{DRAINAGE}}$  has been assigned to all soil types based on all available comparisons of drained and undrained sites. For fens, there is more uncertainty associated with the estimation of DOC flux changes after drainage, therefore countries may choose to apply values of  $\text{DOC}_{\text{FLUX\_NATURAL}}$  given above (multiplied by  $\text{Frac}_{\text{DOC-CO}_2}$  but assuming  $\Delta\text{DOC}_{\text{DRAINAGE}} = 0$ ) or to obtain direct measurements of the DOC flux from drained sites.

483

484 **Tier 2**

485 A Tier 2 approach for estimation of DOC may follow the Tier 1 methodology provided above, but should use  
 486 country-specific information where possible to refine the emission factors used. Possible refinements where  
 487 supporting data are available could include:

- 488 • Use of country-level measurements from natural organic soils to obtain accurate values of  $\text{DOC}_{\text{FLUX\_NATURAL}}$   
 489 for that country, for example by developing specific values for raised bogs versus fens, or for blanket bogs;
- 490 • Use of country-level data on the impacts of organic soil drainage on DOC flux to derive specific values of  
 491  $\Delta\text{DOC}_{\text{DRAINAGE}}$  that reflect local organic soil types, and the nature of drainage practices and subsequent land-  
 492 use. If sufficient, robust, direct measurements are available from representative drained sites, these may be  
 493 used to estimate DOC fluxes from drained sites, replacing  $\text{DOC}_{\text{FLUX\_NATURAL}}$  in Equation 2.5. Specific DOC  
 494 flux estimates from drained organic soils in different land-use categories could also be considered where  
 495 data support this level of stratification;
- 496 • Use of alternative values for  $\text{Frac}_{\text{DOC-CO}_2}$  where evidence is available to estimate the proportion of DOC  
 497 exported from drained organic soils that is transferred to stable long-term carbon stores, such as lake or  
 498 marine sediments.

499 **Tier 3**

500 A Tier 3 approach might include the use of more detailed data to develop and apply process models that describe  
 501 DOC release as a function of vegetation composition, nutrient levels, land-use category, water table level and  
 502 hydrology, as well as temporal variability in DOC release in the years following land-use change (e.g. initial  
 503 drainage) and on-going management activity (e.g., drain maintenance, forest management) (see Annex 3A.2,  
 504 Chapter 3).

505 Guidance is not currently presented for the effects of land-use impacts other than drainage on DOC loss from  
 506 peatlands and organic soils, for example the effects of managed burning or intensity of agricultural use. However,  
 507 these may be included in higher tier methods if sufficient evidence can be obtained to develop the associated  
 508 emission factors.

509 **CHOICE OF ACTIVITY DATA**510 **Tier 1**

511 Activity data consist of areas of land remaining in a land-use category on drained organic soils summarised by  
512 organic soil type, climate zones and land-use type (specifically occurrence of drainage). Total areas should be  
513 determined according to Approaches laid out in Chapter 3 of the *2006 IPCC Guidelines* and should be consistent  
514 with those reported under other sections of the inventory. They also need to be consistent with activity data for  
515 on-site CO<sub>2</sub> emissions. For boreal and temperate raised bogs and fens, additional data on annual mean  
516 precipitation may be used to refine emission estimates, as shown in Table 2.2.

### 517 **Tier 2 and 3**

518 For higher Tier approaches, additional activity data requirements may include specific information on the land-  
519 use type associated with drained organic soils, and intensity of drainage. Use of a variable  $Frac_{DOC-CO_2}$  value at a  
520 country level, or within a country, would require information on the characteristics of downstream river networks  
521 (e.g., water residence time, extent of lakes and reservoirs, lake sedimentation rates). A Tier 3 modelling approach  
522 could include additional information on the timing of drainage, drain maintenance and land-management (e.g.,  
523 forest management, influence of fertiliser application rates on DOC production).

### 524 **CALCULATION STEPS FOR TIER 1**

525 The steps for estimating the off-site emissions from soil carbon on drained inland organic soils are as follows:

526 **Step 1:** Determine areas with drained inland organic soils under each land-use category for land remaining in a  
527 land-use category, disaggregated by climate domain and other appropriate factors as outlined above.

528 **Step 2:** Assign the appropriate values for  $DOC_{FLUX\_NATURAL}$ ,  $\Delta DOC_{DRAINAGE}$   $Frac_{DOC-CO_2}$  from Table 2.2 for each  
529 land-use category and climate domain.

530 **Step 3:** Calculate  $EF_{DOC}$  for each land-use category using Equation 2.5

531 **Step 4:** Multiply activity data by the emission factor for each land-use category and sum across land-use  
532 categories.

### 533 **UNCERTAINTY ASSESSMENT**

534 Three broad sources of uncertainty exist in estimating off-site emissions and removals: 1) uncertainties in land-  
535 use and management activity and environmental data; 2) uncertainties in the emission/removal factors for Tier 1  
536 or 2 approaches; and 3) uncertainties in the fraction of DOC that is emitted as CO<sub>2</sub>. In general, precision of an  
537 inventory is increased and confidence ranges are smaller with more sampling to estimate values for these  
538 categories, while accuracy is more likely to be increased through implementation of higher tier methods that  
539 incorporate country-specific information.

540 Uncertainties for land use and management activities are the same as for on-site emissions and will not be  
541 repeated here. Uncertainty ranges (95% confidence intervals) are provided for DOC emission factors in Table 2.2.  
542 These ranges are calculated from literature data in Annex 2A.2 based on observations from natural peatlands  
543 used to derive values of  $DOC_{FLUX-NATURAL}$  in each of the peat classes used (Table 2A.2); observations of  
544  $\Delta DOC_{DRAINAGE}$  from published studies (Table 2A.3); and an uncertainty range for  $Frac_{DOC-CO_2}$  value of 0.8 to 1.0  
545 as described above. These uncertainty ranges may be adapted or refined under Tier 2 if further sub-classification  
546 according to land-use type or intensity is undertaken, based on additional measurement data.

## 547 **2.2.2 Non-CO<sub>2</sub> emissions and removals from drained** 548 **inland organic soils**

549 In the *2006 IPCC Guidelines*, CH<sub>4</sub> emissions were assumed to be negligible from all drained organic soils. Here  
550 new methodologies and emission factors are provided for soil CH<sub>4</sub> emissions from drained organic soils and  
551 drainage ditches (Section 2.2.2.1).

### 552 **2.2.2.1 CH<sub>4</sub> emissions and removals from drained organic** 553 **soils**

554 In the *2006 IPCC Guidelines*, CH<sub>4</sub> emissions were assumed to be negligible from all drained organic soils.  
555 However, recent evidence suggests that some CH<sub>4</sub> emissions can occur from the drained land surface, and also  
556 from the ditch networks constructed during drainage. Each of these emission pathways is considered here (Best  
557 and Jacobs, 1997; Minkinen and Laine 2006; Schrier-Uijl *et al.*, 2011; Hyvönen *et al.*, 2012).

Final Draft

558 Drainage lowers the water table and exposes formerly saturated organic soil layers to oxidation and, as described  
 559 above, increases CO<sub>2</sub> emissions from the land surface. Drainage alters environmental factors such as temperature,  
 560 reduction–oxidation potential, and the amount of easily decomposable organic matter. Drainage also affects the  
 561 activity of methanogens and methanotrophs (Blodau, 2002; Treat *et al.*, 2007). Drainage increases plant root  
 562 respiration and mitigates CH<sub>4</sub> emission dramatically (Martikainen *et al.*, 1995a; Strack *et al.*, 2004; Hergoualc'h  
 563 and Verchot, 2012) as the methanogenic bacteria thrive only in anoxic conditions. Shifts in vegetation with  
 564 dominant aerenchymous species to other vegetation types will also reduce the transfer of methane from the soil  
 565 profile to the atmosphere (e.g., Tuittila *et al.*, 2000). In general, when the organic soil is drained the natural  
 566 production of CH<sub>4</sub> is reduced and organic soils may even become a CH<sub>4</sub> sink, once methanotrophs dominate the  
 567 CH<sub>4</sub> cycle. As natural CH<sub>4</sub> emissions are not included in the inventory, this emission reduction is not considered  
 568 when natural un-drained organic soils are being drained. However, for completeness any remaining CH<sub>4</sub>  
 569 emission from the land surface of drained organic soils needs to be included in inventories.

570 Ditch networks provide a further source of CH<sub>4</sub> emissions from drained organic soils. This occurs due to a  
 571 combination of lateral CH<sub>4</sub> transfer from the organic soil matrix, and in-situ CH<sub>4</sub> production within the ditches  
 572 themselves (e.g., Roulet and Moore, 1995; Van den Pol 1999c; Van Dasselaar *et al.*, 1999a; Sundh *et al.*, 2000;  
 573 Minkinen and Laine, 2006; Teh *et al.*, 2011; Vermaat *et al.*, 2011). These emissions may approach, or even  
 574 exceed, the CH<sub>4</sub> flux from an undrained organic soil when averaged over the land surface (Roulet and Moore,  
 575 1995; Schrier-Uijl *et al.*, 2011). Emission/removal factors for ditch CH<sub>4</sub> emissions were compiled from available  
 576 published literature (See Annex 2A.1). We present only general factors for ditches because of limited data.  
 577 Effects of ditch maintenance, deepening etc. may be addressed at higher Tiers.

## 578 CHOICE OF METHOD

### 579 Tier 1

580 CH<sub>4</sub> emissions from the land surface are estimated using a simple emission factor approach (See Equation 2.6),  
 581 depending on climate and type of land-use. The default methodology considers boreal, temperate and tropical  
 582 climate zones and nutrient-rich and nutrient-poor organic soils. Different land-uses imply drainage to different  
 583 depths. The CH<sub>4</sub> emission factors depend on gas flux measurements, either from closed chambers or (for land-  
 584 surface emissions) from eddy covariance.

585 Ditch CH<sub>4</sub> emissions should be quantified for any area of drained organic soil where there are ditches or drainage  
 586 canals (note that CH<sub>4</sub> may also be emitted from ditches within re-wetted organic soils, where ditches remain  
 587 present, although at Tier 1 it is assumed that this flux equates to that from the remainder of the re-wetted site; see  
 588 Chapter 3). Estimation of ditch CH<sub>4</sub> emissions requires information on the land-use class and on the area of the  
 589 landscape occupied by the drainage ditch network,  $Frac_{ditch}$ .

590

591

592

**EQUATION 2.6**

**ANNUAL CH<sub>4</sub> EMISSION FROM DRAINED ORGANIC SOILS**

$$CH_{4\_organic} = \sum_{c,n,p} (A_{c,n,p} \cdot ((1 - Frac_{ditch}) \cdot EF_{CH_4\_land_{c,n}} + Frac_{ditch} \cdot EF_{CH_4\_ditch_{c,p}}))$$

593

594 Where:

595  $CH_{4\_organic}$  = Annual CH<sub>4</sub> loss from drained organic soils, kg CH<sub>4</sub> yr<sup>-1</sup>596  $A_{c,n,p}$  = Land area of drained organic soils in a land-use category in climate zone c, nutrient status n  
597 and soil type p, ha598  $EF_{CH_4\_land_{c,n}}$  = Emission factors for direct CH<sub>4</sub> emissions from drained organic soils, by climate zone c  
599 and nutrient status n, kg CH<sub>4</sub> ha<sup>-1</sup> yr<sup>-1</sup>600  $EF_{CH_4\_ditch_{c,p}}$  = Emission factors for CH<sub>4</sub> emissions from drainage ditches, by climate zone c and soil type  
601 p, kg CH<sub>4</sub> ha<sup>-1</sup> yr<sup>-1</sup>602  $Frac_{ditch}$  = Fraction of the total area of drained organic soil which is occupied by ditches. The ditch area  
603 may be calculated as the width of the ditches (from bank to bank) multiplied by their total length.

### 604 Tier 2

605 The Tier 2 approach for CH<sub>4</sub> emissions from drained organic soils incorporates country-specific information in  
 606 Equation 2.6 to estimate the emissions. Tier 2 uses the same procedural steps for calculations as provided for  
 607 Tier 1. Under Tier 2, the emission factors for CH<sub>4</sub> from the surface of drained organic soils can be further  
 608 differentiated by drainage depth, land-use subcategories or vegetation type (such as presence or absence of plant

609 species that act as transporters of CH<sub>4</sub> from the soil to the atmosphere). Guidance for further stratification  
610 follows the principles given in Section 2.2.1.1 of this chapter.

611 Tier 2 approaches for CH<sub>4</sub> emissions from drainage ditches generally follow the Tier 1 approach described above,  
612 with country-specific measurements or estimates of annual mean ditch CH<sub>4</sub> emissions, and national or regional  
613 estimates of fractional ditch area that reflect local drainage practices. The land-use sub-categories in Table 2.4  
614 may be expanded or sub-divided where appropriate to reflect the range of observed land-use on drained organic  
615 soils.

### 616 **Tier 3**

617 Tier 3 methods for estimating CH<sub>4</sub> emissions from drained organic soils involve a comprehensive understanding  
618 and representation of the dynamics of CH<sub>4</sub> emissions and removals on managed peatlands and organic soils,  
619 including the effect of site characteristics, peat/soil type, peat degradation and depth, land-use intensity, drainage  
620 depth, management systems, and the level and kinds of fresh organic matter inputs. Also emission spikes may  
621 occur, for example during spring thaw or strong rains or when debris from ditch dredging is deposited on  
622 adjacent land.

623 For CH<sub>4</sub> emissions from drainage ditches, development of a Tier 3 approach could take account of the influence  
624 of land-management activities (e.g., organic matter additions to agricultural land) on substrate supply for  
625 methane production in ditches, of possible short-term pulses of ditch CH<sub>4</sub> emissions associated with land-use  
626 change, and of the legacy effects of past land-use (e.g. nutrient-enriched soils). Information on drainage ditch  
627 characteristics and maintenance may be used to refine ditch CH<sub>4</sub> emissions estimates, for example taking account  
628 of the potential effects of plant or algal growth within ditches; presence of subsurface drainage in Croplands and  
629 Grasslands; water flow rates, transport length of water and oxygen status; ditch maintenance activities, and the  
630 deposition of organic material removed from ditches onto adjacent land areas.

## 631 **CHOICE OF EMISSION FACTORS**

### 632 **Tier 1**

633 Default emission factors for the Tier 1 method are provided in Table 2.3 for EF<sub>CH<sub>4</sub>\_land</sub> and Table 2.4 for  
634 EF<sub>CH<sub>4</sub>\_ditch</sub>.

635 At present, literature data are sufficient to provide Tier 1 default values of EF<sub>CH<sub>4</sub>\_ditch</sub> for each of the four major  
636 land-use classes on organic soils (Forest Land, Grassland, Cropland and Wetlands used for peat extraction) in  
637 boreal and temperate regions (Table 2.4). For Cropland, because no data are currently available, Tier 1 default  
638 values for deep-drained Grassland may be applied. For tropical organic soils, few data on ditch CH<sub>4</sub> emissions  
639 are currently available, and a single Tier 1 EF is therefore provided for all drained land-use classes.

### 640 **Tier 2**

641 Tier 2 emission factors EF<sub>CH<sub>4</sub>\_land</sub> may be based on country- or region-specific emission factors for CH<sub>4</sub>  
642 emissions from the surface of drained organic soils. These allow a further stratification of land-use categories by  
643 drainage class, nutrient status or vegetation characteristics.

644 Methane emissions from drainage ditches will vary according to peat/soil type, land-use type, drainage intensity,  
645 and (for agriculturally managed areas) land-use intensity. For example labile organic matter and nutrient inputs  
646 from terrestrial areas are likely to increase CH<sub>4</sub> production in ditches (Schrier-Uijl *et al.*, 2011). The Tier 1  
647 emission factors EF<sub>CH<sub>4</sub>\_ditch</sub> provided are based on measurements from ditches located within the organic layer.  
648 Subsurface drainage systems may represent additional sources of CH<sub>4</sub> emissions in Cropland and Grassland, and  
649 could be incorporated in the approach provided that appropriate measurement data are available. Countries are  
650 encouraged to obtain new measurement data for significant land-use classes to enhance the current dataset, and  
651 to develop country-specific Tier 2 emission factors. Sharing of data between countries may be appropriate where  
652 environmental conditions and practices are similar.

### 653 **Tier 3**

654 A Tier 3 approach for CH<sub>4</sub> emissions from drained organic soils might include further details and processes or  
655 capture the seasonal dynamics of CH<sub>4</sub> emissions as additional element of stratification or by dynamic modelling.

656 A Tier 3 approach for CH<sub>4</sub> emissions from drainage ditches might include the use of more detailed data to  
657 develop and apply process models that describe CH<sub>4</sub> emissions as a function of drainage ditch characteristics and  
658 maintenance, for example taking account of the potential effects of plant or algal growth within ditches; water  
659 flow rates, transport length of water and oxygen status; ditch maintenance activities, and the deposition of  
660 organic material removed from ditches onto adjacent land areas.

---

Final Draft

661 A Tier 3 approach to estimating ditch CH<sub>4</sub> emissions could take account of the temporal variability of  
662 hydrological conditions, labile substrate and nutrient supply, and controls on the composition of in-ditch-  
663 vegetation that might enhance or reduce emission rates.

664 Emissions from stockpiles of drying peat are uncertain and stockpiles may release or consume CH<sub>4</sub> at different  
665 rates than the excavation field, but data are not at present sufficient to provide guidance. Methods for estimating  
666 this flux may be developed for Tier 3 approaches

**TABLE 2.3**  
**TIER 1 CH<sub>4</sub> EMISSION/REMOVAL FACTORS FOR DRAINED ORGANIC SOILS (EF<sub>CH<sub>4</sub> LAND</sub>) IN ALL LAND-USE CATEGORIES**

Land-use category		Climate / Vegetation zones	Emission Factor*** (kg CH <sub>4</sub> ha <sup>-1</sup> yr <sup>-1</sup> )	95% Confidence Interval (centred on mean)		No. of Sites	Citations/Comment
Forest Land, drained	Nutrient-poor	Boreal	7.0	2.9	11	47	Komulainen <i>et al.</i> 1998 ; Lohila <i>et al.</i> , 2011; Maljanen <i>et al.</i> , 2006a ; Martikainen <i>et al.</i> , 1992 1993, 1995b; Minkkinen and Laine, 2006 ; Minkkinen <i>et al.</i> , 2006a, 2007a; Nykänen <i>et al.</i> , 1998 ; Ojanen <i>et al.</i> , 2010, 2013
	Nutrient-rich	Boreal	2.0	-1.6	5.5	83	Komulainen <i>et al.</i> , 1998; Laine, <i>et al.</i> , 1996; Mäkiranta <i>et al.</i> , 2007; Maljanen <i>et al.</i> , 2001, 2003b, 2006a ; Martikainen <i>et al.</i> , 1992, 1995b; Minkkinen and Laine, 2006; Minkkinen <i>et al.</i> , 2007a; Nykänen <i>et al.</i> , 1998; Ojanen <i>et al.</i> , 2010, 2013
Forest Land, drained		Temperate	2.5	-0.60	5.7	13	Glenn <i>et al.</i> , 1993; Moore and Knowles, 1990; Sikström <i>et al.</i> , 2009; Von Arnold <i>et al.</i> , 2005a, b; Weslien <i>et al.</i> , 2009; Yamulki <i>et al.</i> , 2013
Forest Land and cleared Forest Land (shrubland****), drained		Tropical/ Subtropical	4.9	2.6**			Jauhainen <i>et al.</i> , 2008; Hirano <i>et al.</i> , 2009; Furukawa <i>et al.</i> , 2005
Forest plantations, drained *****			2.7	3.6**			Basuki <i>et al.</i> , 2012 Jauhainen <i>et al.</i> , 2012c
Grassland, drained		Boreal	1.4	-1.6	4.5	12	Grønlund <i>et al.</i> , 2006; Guðmundsson and Óskarsson 2008; Hyvönen <i>et al.</i> , 2009; Maljanen <i>et al.</i> , 2001, 2003b, 2004; Nykänen <i>et al.</i> , 1995; Regina <i>et al.</i> , 2007
Grassland, drained, nutrient-poor		Temperate	1.8	0.72	2.9	9	Drösler <i>et al.</i> , 2013; Kasimir-Klemmedtsson <i>et al.</i> , 2009; Van Den Bos, 2003
Grassland, deep drained, nutrient- rich		Temperate	16	2.4	29	44	Augustin <i>et al.</i> , 1996; Best & Jacobs, 1997; Drösler <i>et al.</i> , 2013; Flessa <i>et al.</i> 1997, 1998 ; Jacobs <i>et al.</i> 2003; Kroon <i>et al.</i> 2010; Langeveld <i>et al.</i> , 1997; Meyer <i>et al.</i> , 2001; Nykanen <i>et al.</i> , 1995; Petersen <i>et al.</i> , 2012; Schrier-Uijl, 2010a,b; Teh <i>et al.</i> , 2011; Van Den Bos, 2003; Van den Pol-Van Dassel <i>et al.</i> , 1997; Wild <i>et al.</i> , 2001
Grassland, shallow		Temperate	39	-2.9	81	16	Augustin, 2003; Drösler <i>et al.</i> , 2013; Jacobs <i>et al.</i> ,

Final Draft

drained, nutrient-rich						2003, Van den Pol-Van Dasselaar <i>et al.</i> , 1997 667
Grassland	Tropical/ Subtropical	7.0	6.7*			Same emission factor as tropical Cropland
Cropland, drained	Boreal & Temperate	0	-2.8	2.8	38	Augustin, 2003; Augustin <i>et al.</i> , 1998; Drösler <i>et al.</i> , 2013; Elsgaard <i>et al.</i> , 2012; Flessa <i>et al.</i> , 1998; Kasimir-Klemedtsson <i>et al.</i> , 2009; Maljanen <i>et al.</i> , 2003a,b, 2004, 2007a; Petersen <i>et al.</i> , 2012; Regina <i>et al.</i> , 2007; Taft <i>et al.</i> , 2013
Cropland	Tropical/ Subtropical	7.0	6.7*			Furukawa <i>et al.</i> , 2005; Hirano <i>et al.</i> , 2009
Rice*****		143	80*			Furukawa <i>et al.</i> , 2005; Hadi <i>et al.</i> , 2001; Inubushi <i>et al.</i> , 2003
Plantation: oil palm		0	0			Melling <i>et al.</i> , 2005b
Plantation: sago palm		26	19*			Watanabe <i>et al.</i> , 2009; Melling <i>et al.</i> , 2005b; Inubushi <i>et al.</i> , 1998
Peat Extraction	Boreal & Temperate	6.1	1.6	11	15	Hyvönen <i>et al.</i> , 2009; Nykänen <i>et al.</i> , 1996; Strack and Zuback, 2013; Sundh <i>et al.</i> , 2000; Tuittila <i>et al.</i> , 2000; Waddington and Day, 2007
Settlements	All climate zones	There is no fixed default emission/removal factor for Settlements. It is <i>good practice</i> to take the default emission/removal factor in Table 2.1 of the land-use category that is closest to the national conditions of drained organic soils under Settlements. Information about national conditions could include drainage level, vegetation cover, or other management activities. For example, drained organic soils in urban green areas, parks or gardens could use the default Tier 1 emission/removal factor for Grassland, deep-drained in Table 2.1.				
Other Land	All climate zones	<i>Other Land Remaining Other Land: 0</i> <i>Land Converted to Other Land: Maintain emission factor of previous land-use category</i>				
<p>**Standard error</p> <p>*** Mean</p> <p>**** Shrubland refers to any type of land sparsely or fully covered with shrubs or trees, which may or may not fulfil the national forest definition. It extends to degraded lands, which cannot be clearly classified as forest or non-forest.</p> <p>***** Number derived solely from Acacia plantation data.</p> <p>***** The default value applies to countries without data about flooding regime for rice on organic soils. Countries with data about flooding regime for rice on organic soil may continue to use the methodologies and emission factors provided in the 2006 IPCC Guidelines.</p>						

668 Plantations can be defined as Forest Land or Cropland, according to national definitions. Plantations that  
669 produce food should be classified as Cropland; plantations producing wood and fibre should be reported as  
670 Forest Land. It is *good practice* to report plantations in the appropriate national land-use category according to  
671 the national forest definition.

## 672 CHOICE OF ACTIVITY DATA

### 673 Tier 1

674 It is *good practice* to use the same activity data for estimating CO<sub>2</sub>, N<sub>2</sub>O and CH<sub>4</sub> emissions from drained  
675 organic soils. Information on obtaining these data is provided in Section 2.2.1 above. For countries in boreal and  
676 temperate regions using the Tier 1 method, if the available information does not allow stratification by nutrient  
677 status of organic soils, countries may rely on guidance given in Section 2.2.1.1.

678 Activity data required to estimate CH<sub>4</sub> emissions from drainage ditches at Tier 1 consist of areas of managed  
679 organic soils disaggregated by land-use category (Forest Land, Grassland, Cropland, Wetlands used for peat  
680 extraction) as shown in Table 2.4. Fractional ditch areas recorded in published studies are given for individual  
681 sites in Table A2.1, and these data have been used to provide indicative Frac<sub>ditch</sub> values by land-use class in Table  
682 2.4. However it should be noted that these proportions are likely to vary between countries and it is therefore  
683 *good practice* to derive country-specific activity data on fractional ditch areas wherever possible, to reflect local  
684 land-use practices. This fractional ditch area may depend on the topographic situation and organic soil properties  
685 rather than on land-use alone. Fractional ditch area can be calculated from spatially explicit information about  
686 ditch and canal networks. From these the length and width of ditches can be derived, or alternatively, ditch  
687 spacing and ditch width on organic soils, which gives the ditch area on organic soils. This geometrical  
688 information is converted to fractional ditch area by dividing the ditch area on organic soils through the area of  
689 drained organic soils.

### 690 Tier 2 and 3

691 Activity data required for higher Tier methods are likely to include more detailed information on land-use, in  
692 particular land-use intensity within Grassland and Cropland classes. Further stratification may be necessary for  
693 other classes if sufficient data become available to estimate emission factors, e.g., for cleared peat swamp forest,  
694 oil palm or pulpwood plantation in tropical peat areas.

695 Activity data for higher Tier methods may be spatially explicit and consist of areas of organic soils managed for  
696 different forest types, peat extraction, production systems, horticulture and plantations, disaggregated according  
697 to nutrient status of the organic soil if relevant. More sophisticated estimation methodologies will require the  
698 determination of areas in different phases of land-uses with longer term rhythms such as age-classes in Forest  
699 Land or in a peat extraction operation, where on abandoned areas drainage or the effects of former peat  
700 extraction are still present. Land-use intensity, particularly fertilizer and organic matter addition, may be used to  
701 refine CH<sub>4</sub> emission estimates for Grassland and Cropland, as emissions are likely to change under more  
702 intensive management systems.

703 To estimate CH<sub>4</sub> emissions from drainage ditches, additional activity data are required on fractional ditch area  
704 within each land use category. Country-specific values of fractional ditch areas are used to reflect drainage  
705 methodologies such as typical ditch spacing, depth, width and length, maintenance (such as vegetation clearance)  
706 and land-use practices. Fractional ditch area can be stratified by type of organic soil or topographic situation,  
707 peat/soil properties and land-use.

708 Activity data for CH<sub>4</sub> emissions from drainage ditches could incorporate additional information on water table  
709 level and variability (such as seasonal water management regime), flow rates, in-ditch vegetation and land-use  
710 factors affecting substrate supply for methanogenesis, such as livestock density and fertilizer application in  
711 intensive Grasslands and Croplands. Incorporating seasonal and short-term controls on emissions would require  
712 additional activity data on the nature and timing of agricultural activities (such as organic matter additions) and  
713 on hydrological parameters.

## 714 CALCULATION STEPS FOR TIER 1

715 The steps for estimating the CH<sub>4</sub> emissions from drained inland organic soils are as follows:

716 **Step 1:** Determine areas with drained inland organic soils under each land-use category for lands remaining in a  
717 land-use category, disaggregated by climate domain and other appropriate factors as outlined above and  
718 consistently with on-site CO<sub>2</sub> emissions estimates from drained organic soils. Where needed for Tier 1 emission  
719 factors, land areas are further stratified by nutrient-rich and nutrient-poor organic soils. Temperate nutrient-rich  
720 Grasslands are further stratified into shallow-drained and deep-drained classes.

721 **Step 2:** Assign the appropriate value for the fraction of areas covered by ditches using national statistics. If  
722 statistics are not available, values given in Table 2.4 provide appropriate defaults.

---

Final Draft

723 **Step 3:** Assign the appropriate emission factor values ( $EF_{CH_4\_land}$  and  $EF_{CH_4\_ditch}$ ) from Tables 2.3 and 2.4,  
724 respectively.

725 **Step 4:** Multiply each area with the appropriate emission factor by using Equation 2.6 and sum across land use  
726 categories.

727

728

<b>Climate zone</b>	<b>Land-use</b>	<b>EF<sub>CH<sub>4</sub> ditch</sub> (kg CH<sub>4</sub> ha<sup>-1</sup> yr<sup>-1</sup>)</b>	<b>Uncertainty range (kg CH<sub>4</sub> ha<sup>-1</sup> yr<sup>-1</sup>)</b>	<b>No. of sites</b>	<b>Frac<sub>ditch</sub> (indicative values<sup>d</sup>)</b>	<b>References</b>
Boreal /temperate	Drained forest, Drained wetland <sup>a</sup>	217	41 – 393	11	0.025	Cooper & Evans, 2013; Glagolev <i>et al.</i> , 2008; Minkkinnen & Laine, 2006 (two study areas); Roulet & Moore, 1995 (three study areas); Sirin <i>et al.</i> , 2012 (3 study areas); von Arnold <i>et al.</i> , 2005b.
	Shallow-drained Grassland	527	285 – 769	5	0.05	Best & Jacobs, 1997; Hendriks <i>et al.</i> , 2007, 2010; Van den Pol-Van Dasselaar <i>et al.</i> , 1999a; Vermaat <i>et al.</i> , 2011; McNamara, 2013
	Deep-drained Grassland Cropland <sup>b</sup>	1165	335 – 1995	6	0.05	Best & Jacobs, 1997; Chistotin <i>et al.</i> , 2006 ;; Schrier-Uijl <i>et al.</i> , 2010, 2011; Sirin <i>et al.</i> , 2012; Teh <i>et al.</i> , 2011; Vermaat <i>et al.</i> , 2011.
	Peat Extraction	542	102 – 981	6	0.05	Chistotin <i>et al.</i> , 2006; Nykänen <i>et al.</i> , 1996; Sirin <i>et al.</i> , 2012; Sundh <i>et al.</i> , 2000; Waddington & Day, 2007; Hyvönen <i>et al.</i> , 2013
Tropical	All land-use involving drainage	2259	599 – 3919 <sup>c</sup>	2	0.02	Jauhianen & Silvennoinen, 2012 (drained and abandoned, and pulpwood plantation)
Values shown in parentheses represent 95% confidence intervals unless otherwise stated						
<sup>a</sup> Ditch CH <sub>4</sub> emissions from wetlands subject to drainage but no other land-use modification are assumed to be equivalent to those from organic soils drained for forestry.						
<sup>b</sup> Ditch CH <sub>4</sub> emissions from Cropland are assumed to be the same as those from high-intensity Grassland, for which more data exist.						
<sup>c</sup> Due to limited data for CH <sub>4</sub> emissions from tropical drainage channels, the range of measurements is shown, rather than 95% confidence intervals.						
<sup>d</sup> Indicative values for Frac <sub>ditch</sub> within each class are derived from the mean of studies reporting CH <sub>4</sub> emission values for this class. Note that studies from the Netherlands were not included in this calculation, because they are characterised by much higher fractional ditch areas (0.1 to 0.25) that are not typical of drained organic soils in other countries.						

729

## 730 UNCERTAINTY ASSESSMENT

731 The principal sources of uncertainty for CH<sub>4</sub> emissions from drained organic soils are activity data, including  
 732 associated information on the fraction of drained areas covered by ditches, and emission factors. Uncertainty  
 733 ranges are provided in Tables 2.3 for values of EF<sub>CH<sub>4</sub> land</sub> and Table 2.4 for values of EF<sub>CH<sub>4</sub> ditch</sub> for each organic  
 734 soil/land-use category. Uncertainty ranges in Table 2.3 are expressed as 95% confidence intervals or as standard  
 735 errors, depending on the number of studies available. The major source of uncertainty in these values is simply  
 736 the small number of studies on which many Tier 1 estimates are based, and the high degree of heterogeneity in  
 737 measured fluxes between different studies undertaken within some classes. Confidence intervals (95%) have  
 738 been calculated for all classes other than the drained tropical organic soil class, for which only one study  
 739 (Jauhianen and Silvennoinen, 2012) is available, which provides estimates of ditch CH<sub>4</sub> emissions from areas of

Final Draft

740 drained, deforested and abandoned organic soils, and pulpwood plantation. For the drained tropical organic soils  
741 category, the uncertainty range is provided by the lower (abandoned) and higher (pulpwood plantation) emission  
742 values recorded.

743 The final calculation of  $CH_4_{organic}$  is also sensitive to uncertainties in the activity data, and in particular to data  
744 used to estimate the proportion of the land area which is occupied by drainage ditches,  $Frac_{ditch}$ . Many countries  
745 lack such data and although activity data should be country-specific, even for Tier 1, indicative values from  
746 Table 2A.1 can be used at the discretion of the inventory compiler. Uncertainty assessments should therefore  
747 also take account of this source of uncertainty in calculating total  $CH_4$  emissions from drained organic soils.

### 748 2.2.2.2 N<sub>2</sub>O EMISSIONS FROM DRAINED ORGANIC SOILS

749 N<sub>2</sub>O emissions from soils are produced by the microbiological processes of nitrification and denitrification (to  
750 N<sub>2</sub>O or N<sub>2</sub>) (Davidson 1991; Firestone and Davidson, 1989). These processes are controlled by several factors,  
751 including water-filled pore space (Aulakh and Sigh, 1997; Davidson 1991; Dobbie *et al.* 1999; Ruser *et al.*,  
752 2001), temperature (Keeney *et al.*, 1979; Kroon *et al.*, 2010), and concentration of mineral nitrogen (Bremner  
753 1997; Firestone and Davidson, 1989; Ryden and Lund, 1980).

754 Drained organic soils emit significant amounts of N<sub>2</sub>O, whereas emissions from wet organic soils are close to  
755 zero (Kasimir-Klemedtsson *et al.*, 1997; Flessa *et al.*, 1998; Couwenberg *et al.*, 2011). A main reason for  
756 increased N<sub>2</sub>O emissions is nitrogen mineralization associated with organic matter decomposition in drained  
757 organic soils (Höper, 2002). Emissions from this N mineralization will be dealt with here. Other sources of  
758 anthropogenic N in organic soils include nitrogen fertilizer, application of crop residues, organic amendments,  
759 and use of N fixing species. These emissions from other N sources are dealt with in Chapter 11 of Volume 4 of  
760 the 2006 IPCC Guidelines and in all earlier guidance.

761 Most of the published data on N<sub>2</sub>O fluxes from managed organic soils refer to boreal and temperate ecosystems  
762 and these data served as the basis for the emission factors in the 2006 IPCC Guidelines. With new studies  
763 published since 2005, there are enough data to derive separate N<sub>2</sub>O emission factors for Forest Land, Cropland,  
764 Grassland, and peatlands under peat extraction in boreal and temperate zones and these new values replace the  
765 values Table 7.6 in Volume 4, Chapter 7 of the 2006 IPCC Guidelines.

766 There are still limited data available for tropical organic soils. However, the studies that have been published  
767 over the past decade provide enough data to develop Tier 1 emissions factors for the first time.

## 768 CHOICE OF METHOD

### 769 Tier 1

770 This section presents the equation for estimating direct emissions of N<sub>2</sub>O due to drainage of organic soils. The  
771 revisions presented here, as shown in Equation 2.7, are applicable to Equation 11.1 presented in Chapter 11,  
772 Volume 4 of the 2006 IPCC Guidelines. This Equation is used to estimate N<sub>2</sub>O for specific land-use categories,  
773 but there are not enough data available for developing coefficients to modify EFs by condition-specific variables  
774 (e.g., variations of drainage depths). The Equations 11.1 and 11.2 have been modified to include variables for the  
775 boreal climate zone as well by adding terms  $F_{OS,CG,Bor,NR}$ ,  $F_{OS,CG,Bor,NP}$ ,  $F_{OS,F,Bor,NR}$ , and  $F_{OS,F,Bor,NP}$  (the  
776 subscripts CG, F, Bor, NR and NP refer to Cropland and Grassland, Forest Land, Boreal, Nutrient-Rich, and  
777 Nutrient-Poor, respectively) and their respective emissions factors.

778 Direct N<sub>2</sub>O emissions from managed soils are estimated using Equation 11.1 in Chapter 11, Volume 4 of the  
779 2006 IPCC Guidelines. This Equation has three segments: one for emissions associated with N inputs, one for  
780 organic soils, and one for urine and dung inputs during grazing. In this section, updates are provided for the  
781 second segment focusing on organic soils as follows:

782

783

784

$$\begin{array}{c}
 \text{EQUATION 2.7} \\
 \text{DIRECT N}_2\text{O EMISSIONS FROM MANAGED ORGANIC SOILS} \\
 N_2O - N_{OS} = \left[ \begin{array}{l}
 (F_{OS,CG,Bor} \cdot EF_{2CG,Bor}) + (F_{OS,CG,Temp} \cdot EF_{2CG,Temp}) + (F_{OS,CG,Trop} \cdot EF_{2CG,Trop}) + \\
 (F_{OS,F,Bor,NR} \cdot EF_{2F,Bor,NR}) + (F_{OS,F,Temp,NR} \cdot EF_{2F,Temp,NR}) + \\
 (F_{OS,F,Bor,NP} \cdot EF_{2F,Bor,NP}) + (F_{OS,F,Temp,NP} \cdot EF_{2F,Temp,NP}) + (F_{OS,F,Trop} \cdot EF_{2F,Trop})
 \end{array} \right]
 \end{array}$$

785

786 Where:

787  $N_2O-N_{OS}$  = Annual direct  $N_2O-N$  emissions from managed organic soils,  $kg N_2O-N yr^{-1}$   
 788  $F_{OS}$  = Annual area of managed/drained organic soils, ha (Note: the subscripts CG, F, Temp, Trop,  
 789 NR and NP refer to Cropland and Grassland, Forest Land, Temperate, Tropical, Nutrient Rich, and  
 790 Nutrient Poor, respectively)  
 791  $EF_2$  = Emission factor for  $N_2O$  emissions from drained/managed organic soils,  $kg N_2O-N ha^{-1} yr^{-1}$ ;  
 792 (equivalent to Table 11.1, Chapter 11, Volume 4, of the *2006 IPCC Guidelines* but using updated  
 793 emission factor values provided in Table 2.5 below; note: the subscripts CG, F, Temp, Trop, NR and  
 794 NP refer to Cropland and Grassland, Forest Land, Temperate, Tropical, Nutrient Rich, and Nutrient  
 795 Poor, respectively.).

## 796 **Tier 2**

797 Tier 2 estimates are to be based on the Tier 1 Equation 2.7, but use country or region-specific emission factors.  
 798 These can be further stratified by drainage class, nutrient status of organic soils or other criteria used for  
 799 stratifying organic soils for direct  $N_2O$  emissions. The corresponding emission factors are country or region-  
 800 specific and take into account the land management systems. Tier 2 emission factors can follow the Tier 1  
 801 assumption that N mineralization from the degrading organic matter exceeds the amount of N input so that the  
 802 measured  $N_2O$  emissions are entirely attributed to the drained organic soil.

## 803 **Tier 3**

804 Tier 3 approaches can attribute  $N_2O$  emissions from drained organic soils separately to the mineralization of peat  
 805 or organic matter versus N input by fertilizer, crop residues and organic amendments. Attribution could rely on  
 806 the fraction of  $N_2O$  released by  $N_2O$  emission peaks after N fertilization, or by subtracting a fertilizer EF from  
 807 total  $N_2O$  emissions. Nitrogen mineralization from the drained organic soil can be estimated by the  $CO_2-C$   
 808 emission from the drained organic soil and the C/N ratio of the topsoil and this value could be used to predict  
 809  $N_2O$  emissions.

810 Tier 3 methods are based on modelling or measurement approaches. Models can simulate the relationship  
 811 between the soil and environmental variables that control the variation in  $N_2O$  emissions and the size of those  
 812 emissions (Stehfest & Bouwman, 2006; Kroon *et al.*, 2010; Dechow & Freibauer, 2011). These models can be  
 813 used at larger scales where measurements are impractical. Models should only be used after validation against  
 814 representative measurements that capture the variability of land-use, management practices and climate present  
 815 in the inventory (IPCC, 2010).

## 816 **CHOICE OF EMISSION FACTORS**

### 817 **Tier 1**

#### 818 ***Emission factors for boreal and temperate organic soils***

819 The *2006 IPCC Guidelines* provided emission factors that were partly disaggregated for land-use types or  
 820 climatic zones (Table 11.1), Chapter 11, Volume 4). An increased availability of scientific data allows for an  
 821 improved choice of default emission factors (Table 2.5). Nutrient poor and rich organic soils drained for forestry  
 822 have different  $N_2O$  emissions. Croplands and Grasslands are established on nutrient-rich organic soil or are  
 823 amended for better nutrient availability, and are considered here as rich. Peat extraction occurs both on nutrient-  
 824 poor (bogs) and nutrient-rich (fens) peatlands. In all cases the residual bottom organic layers consist of  
 825 minerogenous but recalcitrant nutrient-rich peat. There is not enough data available to disaggregate for the peat  
 826 types in peat extraction areas.

827

Final Draft

828

TABLE 2.5 TIER 1 DIRECT N <sub>2</sub> O EMISSION/REMOVAL FACTORS FOR DRAINED ORGANIC SOILS IN ALL LAND-USE CATEGORIES							
Land-use Category		Climate / Vegetation zone	Emission factor (kg N <sub>2</sub> O-N ha <sup>-1</sup> yr <sup>-1</sup> )	95 % confidence interval		No. of Sites	Citations/Comment
Forest Land, drained	Nutrient- poor	Boreal	0.22	0.15	0.28	43	Lohila <i>et al.</i> , 2011; Maljanen <i>et al.</i> , 2006a; Martikainen <i>et al.</i> , 1993, 1995a; Ojanen <i>et al.</i> , 2010, 2013; Regina <i>et al.</i> , 1996
	Nutrient- rich	Boreal	3.2	1.9	4.5	75	Ernfors <i>et al.</i> , 2011; Mäkiranta <i>et al.</i> , 2007; Maljanen <i>et al.</i> , 2001, 2003a, 2006a, 2010a; Martikainen <i>et al.</i> , 1993, 1995a; Ojanen <i>et al.</i> , 2010, 2013 ; Pihlatie <i>et al.</i> , 2004; Regina <i>et al.</i> , 1998; Saari, <i>et al.</i> , 2009
Forest Land, drained		Temperate	2.8	-0.57	6.1	13	Sikström <i>et al.</i> , 2009; Von Arnold <i>et al.</i> , 2005a, b; Weslien <i>et al.</i> , 2009; Yamulki <i>et al.</i> , 2013
Forest Land and cleared Forest Land (shrubland****), drained		Tropical/ Subtropical	2.4	1.1*			Furukawa <i>et al.</i> , 2005; Jauhiainen <i>et al.</i> , 2012b; Takakai <i>et al.</i> , 2006
Grassland, drained		Boreal	9.5	4.6	14	16	Grönlund <i>et al.</i> , 2006; Hyvönen <i>et al.</i> , 2009; Jaakkola, 1985; Maljanen <i>et al.</i> , 2001, 2003a, 2004, 2009; Nykänen <i>et al.</i> , 1995; Regina <i>et al.</i> , 1996, 2004
Grassland, drained, nutrient-poor		Temperate	4.3	1.9	6.8	7	Drösler <i>et al.</i> , 2013; Kasimir-Klemedtsson <i>et al.</i> , 2009
Grassland, deep drained, nutrient-rich		Temperate	8.2	4.9	11	47	Augustin and Merbach, 1998; Augustin <i>et al.</i> , 1996, 1998; Drösler <i>et al.</i> , 2013; Flessa <i>et al.</i> , 1997, 1998; Jacobs <i>et al.</i> , 2003; Kroon <i>et al.</i> , 2010; Langeveld <i>et al.</i> , 1997; Meyer <i>et al.</i> , 2001; Nykänen <i>et al.</i> , 1995; Petersen <i>et al.</i> , 2012; Teh <i>et al.</i> , 2011; van Beek <i>et al.</i> , 2010; Velthof <i>et al.</i> , 1996; Wild <i>et al.</i> , 2001
Grassland, shallow drained, nutrient-rich		Temperate	1.6	0.56	2.7	13	Drösler <i>et al.</i> , 2013; Jacobs <i>et al.</i> , 2003
Grassland		Tropical/ Subtropical	5.0	2.7*			Emission factor for tropical Cropland can be used
Cropland, drained		Boreal & Temperate	13	8.2	18	36	Augustin <i>et al.</i> , 1998; Drösler <i>et al.</i> , 2013; Elsgaard <i>et al.</i> , 2012; Flessa <i>et al.</i> , 1998; Kasimir-Klemedtsson <i>et al.</i> , 2009; Maljanen <i>et al.</i> , 2003a,b, 2004, 2007a; Petersen <i>et al.</i> , 2012; Regina <i>et al.</i> , 2004; Taft <i>et al.</i> , 2013
Cropland except rice		Tropical/	5.0	2.7*			Furukawa <i>et al.</i> , 2005; Jauhiainen <i>et al.</i> , 2012b; Takakai <i>et al.</i> , 2006

	Subtropical					
Rice		0.4	0.5*			Furukawa <i>et al.</i> , 2005; Hadi <i>et al.</i> , 2005; Inubushi <i>et al.</i> , 2003
Plantation: oil palm		1.2				Melling <i>et al.</i> , 2007b
Plantation: sago palm		3.3				Melling <i>et al.</i> , 2007b
Peatland Managed for Extraction	Boreal & Temperate	0.30	-0.03	0.64	4	Hyvönen <i>et al.</i> , 2009 ; Nykänen <i>et al.</i> , 1996 ; Regina <i>et al.</i> , 1996
Peatlands Managed for Extraction	Tropical/ Subtropical	3.6	0.2 to 5.0***			Emission factor from the 2006 IPCC Guidelines can be used.
Settlements	All climate zones	Emission factor for Cropland can be used*****				
Other Lands	All climate zones	<i>Other Land Remaining Other Land: 0</i> <i>Land Converted to Other Land: Maintain emission factor of previous land-use category</i>				
<p>** Standard error</p> <p>*** Mean</p> <p>**** Shrubland refers to any type of land sparsely or fully covered with shrubs or trees, which may or may not fulfil the national forest definition. I extends to degraded lands, which cannot be clearly classified as forest or non-forest.</p> <p>***** Lands in the settlements were mostly used for gardens.</p>						

---

Final Draft

829 Plantations can be defined as Forest Land or Cropland. The attribution to Cropland made in this table is not  
830 binding. It is *good practice* to report plantations in the appropriate national land-use category according to the  
831 national forest definition.

832 In the *2006 IPCC Guidelines*, emission factors were provided for  $EF_{2CG, Trop}$  and  $EF_{2F, Trop}$ , based on the  
833 expectation that net mineralization was twice as high in tropical soils compared to temperate soils. Research in  
834 tropical soils suggests that net mineralization is not a useful predictor of  $N_2O$  flux and that net nitrification or the  
835 nitrate portion of the inorganic-N pool are better predictors (Verchot *et al.*, 1999, 2006; Ishizuka *et al.*, 2005). It  
836 also needs to be highlighted that all measurements of  $N_2O$  emissions on tropical organic soils to date are from  
837 Southeast Asia and from a very limited number of studies. Nonetheless these EFs are to be used for all tropical  
838 ecosystems until better data become available.

### 839 **Tier 2**

840 Tier 2 emission factors may be based on country- or region-specific emission factors for  $N_2O$  emissions from the  
841 surface of drained organic soils. These allow a further stratification of land-use categories by drainage class,  
842 nutrient status or vegetation characteristics. Countries are encouraged to obtain new measurement data for  
843 significant land-use classes to enhance the current dataset, and to develop country-specific Tier 2 emission  
844 factors. Sharing of data between countries may be appropriate where environmental conditions and practices are  
845 similar.

### 846 **Tier 3**

847 Tier 3 emission factors or relations are based on country-specific emission data and models calibrated for  
848 management practices such as drainage intensity; crop, livestock or forest type; fertiliser or organic matter  
849 additions; peat extraction technology and the phases of peat extraction or other relevant factors for  $N_2O$   
850 emissions.

## 851 **CHOICE OF ACTIVITY DATA**

852 Activity data consist of areas of land remaining in a land-use category on organic soils stratified by major land-  
853 use types, management practices, and disturbance regimes. Total areas should be determined according to  
854 approaches laid out in Chapter 3 of the *2006 IPCC Guidelines* and should be consistent with those reported  
855 under other sections of the inventory. Stratification of land-use categories according to climate regions, based on  
856 default or country-specific classifications, can be accomplished with overlays of land-use on suitable climate and  
857 soil maps.

### 858 **Tier 1**

859 It is *good practice* to use activity data for  $N_2O$  emissions consistent with activity data for  $CO_2$  and  $CH_4$  emissions  
860 from soils. Guidance for activity data is given in the respective sections in this Chapter.

### 861 **Tier 2 and 3**

862 Activity data required for higher Tier methods are likely to include more detailed information on land-use, in  
863 particular land-use intensity within Grassland and Cropland classes. Further stratification may be necessary for  
864 other classes if sufficient data become available to estimate emission factors, e.g., for cleared peat swamp forest,  
865 oil palm or pulpwood plantations in tropical peat areas.

866 Activity data for higher Tier methods may be spatially explicit and consist of areas of organic soils managed for  
867 different forest types, peat extraction, cultivation systems, horticulture and plantations, disaggregated according  
868 to nutrient status of the organic soil if relevant, and annual peat production data. More sophisticated estimation  
869 methodologies will require the determination of areas in different phases of land-uses with longer term rhythms  
870 such as age-classes in Forest Land or in a peat extraction cycle, where on abandoned areas drainage or the effects  
871 of former peat extraction are still present.

## 872 **CALCULATION STEPS FOR TIER 1**

873 The steps for estimating  $N_2O$  emissions on drained inland organic soils are as follows:

874 **Step 1:** Determine areas with drained inland organic soils under each land-use category for lands remaining in a  
875 land-use category, disaggregated by climate domain and other appropriate factors as outlined above. Where  
876 needed for Tier 1 emission factors, land areas are further stratified by nutrient-rich and nutrient-poor organic  
877 soils. Temperate nutrient-rich Grassland is further stratified into shallow-drained and deep-drained classes.

878 **Step 2:** Assign the appropriate values for  $EF_2$  from Table 2.5 for each land-use category, climate domain,  
879 nutrient status and drainage class stratum.

880 **Step 3:** Multiply activity data by the emission factor for each land use category according to Equation 2.7.

Final Draft

## 881 **UNCERTAINTY ASSESSMENT**

882 Uncertainties in estimates of direct N<sub>2</sub>O emissions from drained organic soils are caused by uncertainties related  
883 to the emission factors (see Table 2.5 for uncertainty ranges), inter-annual variability associated with temperature  
884 and precipitation, activity data, lack of coverage of measurements, spatial aggregation, and lack of information  
885 on specific on-farm practices.

886 Additional uncertainty will be introduced in an inventory when emission factors are derived from measurements  
887 that are not representative of the variation of conditions in a country. Because of very high spatial variability of  
888 N<sub>2</sub>O emissions from soils, most estimates have large standard errors relative to the mean flux. In general, the  
889 uncertainty of activity data will be lower than that of the emission factors. Additionally, uncertainties may be  
890 caused by missing information on variation in drainage levels, and changing management practices in farming.  
891 Generally, it is difficult to obtain information on the actual drainage levels and possible emission reductions  
892 achieved as well as information on farming practices. For more detailed guidance on uncertainty assessment  
893 refer to Chapter 3, Volume 1 of the *2006 IPCC Guidelines*.

### 894 **2.2.2.3 CO<sub>2</sub> AND NON-CO<sub>2</sub> EMISSIONS FROM FIRES ON DRAINED** 895 **ORGANIC SOILS**

896 Fires can be a large and variable source of greenhouse gases and significantly affect other feedbacks within the  
897 climate system. When compared to combustion of above-ground vegetation, the emissions from both  
898 uncontrolled wildfires and managed (prescribed) fires in organic (peat) soils are high. On organic soils, fires  
899 comprise both surface fires that consume vegetation, litter and duff, and ground fires which burn into and below  
900 the surface consuming soil organic matter and deadwood mass as a fuel source. The latter are smouldering fires  
901 that may persist for long periods of time, burn repeatedly in response to changing soil moisture and surface  
902 hydrology, and penetrate to different depths. This section addresses the emissions arising from combustion of  
903 soil organic material.

904 In any ecosystem, fire activity is strongly influenced by several factors, namely weather/climate, fuel availability,  
905 drainage and ignition agents, including human activities (Johnson, 1992; Swetnam, 1993). In ecosystems with  
906 organic soils, conditions such as organic soil depth and density, soil moisture, vegetation composition and soil  
907 surface micro-topography (e.g., Benscoter and Wieder., 2003) along with fire characteristics, such as intensity,  
908 frequency and duration (Kasischke *et al.*, 1995), which are affected by fire management practices, influence the  
909 quantity of organic matter consumed and hence the emissions of greenhouse gases (Kuhry 1994; Kasischke *et al.*,  
910 1995; Kasischke and Bruhwiler, 2003).

911 *2006 IPCC Guidelines* covered emissions from burning of above-ground carbon stocks (biomass and dead  
912 organic matter) but did not cover the often substantial release of emissions from combustion of organic soils. It  
913 is *good practice* to report greenhouse gas emissions from fires on all managed lands with organic soils Including  
914 all fire related emissions both from natural fires as well as those that have a human-induced cause(e.g., soil  
915 drainage) even if the initiation of the fire is non-anthropogenic (e.g., lightning strike).

916 This Chapter updates the *2006 IPCC Guidelines* by:

- 917 • Providing default methodologies and emission factors for CO<sub>2</sub>, CH<sub>4</sub> and CO emissions from fires on organic  
918 soils
- 919 • Providing generic guidance for higher Tier methods to estimate these fluxes

920 Change in soil organic carbon following fire is the result of both CO<sub>2</sub> as well as non-CO<sub>2</sub> emissions (principally  
921 of CH<sub>4</sub> and CO). Emissions of both CO<sub>2</sub> and non-CO<sub>2</sub> greenhouse gases are addressed in the following sections.  
922 These deal specifically with below-ground biomass as opposed to vegetation and litter losses (the latter are  
923 included in the estimation of carbon stock changes in the *2006 IPCC Guidelines*).

## 924 **CHOICE OF METHOD**

925 CO<sub>2</sub> and non-CO<sub>2</sub> emissions from burning of drained organic soils can either be directly measured or estimated  
926 using data on area burnt along with the default values for mass of fuel consumed and emission factors provided  
927 in this chapter. Previous IPCC Guidelines have noted that emissions from wildfires on managed (and unmanaged)  
928 land can exhibit large inter-annual variations that may be driven by either natural causes (e.g., climate cycles,  
929 random variation in lightning ignitions), or indirect and direct human causes (e.g., prescribed burning, historical  
930 fire suppression and past forest harvest activities) or a combination of all three causes, the effects of which  
931 cannot be readily separated. This variability is also true for emissions from fires on organic soils which critically  
932 depend on the extent and depth of the organic soil, the fuel moisture, the water table depth, and hence the

Final Draft

933 thickness of the drained layer and the resulting depth of the consumed organics, all of which are affected by site  
 934 characteristics, weather, land management, fire type, and climate. At Tier 1, differentiation by land management  
 935 category and fire type is possible, but reporting at higher Tiers will enable a greater level of differentiation  
 936 between land-use, site characteristics and fire types.

937 The parameters required to calculate the CO<sub>2</sub> and non-CO<sub>2</sub> emissions from burning organic soils are: area burned,  
 938 mass of fuel available for consumption, combustion factor (this is also known as burning efficiency and can be  
 939 used to characterize smouldering vs. flaming fires) and emission factor. Compared with vegetation fires, the  
 940 uncertainties involved in estimating emissions from fires on organic soils are much higher because organic soils  
 941 can burn repeatedly and to different depths. Furthermore, the type and density of the soil organic material  
 942 combined with the combustion efficiency will determine the nature of the gases and other compounds emitted.

943 The mass of fuel that can potentially burn in a fire event on organic soils will be determined by measuring the  
 944 depth of burn, along with soil bulk density and carbon content; the former is strongly controlled by soil water  
 945 content (influenced by position of the water table or permafrost depth) while the latter variables are ideally  
 946 measured in the field. While default values can be used for Tier 1 reporting, for higher Tiers data on the depth of  
 947 burn and soil carbon density need to be determined. The combustion factor describes how much of the fuel mass  
 948 available is actually consumed during a fire event, i.e., converted into CO<sub>2</sub> or non-CO<sub>2</sub> gases. The emission  
 949 factor ( $G_{ef}$ ) determines the mass of CO<sub>2</sub> or non-CO<sub>2</sub> gas emitted per mass of fuel consumed by the fire (e.g., g  
 950 CO<sub>2</sub>/kg dry fuel). The total emissions of CO<sub>2</sub> or non-CO<sub>2</sub> gases are calculated from the product of area burnt and  
 951 the corresponding biomass loading, combustion factor, and emission factor.

**EQUATION 2.8**  
**ANNUAL CO<sub>2</sub>-C AND NON-CO<sub>2</sub> EMISSIONS FROM ORGANIC SOIL FIRE**

$$L_{fire} = A \cdot M_B \cdot C_f \cdot G_{ef} \cdot 10^{-3}$$

955 Where:

956  $L_{fire}$  = amount of CO<sub>2</sub> or non-CO<sub>2</sub> emissions, e.g., CH<sub>4</sub> from fire, tonnes

957  $A$  = total area burned annually, ha

958  $M_B$  = mass of fuel available for combustion, tonnes ha<sup>-1</sup> (i.e. mass of dry organic soil fuel) (default values  
 959 in Table 2.6)

960  $C_f$  = combustion factor, dimensionless

961  $G_{ef}$  = emission factor for each gas, g kg<sup>-1</sup> dry matter burnt (default values in Table 2.7)

962  
 963 Where data for  $M_B$  and  $C_f$  are not available, a default value for the amount of fuel actually burnt (the product of  
 964  $M_B$  and  $C_f$ ) can be used under Tier 1 methodology (Table 2.6). The value  $10^{-3}$  converts  $L_{fire}$  to tonnes.

965 The amount of fuel that can be burned is given by the area burned annually and the mass of fuel available in that  
 966 area.


967 Default values for the Tier 1 method or components of a Tier 2 method are provided in Tables 2.6 and 2.7. For  
 968 higher Tiers, data on the variation in the mass of fuel available (based on site or region-specific data, including  
 969 area of organic soil burnt, depth of organic soil, depth of burn and/or depth of water table/soil moisture content  
 970 values and soil bulk density) are incorporated.

971 Figure 2.1 presents a decision tree that guides the selection of the appropriate Tier level to report CO<sub>2</sub> and non-  
 972 CO<sub>2</sub> emissions from the burning of organic soils.

973

974  
975

**Figure 2.2** Generic decision tree for identification of the appropriate tier to estimate greenhouse gas emissions from fires on organic soils



976  
977  
978  
979

Note:

1: See Chapter 4, “Methodological Choice and Identification of Key Categories” (noting Section 4.1.2 on limited resources), Volume 1 of the 2006 IPCC Guidelines for discussion of *key categories* and use of decision trees.

---

Final Draft**980 Tier 1**

981 Countries may choose to report CO<sub>2</sub> emissions using the Tier 1 method if fires on organic soils are not a *key*  
982 *category*. This approach is based on highly aggregated data and default factors. It does, however, require primary  
983 data on area burned.

984 If burning in ecosystems with organic soils is a *key category*, countries are encouraged to report emissions by  
985 applying the highest tier possible, given national circumstances. For prescribed fires, country-specific data will  
986 be required to generate reliable estimates of emissions.

**987 Tiers 2 and 3**

988 The Tier 1 method is refined by incorporating more disaggregated area estimates (per organic soil and fire type  
989 sub-categories) and country-specific estimates of combustion and emission factors into Equation 1. Tier 2 uses  
990 the same procedural steps for calculations as provided for Tier 1. Potential improvements to the Tier 1 approach  
991 may include:

- 992 • Knowledge of the amount of soil organic matter consumed;
- 993 • The position of the soil water table relative to the surface;
- 994 • Improved information on land-use/management and their effects on organic soil condition, in particular  
995 hydrological status; and
- 996 • Improved data on area burnt, estimated using remotely sensed data of adequate spatial and temporal  
997 resolutions and verified according to a robust sampling design at suitable periodicity to take account of the  
998 monthly variations of area burnt. Estimates of the depth of burn in a representative number of locations.

999 Countries may further stratify the data on area burnt by depth of burn, organic soil condition (e.g., drained vs.  
1000 undrained, with further detail possible through characterisation of the intensity of drainage) and fire types  
1001 (wildfire vs. prescribed).

1002 It may also be possible to develop models with algorithms to generate regional scale maps of area burnt using  
1003 satellite data of multiple sources and of moderate spatial resolution. Model results should be validated, for  
1004 example, by using high spatial resolution data augmented by field observations, and refined based on the  
1005 validation results whenever possible. A sampling approach can be designed to generate estimates of area burnt.  
1006 This reporting method should provide estimates (fluxes) of the impact of burning on below-ground biomass,  
1007 particularly including the depth of burn, and if feasible the variation of depth within the area burned. Reporting  
1008 at higher Tiers should differentiate fires burning at different intensities (critical for Tier 3) and with different  
1009 proportions of smouldering vs. flaming combustion (i.e. different Modified Combustion Efficiency (MCE)  
1010 defined as  $\Delta\text{CO}_2/(\Delta\text{CO}_2 + \Delta\text{CO})$  which is an index of the relative proportion of smouldering vs. flaming  
1011 combustion). The development of robust methodologies to assess burn severity in organic soils would enable  
1012 more accurate quantification of greenhouse gas emissions from below-ground fires.

**1013 CHOICE OF EMISSION FACTORS****1014 Tier 1**

1015 The Tier 1 method uses default values for  $M_B$  along with default emissions factors provided in Tables 2.6 & 2.7  
1016 respectively.

1017 Due to the limited data available in the scientific literature, organic soils have been very broadly stratified  
1018 according to climate domain (boreal/temperate and tropical) and fire type (wild vs. prescribed). Values are  
1019 derived from the literature for all categories with the exception of prescribed fires.

1020 For all organic soil fires, the default combustion factor is 1.0, since the assumption is that all the fuel is  
1021 combusted (Yokelson *et al.*, 1997).

1022

Final Draft

1023

<b>Climate/vegetation zone</b>	<b>Sub-category</b>	<b>Mean (t d.m. ha<sup>-1</sup>)</b>	<b>95% CI (t d.m. ha<sup>-1</sup>)</b>	<b>References</b>
Boreal/temperate	Wildfire (undrained peat)	66.4	19.2	Zoltai <i>et al.</i> , 1998; Turetsky & Wieder, 2001; Benscoter & Wieder, 2003; Kasischke & Bruhwiler, 2003; Amiro <i>et al.</i> , 2001; Kajii <i>et al.</i> , 2002; Kasischke <i>et al.</i> , 1995; Pitkänen <i>et al.</i> , 1999; Cahoon <i>et al.</i> , 1994; Turetsky <i>et al.</i> , 2011a; Turetsky <i>et al.</i> , 2011b; Poulter <i>et al.</i> , 2006; de Groot & Alexander, 1986; Kuhry, 1994
	Wildfire (drained peat)	336.0	4.0	Turetsky <i>et al.</i> , 2011b
	Prescribed fire (land management)	-	-	No literature found
Tropical	Wildfire (undrained peat)	-	-	No literature found.
	Wildfire (drained peat)	353	183	Page <i>et al.</i> , 2002; Usop <i>et al.</i> , 2004; Ballhorn <i>et al.</i> , 2009
	Prescribed fire (agricultural land management)†	155	73	Saharjo & Munoz, 2005; Saharjo & Nurhayati, 2005

Note: Where fuel consumption values have been reported as t C ha<sup>-1</sup>, default values for organic soil bulk density (0.1 g cm<sup>-3</sup>)\* and carbon density (50% mass dry weight)\*\* have been applied to derive a value for mass of fuel (t ha<sup>-1</sup>) (following Akagi *et al.* 2011). At higher Tier levels, country or ecosystem specific values for both these variables are used.

\*The value for surface organic soil bulk density is an average derived from Gorham (1991) who provides a default value of 1.12 g cm<sup>-3</sup> for all northern peatlands and Page *et al.* (2011) who provide a default value of 0.09 g cm<sup>-3</sup> for all tropical peats.

\*\*The value for surface organic soil carbon content is an average derived from the typical average for eutrophic peat of 48% and the typical average for oligotrophic peat of 52% (after Lucas (1982), Immirzi *et al.* (1992) as reported in Charman (2002)).

† The consumption value excludes crop residues.

1024

1025

1026

Climate/vegetation zone	CO <sub>2</sub> -C	CO	CH <sub>4</sub>	References
Boreal/temperate	362 ± 41	207 ± 70	9 ± 4	Ward & Hardy, 1984; Yokelson <i>et al.</i> , 1997; Yokelson <i>et al.</i> , 2013
Tropical	464	210	21	Christian <i>et al.</i> , 2003
<p>1. These values have been derived from a very limited number of studies. The EF values for boreal/temperate fires are arithmetic means of the two values reported by Yokelson <i>et al.</i> (1997) for Alaska and Minnesota organic soils (carbon content 49% for Minnesota; n.d. for Alaska); of the minimum and maximum values reported by Ward and Hardy (1984) (no carbon contents reported) and the single value reported by Yokelson <i>et al.</i> (2013) for Alaskan organic soil (carbon content 42%). Surface (flaming) and deep (smouldering) organic soil fires produce a complex mixture of gases and fine particles, the nature of which will reflect vegetation type, fire behaviour, soil physical and chemical characteristics as well as the combustion conditions (in particular combustion efficiency) (Itkonen and Jantunen, 1986; NCDENR, 1998). The combustion of organic material leads to a loss of carbon; most of this is in the form of CO<sub>2</sub>, but quantities of CO, CH<sub>4</sub>, long-chain hydrocarbons, and carbon particulate matter are also emitted. Other greenhouse gases along with ozone precursors (NO<sub>x</sub>), volatile organic compounds (VOCs) and polycyclic aromatic hydrocarbons are also released (Ramadan <i>et al.</i>, 2000; Gebhart <i>et al.</i>, 2001; Honrath <i>et al.</i>, 2004; Val Martin <i>et al.</i>, 2006; Lapina <i>et al.</i>, 2008; Akagi <i>et al.</i>, 2011). Emission factors for N<sub>2</sub>O and NO<sub>x</sub> are not provided at Tier 1. There are very limited data for N<sub>2</sub>O and NO<sub>x</sub> emissions from organic soil fires and it should be noted that N<sub>2</sub>O can be produced in canisters during sample storage (e.g. Cofer <i>et al.</i>, 1990). At higher Tiers, N<sub>2</sub>O and NO<sub>x</sub> can either be measured directly or could be calculated using published emission ratios for organic soil fires (e.g. Christian <i>et al.</i>, 2003; Hamada <i>et al.</i>, 2013).</p> <p>2. The composition of organic soil fire emissions differs substantially from forest fires on mineral soils; in part this is a function of the fact that organic soil fires are dominated by smouldering rather than flaming combustion owing to the moist and often oxygen-limiting substrate conditions. Fire temperatures also differ: the typical peak temperature of smouldering organic soil fires is in the range 500-700°C, while for flaming fires it can be 1000-1500°C (Usup <i>et al.</i>, 2004; Rein, 2008). The lower temperatures and smouldering combustion associated with organic soil fires makes them harder to detect by satellites and leads to the emission of high amounts of CO relative to CO<sub>2</sub> as well as large amounts of fine particulate matter (PM<sub>2.5</sub>); fires on tropical organic soils, for example, emit as much as 3 to 6 times more particulate matter per amount of biomass consumed than other types of biomass fires (grassland, forest, plantation fires) (Heil <i>et al.</i>, 2006). The emission ratio of CO to CO<sub>2</sub> (ER<sub>CO/CO2</sub>) can be used as an indicator of the relative amount of flaming versus smouldering combustion during biomass burning with higher ER<sub>CO/CO2</sub> observed in smouldering fires (Cofer <i>et al.</i>, 1989, 1990; Christian <i>et al.</i>, 2007; Yokelson <i>et al.</i>, 2007).</p>				

1027

### 1028 Tier 2 and Tier 3

1029 At higher Tiers the approach for estimating greenhouse gas emissions from fires on organic soils incorporates  
 1030 country-specific information in Equation 2.8. When deriving higher Tier emission factors, country-specific  
 1031 combustion factors need to be developed. Regional factors for stratification could include:

- 1032 • Stratification by drainage class. Position of the soil water table is a proxy for soil moisture which determines  
 1033 depth of burn.
- 1034 • Stratification by depth of burn. This can be measured in the field post-fire (e.g., Page *et al.*, 2002; Turetsky  
 1035 & Wieder, 2003; Turetsky *et al.*, 2011a, b) or using remote sensing approaches (e.g. LiDAR) (Ballhorn *et al.*,  
 1036 2009).
- 1037 • Stratification by different fire types (wild vs. prescribed fires). GIS techniques of interpolation may be  
 1038 helpful in this analysis. Under Tier 3, one might consider annual sampling of a number of control sites.
- 1039 • Stratification by organic soil type taking into account general hydrology (e.g., bog vs. fen); vegetation  
 1040 structure (open, shrubby, forested) whenever possible.
- 1041 • Use of regionally-specific values for organic soil bulk density and carbon concentration.
- 1042 • Stratification by different land-use and management types, including differences in drainage lay-out and  
 1043 intensity, land-use intensity and practices, all of which will influence the mass of fuel available for  
 1044 combustion.

1045 Emission factors can be derived from measurements (field or laboratory based) or calculations validated against  
 1046 country-specific measurements. The literature on emissions from fires on organic soils is very sparse and

---

Final Draft

1047 countries are encouraged to share data when organic soil quality, environmental conditions and land-use  
1048 practices are similar.

1049 A higher tier approach might also use process-based models, adequately validated using observation data that  
1050 take into account temporal and spatial variations in the differences between fires on different types of organic  
1051 soils and conditions and fuel combustion efficiencies. This approach will involve a comprehensive mechanistic  
1052 understanding of combustion of organic soils, including the effects of site characteristics, drainage intensity,  
1053 vegetation cover, soil type and depth, management practices, depth of water table and soil moisture among others.  
1054 Higher Tier approaches could start by developing robust relationships between drainage and depth of burn which  
1055 could then be further refined by land management category. Models ideally also take into account fire return  
1056 interval. Fire changes organic soil chemical and physical characteristics (Yefremova & Yefremov, 1996; Zoltai  
1057 *et al.*, 1998; Milner *et al.*, 2013) as well as the rate and nature of post-fire vegetation recovery, and thus can alter  
1058 total net ecosystem productivity.

## 1059 **CHOICE OF ACTIVITY DATA**

1060 Activity data consist of areas of land remaining in a land-use category with organic soils stratified by climate  
1061 zone and fire type. Total areas should be determined according to approaches laid out in Chapter 3 of the 2006  
1062 *IPCC Guidelines* and be consistent with those reported under other sections of the inventory. The assessment of  
1063 fire-driven changes in soil carbon will be greatly facilitated if this information can be used in conjunction with  
1064 national soils and climate data, vegetation inventories, maps of burned area and other biophysical data.  
1065 Stratification of land-use categories according to climate zones, based on default or country-specific  
1066 classifications can be accomplished with overlays of land-use on suitable climate and soil maps.

### 1067 **Tier 1**

1068 Tier 1 methods require data on burned area of organic soils stratified by climate domain and fire type (wild vs.  
1069 prescribed). Data on burned area can be obtained from ground-based inventories, which can be very valuable in  
1070 areas of small fire. Some countries/regions may have an established fire inventory method in place which they  
1071 are encouraged to maintain rather than go with less comprehensive satellite methods. For larger and/or less  
1072 accessible locations, burned area data are often obtained from a time series of images from remote sensors. In  
1073 country burned area maps should ideally be mapped at Landsat TM scale (30-50 m resolution). If not available,  
1074 this could be degraded to 250 m and even 1 km data. Box 2.3 provides more details on the remote-sensing  
1075 platforms currently used for obtaining burnt area data. Other methods, such as national statistics and forest  
1076 inventory fire data can also produce suitable information in some cases, but may not be as reliable or as  
1077 comprehensive as remotely sensed data. Caution is advised regarding the use of detecting thermal anomalies  
1078 using data sets derived from satellite data. Whilst providing a reasonable indicator of the presence of a fire, one  
1079 cannot proceed to easily derive the burned area parameters required in the emission estimate equations.

1080

**Box 2.1****RECENT ADVANCES IN SATELLITE-DERIVED FIRE PRODUCTS**

Recent advances in satellite-derived fire products using MODerate resolution Imaging Spectroradiometer (MODIS) data from the Terra and Aqua satellites (Roy *et al.*, 2008; Giglio *et al.*, 2009); the Advanced Very High Resolution Radiometer (AVHRR) sensor on the National Oceanic and Atmospheric Administration (NOAA); Polar Operation Environmental Satellite (POES); the European AATSR and VEGETATION/PROBA satellites, and the Geostationary Operational Environmental Satellite (GOES) have all enabled the derivation of burned area data in near real-time and thereby enhanced the ability to estimate the areal extent of regional and global wildland fires and hence the scale of emissions (e.g. Gregoire *et al.*, 2003; Simon *et al.*, 2004; Tansey *et al.*, 2008; Giglio *et al.*, 2009; Kasischke *et al.*, 2011). Products derived from the satellite data sets either provide an indication of the area burned or an indication that a possible active fire is burning within the grid cell, which is based on a high surface temperature signal at thermal wavelengths. At the global scale, these data sets are coarse resolution (a pixel size larger than 500 m). The resulting uncertainties and particular challenges associated with commission and omission errors in remote sensing approaches to peat fire detection and characterization, however, need to be recognized and acknowledged. In normal years, for example, fires on tropical organic soils are relatively small (several hectares would be towards the upper end), and it is therefore necessary to consider using satellite data sets acquiring imagery at an appropriate resolution. During extended smouldering, fires in organic soils may be particularly difficult to pick up by sensors sensitive to thermal wavelengths. There are on-going issues with cloud cover, which are being addressed with increasing use of radar imagery. Furthermore, there are very few operational systems that can be used to develop robust and temporally stable products. The Landsat-8 mission and the forthcoming European Space Agency/European Commission Sentinel programme will help address this issue. The size of the study area is also very important as there may be existing data sets available from which a long term time series of fire disturbance can be reconstructed (e.g. 40 years of Landsat data with gap filling with radar imagery). There are useful materials on fire assessment and standards produced by the UN World Meteorological Organisation (e.g. GTOS 68, 2010).

Data on the location of organic soils can be obtained from several institutions, including ISRIC and FAO who have country-specific and global maps that include organic soils (FAO, 2012) (<http://webarchive.iiasa.ac.at/Research/LUC/External-World-soil-database/HTML/>; <http://www.fao.org/geonetwork/srv/en/main.home>; or <http://www.isric.org/>). A global consortium has been formed to make a new digital soil map of the world at fine resolution (<http://www.globalsoilmap.net/>).

**Tiers 2 and 3**

Higher Tier methods require more disaggregated and spatially explicit activity data than lower Tiers. This includes disaggregation according to drainage classes, vegetation type and condition (the latter refers to moisture, leaf on/off, and other factors); drainage depth, and land management status to improve Tier 1 estimates and may also take into account such variables as seasonal norms and modifications in water table level due to seasonal weather patterns etc. Data on depth of burn (obtained from in situ field measurements), along with country-specific data on organic soil bulk density and carbon content will also greatly improve knowledge of the mass of fuel consumed and the scale of carbon emissions. Seasonal variations in fire-driven emissions are then aggregated to annual emissions.

The accuracy of emission estimates will be further improved if information is available on land-use and its effect on organic soil condition, since fire extent and severity and hence quantity of emissions increase according to the scale of disturbance (e.g., disturbance of vegetation cover, and the presence of drainage structures associated with agriculture, forestry, peat extraction, oil and gas extraction, roads etc. (e.g., Turetsky *et al.*, 2011a, b)). Remote sensing techniques (e.g., Kasischke *et al.*, 2009) can also be used to provide an indication of the likely fire risk by estimating soil water conditions and providing an accurate proxy measure of organic soil surface water content levels and hence likely depth of burn at a landscape scale.

**CALCULATION STEPS FOR TIER 1**

The steps for estimating the CO<sub>2</sub> and non-CO<sub>2</sub> emissions from fires on drained organic soils for land remaining in a land-use category are as follows:

Final Draft

1134 **Step 1:** Using guidance in Chapter 3 of the *2006 IPCC Guidelines*, stratify areas with drained organic soils of  
 1135 land remaining in a land-use category for each land-use category according to climate domain and fire type.  
 1136 Obtain estimates of A (area burnt) from national sources or, if those are not available, from global databases.

1137 **Step 2:** Assign the appropriate fuel load ( $M_B$ ) and emission factor ( $G_{ef}$ ) from Table 2.6 and 2.7 respectively for  
 1138 the gas.

1139 **Step 3:** Estimate the CO<sub>2</sub> or non-CO<sub>2</sub> emissions by multiplying burnt area with the appropriate fuel load ( $M_B$ )  
 1140 and emission factor ( $G_{ef}$ ) from Tables 2.6 and 2.7 using Equation 2.8.

1141 **Step 4:** Repeat step 3 for each greenhouse gas using emission factors ( $G_{ef}$ ) in Table 2.7.

## 1142 **UNCERTAINTY ASSESSMENT**

1143 There are several sources of uncertainty related to estimates of CO<sub>2</sub> and non-CO<sub>2</sub> emissions from fires on organic  
 1144 soils. Fire behaviour varies greatly among wetland types and hence, disaggregation of vegetative formations will  
 1145 lead to greater precision. The fraction of fuel that is actually combusted during burning (the combustion factor)  
 1146 varies, not only between ecosystems, but also between fires, between years, and as a function of land  
 1147 management practices. Measurements from a given fire, year, and/or region cannot be extrapolated with  
 1148 confidence to other locations or years, or to the biome scale. An important cause of uncertainty is the choice of  
 1149 emission factor that partitions the smoke into CO<sub>2</sub>, CO and other trace gasses, since this is strongly driven by the  
 1150 amount of flaming versus smouldering combustion that occurs, and this can vary widely in organic soils, and is  
 1151 not well characterized from field data. In addition, the accuracy of the estimates of area burnt, proportion of the  
 1152 available fuel oxidized, and the biomass fuel available also contribute to the emissions uncertainty. Uncertainties  
 1153 of estimates of areas burnt can vary markedly depending on the methodology employed – for example, where  
 1154 very high resolution remote-sensing is used it may be of the order of ±20%, whereas the use of global fire maps  
 1155 may result in uncertainties of up to two-fold. Uncertainties in estimates of greenhouse gas emissions over large  
 1156 regions from fire are likely to be at least ±50%, even with good country-specific data, and at least two-fold where  
 1157 only default data are used. The calculation of emission errors is addressed by French *et al.* (2004). The study  
 1158 looked at the possible ranges of error in the input variables, since robust data are not available for the range of  
 1159 fire conditions and vegetation types that can burn. The sensitivity analysis revealed that ground-layer fraction  
 1160 consumed is the most important parameter in terms of output uncertainty, indicating that burning in sites with  
 1161 deep organic soils can be the most problematic in terms of uncertainty. The results of this work showed that  
 1162 input data sets are incomplete in describing the possible variability in conditions for both pre-burn and during the  
 1163 fire, and attention to improving measurements and obtaining a range of measurements is a priority for modelling  
 1164 emissions from fire in organic soils.  
 1165

## 1166 **2.3 LAND CONVERTED TO A NEW LAND-USE** 1167 **CATEGORY**

### 1168 **2.2.3 CO<sub>2</sub> emissions and removals from drained inland** 1169 **organic soils**

1170 CO<sub>2</sub> emissions/removals from land converted to a new land-use category on organic soils within the inventory  
 1171 time period are calculated in the same way as CO<sub>2</sub> emissions/removals from land remaining in a land-use  
 1172 category.<sup>1</sup> CO<sub>2</sub> emissions/removals for the lands in the conversion category are calculated using Equations 2.1  
 1173 and 2.2.

1174 On-site CO<sub>2</sub> emissions after land-use change on drained organic soils can occur from all five carbon pools. Land-  
 1175 use change can result in direct losses/gains because of biomass clearance/(re)planting. This is addressed by  
 1176 guidance for changes in the carbon pools in above-ground and below-ground biomass and dead organic matter  
 1177 on lands converted to a new land-use category provided in the *2006 IPCC Guidelines*.

1178 Land-use change can indirectly affect carbon gains and losses because of altered growth of woody biomass and  
 1179 altered respiration and organic matter oxidation through altered soil temperature. These effects are included in  
 1180 the guidance for lands remaining in a land-use category provided in the *2006 IPCC Guidelines* for above-ground  
 1181 and below-ground biomass and dead organic matter and updated emission factors in Table 2.1 of section 2.2.1.1.

<sup>1</sup> For example if a Forest Land is converted to Cropland, methodology and emission factors for Cropland are to be used.

Final Draft

1182 Additional carbon losses from biomass and soil can occur through altered fire frequency after drainage and land-  
1183 use change. These CO<sub>2</sub> emissions from fire are addressed in section 2.3.2.3.

### 1184 **2.2.3.1 ON-SITE CO<sub>2</sub> EMISSIONS/REMOVALS FROM DRAINED INLAND** 1185 **ORGANIC SOILS (CO<sub>2</sub>-C<sub>SOIL-ONSITE</sub>)**

#### 1186 **CHOICE OF METHOD**

1187 CO<sub>2</sub> emissions/removals from land converted to a new land-use category on organic soils within the inventory  
1188 time period are calculated in the same way as CO<sub>2</sub> emissions/removals from land remaining in a land-use  
1189 category. CO<sub>2</sub> emissions/removals for the lands in the conversion category are calculated using Equation 2.3 if  
1190 the soils are drained. Specific guidance by other land-use categories is given in the *2006 IPCC Guidelines*,  
1191 Chapters 5, 6, 8 and 9.

1192 At Tier 1, there is no transition period for CO<sub>2</sub> emissions from organic soils because the land immediately  
1193 switches to the methods for the new land-use category. High carbon loss from organic soils can occur after  
1194 converting natural vegetation to another land use, e.g. after converting tropical forest land to palm plantations, or  
1195 converting grassland to cropland. These additional CO<sub>2</sub>-C<sub>soil-onsite</sub> emissions in the transition phase are not  
1196 captured by the Tier 1 default emission factors shown in Table 2.1. A transitional phase is not captured by the  
1197 Tier 1 methodology due to lack of scientific data for deriving default emission factors. If a transitional phase  
1198 occurs, it should be addressed by higher tier methods. Additional guidance on the Tiers 1, 2 and 3 approaches is  
1199 given in Section 2.2.1.1.

#### 1200 **CHOICE OF EMISSION/REMOVAL FACTORS**

1201 At Tier 1, CO<sub>2</sub> emissions/removal factors for lands in the conversion category are the same as for land remaining  
1202 in a land-use category. For Tier 1 these are given in Table 2.1. Additional guidance on the Tiers 1, 2 and 3  
1203 emission/removal factors is given in Section 2.2.1.1.

1204 If land conversions on organic soils contribute significantly to CO<sub>2</sub> emissions from soils and if CO<sub>2</sub> emissions  
1205 from soils are a key category, it is *good practice* to develop country specific Tier 2 emission factors that include  
1206 the additional CO<sub>2</sub>-C<sub>soil-onsite</sub> emissions in the transition phase. Unless other country specific evidence is available  
1207 the default length of 20 years can be used for the transition phase.

1208 Tier 3 methodologies could further consider the dynamic nature of the additional CO<sub>2</sub>-C<sub>soil-onsite</sub> emissions in the  
1209 transition phase, which may be highest in the first years after the transition.

#### 1210 **CHOICE OF ACTIVITY DATA**

1211 Guidance is the same as for land remaining in a land-use category in Section 2.2.1.1.

#### 1212 **UNCERTAINTY ASSESSMENT**

1213 Guidance is the same as for land remaining in a land-use category in Section 2.2.1.1.

### 1214 **2.2.3.2 OFF-SITE CO<sub>2</sub> EMISSIONS VIA WATERBORNE CARBON** 1215 **LOSSES FROM DRAINED ORGANIC SOILS (CO<sub>2</sub>-C<sub>SOIL-ONSITE</sub>)**

#### 1216 **CHOICE OF METHOD**

##### 1217 **Tier 1**

1218 At Tier 1, CO<sub>2</sub> emissions/removals from land converted to a new land-use category on organic soils within the  
1219 inventory time period are calculated in the same way as CO<sub>2</sub> emissions/removals from land remaining in a land-  
1220 use category. Guidance is given in Section 2.2.1.2 for DOC. CO<sub>2</sub> emissions/removals for the lands in the  
1221 conversion category are calculated using Equations 2.4 and 2.5.

##### 1222 **Tier 2**

1223 The Tier 2 approach for waterborne carbon losses from organic soils incorporates country-specific information to  
1224 estimate the emissions. Tier 2 uses the same procedural steps for calculations as provided for Tier 1. Tier 2  
1225 emission factors can be developed following the same principles as for land remaining in a land-use category.  
1226 Guidance is found in Section 2.2.1.2. Generally, the same stratification should be used for land converted to a  
1227 new land-use category as is used for land remaining in a land-use category. Tier 2 approaches for land-use  
1228 changes can be further stratified according to the time since land-use change. Specific transition periods can be

Final Draft

1229 considered depending on the type of land-use change and the persistence of emissions or removals which differ  
 1230 from those on lands that have been in the new land-use category for a long time. Alternatively, the default  
 1231 transition period applicable to the new land-use category in the *2006 IPCC Guidelines* can be applied.

### 1232 **Tier 3**

1233 The development of Tier 3 approaches follows the guidance given in section 2.2.1.2 including the guidance for  
 1234 transparent documentation of Tier 3 approaches given in Section 2.2.1.1. Generally, the same approach should be  
 1235 used for land converted to a new land-use category as is used for land remaining in a land-use category. Tier 3  
 1236 methods should further differentiate transition effects of increased or reduced waterborne carbon losses after  
 1237 land-use change and the time since land-use change.

1238 Additional guidance on the Tiers 1, 2 and 3 approaches is given in Section 2.2.1.2.

### 1239 **CHOICE OF EMISSION/REMOVAL FACTORS**

1240 CO<sub>2</sub> emissions/removal factors for the lands in the conversion category are the same as for land remaining in a  
 1241 land-use category. For Tier 1 these are given in Table 2.2. Additional guidance on the Tiers 1, 2 and 3  
 1242 emission/removal factors is given in Section 2.2.1.2.

### 1243 **CHOICE OF ACTIVITY DATA**

1244 Guidance is the same as for land remaining in a land-use category in Section 2.2.1.2.

### 1245 **UNCERTAINTY ASSESSMENT**

1246 Guidance is the same as for land remaining in a land-use category in Section 2.2.1.2.

## 1247 **2.2.4 Non-CO<sub>2</sub> emissions and removals from drained** 1248 **inland organic soils**

### 1249 **2.2.4.1 CH<sub>4</sub> EMISSIONS FROM DRAINED INLAND ORGANIC SOILS**

#### 1250 **CHOICE OF METHOD**

1251 CH<sub>4</sub> emissions/removals from land converted to a new land-use category on organic soils within the inventory  
 1252 time period are calculated in the same way as CH<sub>4</sub> emissions/removals from land remaining in a land-use  
 1253 category<sup>2</sup>. CH<sub>4</sub> emissions/removals for the lands in the conversion category are calculated using Equation 2.5.  
 1254 Additional guidance on the Tiers 1, 2 and 3 approaches is given in Section 2.2.2.1.

#### 1255 **CHOICE OF EMISSION/REMOVAL FACTORS**

1256 CH<sub>4</sub> emissions/removal factors for the lands in the conversion category are the same as for land remaining in a  
 1257 land-use category. For Tier 1 these are given in Tables 2.3 and 2.4. Additional guidance on the Tiers 1, 2 and 3  
 1258 emission/removal factors is given in Section 2.2.2.1.

#### 1259 **CHOICE OF ACTIVITY DATA**

1260 Guidance is the same as for land remaining in a land-use category in Section 2.2.2.1.

#### 1261 **UNCERTAINTY ASSESSMENT**

1262 Guidance is the same as for land remaining in a land-use category in Section 2.2.2.1.

### 1263 **2.2.4.2 N<sub>2</sub>O EMISSIONS FROM DRAINED ORGANIC SOILS**

#### 1264 **CHOICE OF METHOD**

1265 N<sub>2</sub>O emissions from land converted to a new land-use category on organic soils within the inventory time period  
 1266 are calculated in the same way as N<sub>2</sub>O emissions from land remaining in a land-use category. N<sub>2</sub>O emissions for  
 1267 lands in the conversion category are calculated using Equation 2.7. Additional guidance on the Tiers 1, 2 and 3  
 1268 approaches is given in Section 2.2.2.2.

<sup>2</sup> For example if a Forest Land is converted to Cropland, methodology and emission factors for Cropland are to be used.

---

Final Draft

1269 **CHOICE OF EMISSION/REMOVAL FACTORS**

1270 N<sub>2</sub>O emission factors for the lands in the conversion category are the same as for land remaining in a land-use  
1271 category. For Tier 1 these are given in Table 2.5. Additional guidance on the Tiers 1, 2 and 3 emission/removal  
1272 factors is given in Section 2.2.2.2.

1273 **CHOICE OF ACTIVITY DATA**

1274 Guidance is the same as for land remaining in a land-use category in Section 2.2.2.2.

1275 **UNCERTAINTY ASSESSMENT**

1276 Guidance is the same as for land remaining in a land-use category in Section 2.2.2.2.

1277 **2.2.4.3 NON-CO<sub>2</sub> EMISSIONS FROM BURNING ON ORGANIC SOILS**

1278 **CHOICE OF EMISSION/REMOVAL FACTORS**

1279 Non-CO<sub>2</sub> emission factors for the lands in the conversion category are the same as for land remaining in a land-  
1280 use category. For Tier 1 these are given in Tables 2.6 and 2.7. Additional guidance on the Tiers 1, 2 and 3  
1281 emission/removal factors is given in Section 2.2.2.3.

1282 **CHOICE OF ACTIVITY DATA**

1283 Guidance is the same as for land remaining in a land-use category in Section 2.2.2.3.

1284 **UNCERTAINTY ASSESSMENT**

1285 Guidance is the same as for land remaining in a land-use category in Section 2.2.2.3.

1286

1287 **Annex 2A.1 Scientific background for developing CO<sub>2</sub>-C**  
 1288 **emission/removal factors for organic soil from the scientific**  
 1289 **literature in Table 2.1**

1290 The Tier 1 CO<sub>2</sub> emission factors presented in Table 2.1 were calculated as annual net change of the soil organic  
 1291 carbon (SOC) plus the belowground portion of the litter carbon in the different land-uses. CO<sub>2</sub> emissions were  
 1292 obtained by two well established methodologies: (1) **Flux method**: flux measurements are commonly used on all  
 1293 types of organic soils to determine gas exchange at frequencies from minutes to weeks over monitoring periods  
 1294 of up to a few years; or (2) **Subsidence method**: determining subsidence rates of drained organic soils at  
 1295 frequencies of months to years, over periods representing one to many years of subsidence.

1296 **Flux methods**

1297 The flux method uses chamber based techniques or eddy covariance in combination with auxiliary carbon pool  
 1298 data from the study sites.

1299 *Dark chamber measurements*

1300 Chamber flux measurements are made with varying frequency over short periods with dark chambers to  
 1301 determine total respiration (R<sub>t</sub>) which includes autotrophic (R<sub>a</sub>) plus heterotrophic (R<sub>h</sub>) respiration from the soil  
 1302 and heterotrophic respiration from litter. To obtain organic soil CO<sub>2</sub> emissions the observed flux (R<sub>t</sub>) must be  
 1303 adjusted for the contributions from other carbon pools (*e.g.*, litter) and autotrophic (plant root) respiration needs to  
 1304 be subtracted. For these calculations, the proportion of R<sub>h</sub> to R<sub>t</sub> was estimated from a limited number of studies.

1305 As with any mass balance approach, outputs must be balanced against inputs to calculate a net flux to the  
 1306 atmosphere. Thus, inputs in the form of root mortality and aboveground litter fall are important in calculating net  
 1307 carbon loss or gain. Tier 1 assumes that the litter pool remains constant in a land use remaining in a land use, so  
 1308 litter inputs to the SOC are equal to litterfall plus root mortality. While litterfall is relatively easy to measure,  
 1309 belowground litter inputs are hard to measure directly (Finér *et al.*, 2011; Gaudinski *et al.*, 2010; Sah *et al.*,  
 1310 2010). Estimates of litter inputs were made from a limited number of studies and were subtracted from R<sub>h</sub> to  
 1311 estimate the net flux of carbon to the atmosphere. On Peatlands Managed for Extraction no vegetation is present  
 1312 so that the net change in soil carbon was assumed to be R<sub>h</sub>.

1313 *Transparent chambers*

1314 CO<sub>2</sub> emission measurements using transparent chambers determine net ecosystem exchange (NEE) *i.e.*, the  
 1315 balance between R<sub>t</sub> and the gross primary productivity (GPP). To obtain SOC emissions the observed flux, NEE  
 1316 must be corrected for the contributions from other carbon pools (*e.g.*, litter, above-ground biomass, etc.). Design  
 1317 and use of transparent chambers is described in detail by Drösler (2005)

1318 *Eddy Covariance flux measurements*

1319 The Eddy Covariance (EC) method finds its greatest utility over larger site or landscape scales. Sophisticated  
 1320 instrumentation and data processing software calculate fluxes of gases by the covariance of gas concentrations  
 1321 with the upward and downward movements of air parcels. In its simplest interpretation for CO<sub>2</sub> fluxes the EC  
 1322 method measures NEE (the balance of ecosystem respiration and GPP). Whenever photosynthetically active  
 1323 radiation (PAR) is zero (such as at night) GPP is zero and NEE is equivalent to ecosystem respiration or R<sub>t</sub>. In  
 1324 essence the strategy for obtaining R<sub>h</sub> from EC results are the same as for transparent chambers - correction is  
 1325 required for R<sub>a</sub> (above and below ground), removals of biomass carbon, inputs of carbon from fertilizers, etc.

1326 **Subsidence Method**

1327 Drainage of an organic soil leads to subsidence or loss of elevation (Armentano and Menges, 1986; Grønlund *et al.*,  
 1328 2008; Leifeld *et al.*, 2011). Oxidative loss of carbon can be related to volume loss of the organic soil using  
 1329 bulk density and soil carbon content obtained from soil cores or pits. Total subsidence of the drained organic soil  
 1330 surface is tracked over time using elevation markers. Other markers, such as pollen have been used to correlate  
 1331 horizons among cores (Minkinen *et al.*, 1999) as an aid to determining subsidence rates.

1332 The parameters used for calculating emission in each study varied slightly. We applied a standardized approach  
 1333 to calculating the emissions from each study so that assumptions across sites would be consistent. CO<sub>2</sub> emission  
 1334 estimates are obtained by converting the volume loss to carbon via bulk density, carbon content and estimates of  
 1335 the oxidized fraction of the volume lost as compared to compaction. Bulk density was considered to remain  
 1336 constant over short periods of time and oxidation fractions were calculated from data in each paper, when  
 1337 available, or data from similar sites were used when data were not available. In all papers, carbon content was

---

Final Draft

1338 not measured, so these values were estimated using the relationship of Warren *et al.* (2012). Subsidence  
1339 emissions were corrected for dissolved organic carbon losses using Tier 1 default factors from Section 2.2.1.2.

1340 *Tropical emission/removal factors*

1341 Two types of data were available for the tropical climate zone: flux studies and studies based on subsidence.  
1342 Integrating the two approaches was problematic because the data for each approach were different. The  
1343 approach that was finally adopted was to calculate one estimate using a gain-loss approach based on flux data for  
1344 each of the gain and loss terms of the mass balance for each land use. A second estimate was calculated using  
1345 the subsidence approach, aggregated by site, with all of the available data for a standardized approach. Each  
1346 approach provided an independent estimate of the EF.

1347  
1348 There was a divergence of opinion on several points; the general approach adopted by the authors was to split the  
1349 difference on the points where this occurred. One point of divergence was over the importance of consolidation  
1350 of peat layers below the water table. Another was over the ability of surface flux measurements to adequately  
1351 capture respiration of belowground litter. Two calculations were made, one excluding one subsidence site and  
1352 including the belowground carbon inputs to the measured surface fluxes. A second calculation was made  
1353 including the site previously excluded and excluding below-ground inputs. The final EF was derived from the  
1354 average of these two calculations.

1355  
1356 Errors were propagated by the quadrature of absolute errors method (Malhi *et al.*, 2009) for each calculation.  
1357 Most estimates converged, but several estimates differed by more than 4 tonnes C ha<sup>-1</sup> y<sup>-1</sup>. These differences  
1358 were not statistically significant and means from each approach were within the 95% CI of each other. To  
1359 resolve the discrepancy between the two approaches, the final EF was determined to be the mean of the two  
1360 approaches. The uncertainty interval was taken from the highest and lowest value of the 95% CI for either  
1361 approach.

1362 **Selection of studies**

1363 A dramatic increase in published studies of CO<sub>2</sub> fluxes occurred recently but not all studies reported results that  
1364 could be used to develop Table 2.1. Many experimental studies involved manipulations other than drainage so  
1365 often their results could not be used; exceptions are results from a “control” drained site. Survey studies,  
1366 particularly on Cropland and Grassland, often involved fertilization or annual cropping where corrections were  
1367 often possible to determine Rh. Most studies in the boreal climate region and many in the temperate were  
1368 conducted seasonally – typically from April/May through September/October (in the N. Hemisphere).  
1369 Annualization of seasonal results were guided by several studies that specifically targeted winter fluxes (*e.g.*,  
1370 Alm *et al.*, 1999; Heikkinen *et al.*, 2002; Saarmio *et al.*, 2007).

Final Draft

1371 **Annex 2A.2 Derivation of ditch CH<sub>4</sub> emission factors**

1372 The Tier 1 default EFs presented in Table 2A.1 were derived from the published studies listed. The number of  
 1373 studies available remains relatively small, although some include a substantial number of individual  
 1374 measurement sites. Measured fluxes are generally quite variable within each soil/land-use type, and are not  
 1375 evenly distributed across different organic soil types (for example, most of the data for deep-drained and  
 1376 shallow-drained Grassland on organic soils are obtained from studies in The Netherlands). Tier 1 defaults for  
 1377  $EF_{CH_4-ditch}$  were derived from the mean of all data within each land-use class, and uncertainty ranges were  
 1378 calculated as 95% confidence intervals. Indicative Tier 1 default values for the fractional area of ditches within  
 1379 drained organic soils were calculated in the same way, except that data from the Netherlands were omitted from  
 1380 the Grassland classes, on the basis that fractional ditch areas are considered to be higher here than elsewhere, and  
 1381 that their inclusion would therefore lead to atypically high default values. Note that here are currently few data  
 1382 on CH<sub>4</sub> emissions from ditches in tropical organic soils or from blanket bogs. Further published data on ditch  
 1383 CH<sub>4</sub> emissions may be used to refine the default values presented in Table 2.4, or to derive country-specific Tier  
 1384 2 emission factors.

**TABLE 2A.1**  
**COLLATED DATA ON DITCH CH<sub>4</sub> EMISSIONS FROM DRAINED AND RE-WETTED ORGANIC SOILS**

<b>Organic soil/land-use type</b>	<b>Country</b>	<b>Reference</b>	<b>EF<sub>CH<sub>4</sub>-ditch</sub> (t CH<sub>4</sub>-C ha<sup>-1</sup> yr<sup>-1</sup>)</b>	<b>Frac<sub>ditch</sub></b>
Deep-drained Grassland	The Netherlands	Schrier-Uijl <i>et al.</i> , 2010, 2011	0.435	0.21
Deep-drained Grassland	The Netherlands	Vermaat <i>et al.</i> , 2011	0.592	0.25
Deep-drained Grassland	The Netherlands	Best & Jacobs, 1997	0.072	0.06
Deep-drained Grassland	UK	McNamara, 2013	0.580	0.04
Dee-drained Grassland	Russia	Sirin <i>et al.</i> , 2012	0.450	0.04
Deep-drained Grassland	Russia	Chistotin <i>et al.</i> , 2006	1.989	0.04
Deep-drained Grassland	USA	Teh <i>et al.</i> , 2011	1.704	0.05
Shallow-drained Grassland	The Netherlands	Vermaat <i>et al.</i> , 2011	0.592	0.25
Shallow-drained Grassland	The Netherlands	Best & Jacobs, 1997	0.345	0.06
Shallow-drained Grassland	The Netherlands	Van den Pol-Van Dasselaar <i>et al.</i> , 1999a,b,c	0.085	0.25
Shallow-drained Grassland	The Netherlands	Hendriks <i>et al.</i> (2007, 2010)	0.375	0.10
Drained treed bog	Canada	Roulet & Moore, 1995	0.114	0.03
Drained treed fen	Finland	Minkkinen & Laine, 2006	0.783	0.03
Drained afforested fen	Russia	Sirin <i>et al.</i> , 2012	0.139	0.02
Drained afforested fen	Russia	Glagolev <i>et al.</i> , 2008	0.088	0.04
Drained treed bog	Canada	Roulet & Moore, 1995	0.028	0.03
Drained afforested bog	Russia	Sirin <i>et al.</i> , 2012	0.301	0.01
Drained afforested bog	Russia	Sirin <i>et al.</i> , 2012	0.011	0.01
Drained afforested bog	Canada	Roulet & Moore, 1995	0.192	0.03
Drained afforested bog	Sweden	Von Arnold <i>et al.</i> , 2005b	0.013	0.02

Final Draft

Drained afforested bog	Finland	Minkinen & Laine, 2006	0.053	0.03
Peat extraction site	Finland	Nykänen <i>et al.</i> , 1995	0.133	0.02
Peat extraction site	Sweden	Sundh <i>et al.</i> , 2000	0.356	0.03
Peat extraction site	Russia	Sirin <i>et al.</i> , 2012	1.022	0.04
Peat extraction site	Russia	Chistotin <i>et al.</i> , 2006	0.797	0.04
Peat extraction site (inactive)	Finland	Hyvönen <i>et al.</i> , 2013	0.011	0.06
Peat extraction (inactive)	Canada	Waddington & Day, 2007	0.110	0.05
Drained blanket bog	UK	Cooper & Evans, 2013	0.070	0.03
Drained tropical peat (abandoned)	Indonesia	Jauhiainen & Silvennoinen, 2012	0.449	0.02
Drained tropical peat (pulpwood plantation)	Indonesia	Jauhiainen & Silvennoinen, 2012	2.939	0.02

1385

### 1386 **Annex 2A.3 Derivation of DOC emission factors**

1387 Dissolved organic carbon (DOC) is commonly the largest component of waterborne carbon loss from peatlands  
1388 and organic soils, with measured fluxes from natural peatlands ranging from 0.04 to 0.63 t C ha<sup>-1</sup> yr<sup>-1</sup>. In many  
1389 peatlands, this flux is of comparable magnitude to the rate of long-term carbon accumulation (e.g., Gorham, 1991;  
1390 Turunen *et al.*, 2004), and the size of waterborne carbon flux can therefore determine whether the site is a carbon  
1391 sink or carbon source (e.g., Billett *et al.*, 2004; Rowson *et al.*, 2010). If this DOC is subsequently converted to  
1392 CO<sub>2</sub> via photochemical or biological breakdown processes, this flux will also contribute to overall CO<sub>2</sub> emissions  
1393 from the organic soil (as an ‘off-site’ emission). This section describes the methodology that has been used to  
1394 derive emission factors for DOC losses from drained peatlands and organic soils. At present, it is not considered  
1395 possible to set reliable emission factor estimates for other forms of waterborne carbon loss, or for the effects of  
1396 specific land-use and land-use changes (other than drainage) on DOC loss. Methodological requirements to  
1397 develop these emission factors in future are described in Appendix 2a.1. The approach is based on Equation 2.5.

#### 1398 **Estimation of DOC<sub>FLUX-NATURAL</sub>**

1399 Most of the available published studies of drainage impacts on DOC loss report concentration changes relative to  
1400 undrained comparison sites, rather than direct (robust) flux measurements. On the other hand, a larger number of  
1401 studies provide reliable DOC flux estimates from natural, or near-natural, peatland systems. These two data  
1402 sources (DOC fluxes from natural sites, and DOC changes from drained-natural comparisons) were therefore  
1403 combined to derive best estimates of the DOC flux from drained sites, following Equation 2.5.

1404 Default values for DOC<sub>FLUX-NATURAL</sub> were derived from 23 published studies reporting DOC fluxes for 26 sites in  
1405 total, including natural boreal and temperate raised bogs and fens, temperate blanket bogs, and tropical peat  
1406 swamp forests (Table 2A.2). Most data were derived from catchment-scale studies with natural drainage  
1407 channels, for which accurate hydrological data are available, and to avoid double-counting of reactive DOC  
1408 exports from peatlands that are rapidly converted to CH<sub>4</sub> or CO<sub>2</sub> within the ditch network (i.e., on-site emissions).  
1409 Clear differences in flux were observed according to climate zone, with the lowest fluxes from boreal sites and  
1410 the highest fluxes from tropical sites, supporting a simple Tier 1 classification system for natural DOC flux  
1411 estimates based on this classification.

1412

Climate zone	Country	Study	DOC flux (t C ha <sup>-1</sup> yr <sup>-1</sup> )
Boreal	Finland	Juutinen <i>et al.</i> , 2013	0.037
Boreal	Canada	Moore <i>et al.</i> , 2003	0.043
Boreal	Canada	Koprivnjak & Moore, 1992	0.052
Boreal	Canada	Moore <i>et al.</i> , 2003	0.060
Boreal	Finland	Kortelainen <i>et al.</i> , 2006	0.060
Boreal	Finland	Jager <i>et al.</i> , 2009	0.078
Boreal	Sweden	Agren <i>et al.</i> , 2007	0.099
Boreal	Finland	Rantakari <i>et al.</i> , 2010	0.120
Boreal	Sweden	Nilsson <i>et al.</i> , 2008	0.130
Boreal	Finland	Kortelainen <i>et al.</i> , 2006	0.159
Temperate	Canada	Strack <i>et al.</i> , 2008	0.053
Temperate	Canada	Roulet <i>et al.</i> , 2007	0.164
Temperate	USA	Urban <i>et al.</i> , 1989	0.212
Temperate	USA	Kolka <i>et al.</i> , 1999	0.235
Temperate	Canada	Moore <i>et al.</i> , 2003	0.290
Temperate	Canada	Clair <i>et al.</i> , 2002	0.360
Temperate	UK	Dawson <i>et al.</i> , 2004	0.194
Temperate	UK	Dinsmore <i>et al.</i> , 2011	0.260
Temperate	UK	Billett <i>et al.</i> , 2010	0.234
Temperate	UK	Billett <i>et al.</i> , 2010	0.276
Temperate	Ireland	Koehler <i>et al.</i> , 2009,2011	0.140
Temperate	Australia	Di Folco & Kirkpatrick, 2011	0.134
Tropical	Indonesia	Baum <i>et al.</i> , 2008	0.470
Tropical	Indonesia	Alkhatib <i>et al.</i> , 2007	0.549
Tropical	Malaysia	Yule <i>et al.</i> , 2009; Zulkifli, 2002	0.632
Tropical	Indonesia	Moore <i>et al.</i> , 2013	0.625

1413

#### 1414 **Estimation of $\Delta\text{DOC}_{\text{DRAINAGE}}$**

1415 A total of eleven published studies were identified which provided sufficient data to calculate ratios of either  
 1416 DOC concentration or DOC flux between comparable drained and un-drained peat sites (Table 2A.3). These  
 1417 included data from boreal and temperate raised bogs and fens, blanket bogs, and tropical peats, and drainage for  
 1418 both peat extraction and land-use change to agriculture. There is a reasonable degree of consistency among the  
 1419 studies included; all show an increase in DOC following drainage, with an overall range of 15% to 118%. Most  
 1420 of the published studies suggest a DOC increase close to the mean (across all studies) of 60%, and there was  
 1421 insufficient evidence to support the use of different Tier 1  $\Delta\text{DOC}_{\text{DRAINAGE}}$  values for different peat types, climate  
 1422 zones, drainage type or drainage intensity. The use of concentration data to estimate  $\Delta\text{DOC}_{\text{DRAINAGE}}$  does,  
 1423 however, assume no corresponding change in total water flux as a result of drainage, which adds uncertainty to  
 1424 the calculated flux changes. This uncertainty should be relatively small for high-precipitation boreal/temperate  
 1425 bogs, as a large change in water flux could only occur if there is a correspondingly large change in  
 1426 evapotranspiration. For drier bog sites, drainage might be expected to increase water fluxes, therefore amplifying  
 1427 the observed concentration differences between drained and undrained sites (e.g., Strack and Zuback, 2013).  
 1428 However for fens, which are fed by external groundwater or surface water inputs rather than solely by

---

Final Draft

1429 precipitation, there is greater potential for drainage to lead to fundamental changes in hydrological functioning  
1430 (e.g., by routing lateral water inputs around the fen rather than through it), thus altering the water flux.  
1431 Consequently, although observed DOC concentration changes in drained fens are similar to those from drained  
1432 bogs (Table 2A.3), the appropriate default value of  $\Delta\text{DOC}_{\text{DRAINAGE}}$  for fens is more uncertain. At Tier 1, it could  
1433 therefore be assumed that the DOC flux from a drained fen is unchanged from the natural flux (i.e., that  
1434  $\Delta\text{DOC}_{\text{DRAINAGE}}$  is equal to zero, and the DOC export is thus equal to  $\text{DOC}_{\text{FLUX-NATURAL}}$ ). At Tier 2 it may be  
1435 possible to develop specific estimates of  $\Delta\text{DOC}_{\text{DRAINAGE}}$  based on paired comparisons between reliable DOC flux  
1436 measurements for undrained and drained fens, either on a country-specific basis or by pooling studies in different  
1437 countries. Alternatively, direct measurements of DOC export flux could be used to derive Tier 2 EFs for DOC  
1438 emissions from drained fens.

1439 Overall, the available data support a Tier 1 default  $\Delta\text{DOC}_{\text{DRAINAGE}}$  value of 0.60 for drained bogs and tropical  
1440 organic soils. Given difficulties of quantifying the water budget of drained fens, there is greater uncertainty about  
1441 the applicable value for  $\Delta\text{DOC}_{\text{DRAINAGE}}$  for this organic soil type. Therefore, countries may choose to apply the  
1442 same Tier 1 default value as in other soil types, or to make the assumption that DOC export does not increase  
1443 with drainage from fens, i.e., to apply the natural DOC flux value to calculate  $\text{EF}_{\text{DOC}}$ . An exception may also be  
1444 made where drainage channels are cut into underlying mineral soils, as this has been found to reduce DOC loss  
1445 (e.g., Moore, 2007).

1446

TABLE 2A.3 DOC CONCENTRATION (ABOVE) OR FLUX (BELOW) COMPARISONS BETWEEN DRAINED AND UNDRAINED ORGANIC SOILS, USED TO DERIVE DEFAULT VALUE FOR $\Delta\text{DOC}_{\text{DRAINAGE}}$						
Organic Soil type	Land-use	Country	Study	DOC		$\Delta\text{DOC}_{\text{DRAINAGE}}$ (%)
				Undrained	Drained	
<i>Concentration-based studies (DOC mg l<sup>-1</sup>)</i>						
Boreal bog	Drainage (peat extraction)	Canada	Glatzel <i>et al.</i> , 2003	60	110	83%
Boreal fen	Drainage	Canada	Strack <i>et al.</i> , 2008	16	24.29	53%
Boreal fen	Drainage	USA	Kane <i>et al.</i> , 2010	56	71.7	29%
Boreal fen	Drainage (peat extraction)	Finland	Heikkinen, 1990	17	20	15%
Temperate bog	Drainage	Poland	Banaś & Gos, 2004	48	71	49%
Temperate bog	Drainage (peat extraction)	New Zealand	Moore & Clarkson, 2007	70	108	54%
Temperate bog	Drainage	Czech Republic	Urbanová <i>et al.</i> , 2011	36	53.9	51%
Temperate fen	Drainage	Czech Republic	Urbanová <i>et al.</i> , 2011	17	37.5	118%
Temperate blanket bog	Drainage	UK	Wallage <i>et al.</i> , 2006	28	42.9	55%
<i>Flux-based studies (DOC g m<sup>-2</sup> yr<sup>-1</sup>)</i>						
Tropical peat	Drainage (sago palm)	Malaysia	Inubushi <i>et al.</i> , 1998	33	63	91%
Tropical peat	Drainage (agriculture)	Indonesia	Moore <i>et al.</i> , 2013	62	97	54%

1447

1448 **Estimation of  $\text{Frac}_{\text{DOC-CO}_2}$** 

1449 The significance of DOC export in terms of greenhouse gas estimation depends on its ultimate fate, i.e., whether  
1450 it is returned to the atmosphere as CO<sub>2</sub> (or even CH<sub>4</sub>), or deposited in stable forms such as lake or marine  
1451 sediments. The latter simply represents a translocation of carbon between stable stores, and should not therefore  
1452 be included in the estimation. The parameter  $\text{Frac}_{\text{DOC-CO}_2}$  sets the proportion of DOC exported from organic soils  
1453 that is ultimately converted to CO<sub>2</sub>. While uncertainty remains in the estimation of this parameter, there is  
1454 growing evidence that fluvial systems process a high proportion of incoming terrestrial carbon, and that much of  
1455 this is converted to CO<sub>2</sub> (e.g., Cole *et al.*, 2007; Wickland *et al.*, 2007; Battin *et al.*, 2009; Algesten *et al.*, 2003).  
1456 Both Jonsson *et al.* (2007) and Algesten *et al.* (2003) estimated that around 50% of all terrestrially-derived  
1457 organic carbon was mineralised within large, lake-influenced catchments in Sweden. Wickland *et al.* (2007)  
1458 measured 6% to 15% conversion of pore-water DOC to CO<sub>2</sub>, and 10% to 90% conversion of the vegetation-  
1459 derived DOC, during one-month dark incubations, while Raymond & Bauer (2001) measured 63%  
1460 biodegradation of riverine DOC during a one-year dark incubation. Multiple studies showing a strong correlation  
1461 between lake DOC concentration and dissolved CO<sub>2</sub> concentrations (e.g., Sobek *et al.*, 2003; Stutter *et al.*, 2011  
1462 and references therein) all suggest widespread conversion of DOC to CO<sub>2</sub> in lakes. Dawson *et al.* (2001)  
1463 estimated that 12-18% of DOC was removed within a 2 km stream reach. Experiments undertaken on light-  
1464 exposed samples of peat-derived waters (Köhler *et al.*, 2002; Worrall *et al.*, 2013; Jones *et al.*, 2013)  
1465 consistently show rapid and extensive DOC loss, with averages ranging from 33% to 75% over periods of up to  
1466 10 days. Both Köhler *et al.* (2002) and Jones *et al.* (2013) found that peat-derived DOC was more susceptible to  
1467 photo-degradation compared to DOC from other water sources, and Köhler *et al.* (2002) found that most of the  
1468 DOC lost was converted to CO<sub>2</sub> (e.g., Opsahl and Benner, 1998). Jones *et al.* (2013) observed that since much  
1469 of this degradation occurs within the first 48 hours, this would be sufficient to convert most peat-derived DOC to  
1470 CO<sub>2</sub> before it enters the sea. Overall, Algesten *et al.* (2003) estimated that 90% of the DOC removal in their large  
1471 catchments was due to mineralisation to CO<sub>2</sub>, with only 10% buried in lake sediments. Terrestrially-derived  
1472 DOC which does reach the sea largely appears to be photo-chemically or microbially processed in the marine  
1473 system, mostly within years to decades (Bianchi, 2011; Opsahl and Benner, 1997).

1474 In summary, there is strong evidence that a high proportion of peat-derived DOC is mineralized rapidly in  
1475 headwaters; that this processing continues at a relatively high rate through rivers and lakes; and that any peat-  
1476 derived DOC that does reach the sea will nevertheless largely be mineralized in the marine ecosystem. These

---

Final Draft

1477 observations support the use of a high value for  $Frac_{DOC-CO_2}$ . Taking the ratio of mineralisation to sediment burial  
1478 obtained by Algesten *et al.* (2003), and assuming that a similar ratio applies to any DOC exported to the ocean,  
1479 would suggest that around 90% of peat-derived DOC is eventually converted to  $CO_2$ . On this basis a Tier 1  
1480 default value of 0.9 is proposed, with an uncertainty range of 0.8-1.0 to reflect uncertainties in the proportion of  
1481 DOC returned to burial in lake or marine sediments.

1482 There is some evidence that controlled burning (for moorland management) also increases DOC losses (e.g.,  
1483 Yallop *et al.*, 2010; di Folco & Kirkpatrick, 2011), although other experimental studies have shown no effect  
1484 (e.g., Ward *et al.*, 2007; Worrall *et al.*, 2007). A precautionary estimate is that managed burning may increase  
1485 mean DOC loss by 20-50%, but further work is required to resolve uncertainties on this issue (Holden *et al.*,  
1486 2012). Grazing levels on semi-natural vegetation have not been shown to affect DOC loss (Ward *et al.*, 2007;  
1487 Worrall *et al.*, 2007), and data on the effects of more intensive agricultural (Grassland and Cropland)  
1488 management on DOC loss are currently insufficient to estimate an emissions factor. Therefore, generic values for  
1489 the effects of drainage may be used.

1490

1491 **Annex 2A.4 Derivation of CO<sub>2</sub>-C and non-CO<sub>2</sub> emission**  
1492 **factors for emissions from burning of drained organic soils from**  
1493 **scientific literature in Tables 2.6 and 2.7**

1494 CO<sub>2</sub> emission factors for fires on drained organic soils were obtained by a consideration of the available  
1495 scientific literature. The data presented in Table 2.6 and Table 2.7 provide default values for mass of available  
1496 fuel and emissions factors.

1497 The data in Table 2.6 were obtained using a variety of different approaches to calculate the mass of fuel  
1498 combusted. It should be noted that there are only a limited number of publications providing ground- or  
1499 laboratory-based data on the depth (i.e. volume) of soil organic material consumed. Quantitative estimation of  
1500 depth of burn as well as organic soil characteristics (i.e. bulk density and carbon content) are not easy to  
1501 determine in the field, thus information on these key parameters is often based on theoretical assumptions or  
1502 limited ground measurements. This knowledge gap contributes considerably to the overall uncertainties related  
1503 to emissions from fires on organic soils because it is difficult to accurately assess the amount of fuel that is  
1504 consumed. Field data of depth of burn are available from a number of studies of fires on organic soils in northern  
1505 forests and peatlands in North America, Europe and Asia (e.g., Zoltai *et al.*, 1998; Turetsky & Wieder, 2001;  
1506 Page *et al.*, 2002; Benscoter & Wieder, 2003; Ballhorn *et al.*, 2009; de Groot *et al.*, 2009; Turetsky *et al.*, 2011a,  
1507 b), while in other cases, data have been extrapolated from previous studies.

1508 Obtaining accurate field data on the depth of combustion on organic soils is problematic since there is usually a  
1509 lack of reference data. Turetsky & Wieder (2001) developed a method for field assessment that considered the  
1510 rooting depth of trees, while other studies have used comparison of adjacent unburned sites to quantify  
1511 combustion depth (e.g., Kasischke, 2000; Page *et al.*, 2002; de Groot *et al.*, 2009; Turetsky *et al.*, 2011a) or  
1512 measurement of fuel loads before and after experimental fires (e.g., Usup *et al.*, 2004). The use of LiDAR remote  
1513 sensing has also been applied in one study (Ballhorn *et al.*, 2009).

1514 Nearly all the data presented in Table 2.6 for the boreal and temperate zones are actually from the boreal zone,  
1515 with only one study in the temperate zone (Poulter *et al.*, 2006) and two studies in tropical zone (Ballon *et al.*,  
1516 2009; Page *et al.*, 2002). Most studies are of wildfires (i.e. unwanted and unplanned fires ignited other than by  
1517 prescription (e.g., by lightning or as a result of human activities, including escaped prescribed fires as well as  
1518 those started through negligence or by arson) and are for fires on undrained peatland organic soils. Only  
1519 Turetsky *et al.* (2011b) provide depth of burn data for a wildfire on a drained boreal organic soil. In addition,  
1520 there are no data for organic soil losses associated with prescribed fires in the boreal/temperate zone but some  
1521 studies to suggest that DOC increases following fire (see also Annex 2A.2). Most prescribed (i.e. managed) fires  
1522 on the vegetation of organic soils probably result in either no or only minimal ignition loss of soil carbon.

1523 Fuel moisture content, depth of water table and burn history will all determine the extent of organic soil  
1524 combustion during a prescribed fire but the scale of loss will often depend on the skill and experience of the fire  
1525 manager. In some parts of the temperate zone, prescribed rotational burning of vegetation on organic soils is a  
1526 long-established land management practice. In the UK it is carried out on about 18% of peatlands,  
1527 predominantly in the uplands (Marsden & Ebmeier, 2012), with the aim of removing the older, less productive  
1528 vegetation and encouraging new growth for livestock grazing and cover for game birds (Worrall *et al.* 2010). In  
1529 North America, prescribed burning of vegetation on organic soils is also practiced, with a range of benefits  
1530 including the reduction of wildfire hazards, improvement of wildlife habitats and restoration of ecosystem  
1531 diversity and health (e.g., Christensen, 1977). Typically prescribed burning will be carried out when fuel  
1532 moisture is high enough to prevent combustion of the organic soil but low enough to carry a surface fire, thus  
1533 reducing the risk of soil ignition. Shifts in climate have narrowed the window of opportunity for prescribed  
1534 burning and changes in weather patterns have resulted in unexpected drying of peatlands during on-going  
1535 prescription burns. Some local fire managers have recognised this shift, but unfortunately this is a minimally  
1536 studied area and little information exists on the scale of emissions arising from the combustion of organic soils  
1537 during prescription burns. At Tier 1, it is assumed that there is either no or very little combustive loss of soil  
1538 organic matter during prescribed fires on organic soils.

1539 For tropical organic soils, the average depth of burn has not been explored in a consistent way that  
1540 representatively covers the different geographical regions, vegetation types or the different fire types (i.e. wild vs.  
1541 prescribed fires). There have been a limited number of field measurements of depth of burn and estimates of  
1542 organic soil combustion losses. These have used either direct field measurements (e.g., Page *et al.*, 2002; Usup *et al.*,  
1543 2004) or a combination of field measurements and LiDAR data (e.g., Ballhorn *et al.*, 2009). There are only  
1544 three studies of wildfires on drained organic soils and none in undrained organic soils, although studies have  
1545 demonstrated that in an intact condition tropical peat swamp forest is at very low risk of fire (e.g., Page *et al.*,

---

Final Draft

1546 2002). There have been a limited number of studies investigating depth of burn on drained organic soils under  
1547 agricultural management (e.g., Saharjo & Munoz, 2005). Prescribed agricultural burning is undertaken on both a  
1548 small and large scale to improve soil fertility and/or to remove forest or crop residues during land preparation  
1549 activities. For example, traditional ‘sonor’ rice cultivation on shallow organic soils involves regular burning of  
1550 crop residues along with the soil surface to enhance soil fertility. In addition to field measurements, there have  
1551 been limited laboratory-based burn tests aimed at establishing the environmental controls on depth of organic  
1552 soil combustion (e.g., Benschoter *et al.*, 2011). While more field and laboratory experiments to determine fuel  
1553 consumption during fires on organic soils are needed (French *et al.*, 2004) there is also a need for improved  
1554 remote sensing methods to aid burn severity mapping in peatlands (defined as the magnitude of ecological  
1555 changes between pre- and post-fire conditions) which can provide an indication of the likely depth of burn. Burn  
1556 severity is not easy to either investigate or quantify but there have been a limited number of studies using  
1557 spectral indices to discriminate different levels of burn severity in boreal and temperate forests (e.g., van  
1558 Wagtendonk *et al.*, 2004; Epting *et al.*, 2005; Hall *et al.*, 2008) but only one study to date of tropical organic  
1559 soils (Hoscilo *et al.*, 2013). Even regionally developed consumption models can have large uncertainties with  
1560 respect to organic soils consumption. The development of robust methodologies to assess burn severity and total  
1561 organic soil consumption in wetlands would enable more accurate quantification of carbon emissions from both  
1562 above and below-ground fires for reporting at higher tiers.

1563 Accurate assessment of the volume of organic soil combusted during a fire will only be feasible at higher Tier 2  
1564 and Tier 3 levels, while at Tier 1 level some simplifying assumptions are required.

1565 **Appendix 2a.1 Estimation for Particulate Organic Carbon**  
 1566 **(POC) and Dissolved Inorganic Carbon (DIC) loss from**  
 1567 **peatlands and drained organic soils: Basis for future**  
 1568 **methodological development**

1569 This Appendix provides a basis for future methodological development rather than complete guidance.

1570 **Particulate Organic Carbon**

1571 Particulate organic carbon (POC) is generally a negligible component of the carbon balance of natural peatlands  
 1572 and organic soils. However, disturbance of organic soils through land-use change, including drainage (which can  
 1573 include the dredging of peat from drains and canals), burning (managed burning and wildfire), conversion to  
 1574 arable and peat extraction, can all result in high rates of POC loss via waterborne erosion and also wind erosion.  
 1575 In actively eroding blanket bogs, POC losses in excess of  $100 \text{ g C m}^{-2} \text{ yr}^{-1}$  may represent the dominant form of  
 1576 soil carbon loss (e.g., Pawson *et al.*, 2008; Worrall *et al.*, 2011).

1577 Available data suggest that the key determinant of POC loss is the proportion of the total area occupied by  
 1578 exposed (bare) peat, according to Equation 2A.1. The bare peat area,  $PEAT_{BARE}$ , would include unvegetated  
 1579 drainage ditches, erosion gullies, peat extraction surfaces, and areas of the soil surface exposed by burning,  
 1580 intensive grazing or the deposition of peat dredged from drainage channels onto the land surface. For Cropland,  
 1581 some estimation of the annual average proportion of the organic soil surface exposed over the full crop rotation  
 1582 would be required. Data from eroding UK blanket bogs suggest that waterborne POC exports can be reasonably  
 1583 well-predicted based on a POC flux from bare peat surfaces ( $POC_{FLUX\_BAREPEAT}$ ) of around  $4 \text{ t C ha}^{-1} \text{ yr}^{-1}$   
 1584 (Goulsbra *et al.*, 2013). Further work is required to establish whether different values would be applicable to  
 1585 other soil types, land-use types and climate regimes (in particular whether it is dependent on precipitation  
 1586 amount or intensity). At present there are few data on which to base an estimate of airborne POC loss, and  
 1587 further work is required to quantify this loss term, which may be large in peat extraction and cropland sites.

1588 Finally, there is limited information currently available from which to derive a value for the proportion of POC  
 1589 ultimately converted to  $CO_2$ , ( $Frac_{POC-CO_2}$ ). Unlike DOC, a substantial proportion of POC is mobilized from  
 1590 organic soils through physical erosion processes, and its reactivity in fluvial systems is uncertain. Some studies  
 1591 have shown fairly high rates of POC turnover in river and estuarine systems (e.g., Sinsabaugh and Findlay, 1995),  
 1592 and POC redeposited on floodplains may be subject to moderate rates of oxidation (Goulsbra *et al.*, 2013).  
 1593 However, it is likely that a significant proportion of waterborne POC loss from organic soils may simply be  
 1594 transferred to lake or coastal sediments, re-deposited on floodplains, or transported to other land areas via aeolian  
 1595 transport, rather than converted to  $CO_2$ . Further research is therefore needed to establish realistic ranges for  
 1596  $Frac_{POC-CO_2}$  in different systems.

1597 **EQUATION 2A.1**  
 1598 **CALCULATION OF POC EXPORT FROM DRAINED ORGANIC SOILS**

$$1599 \quad EF_{POC} = POC_{FLUX\_BAREPEAT} \bullet PEAT_{BARE} \bullet Frac_{POC-CO_2}$$

1600  
 1601 Where:

1602  $EF_{POC}$  = POC emission factor,  $\text{t C ha}^{-1} \text{ yr}^{-1}$

1603  $POC_{FLUX\_BAREPEAT}$  = Flux of POC from a bare peat surface,  $\text{t C ha}^{-1} \text{ yr}^{-1}$

1604  $PEAT_{BARE}$  = Proportion of the ground surface occupied by exposed peat

1605  $Frac_{POC-CO_2}$  = Conversion factor for the fraction of POC converted to  $CO_2$  following export from site

1606

1607 **Dissolved Inorganic Carbon**

1608  
 1609 Waterborne carbon fluxes from organic soils, comprising bicarbonate ion ( $HCO_3^-$ ), carbonate ions ( $CO_3^{2-}$ ) and  
 1610 free  $CO_2$ , are collectively termed dissolved inorganic carbon (DIC). These different carbon species exist in  
 1611 equilibrium, depending primarily on the pH of the water. In water draining low-pH organic soils (i.e. bogs),  
 1612 almost all DIC exists is present as  $CO_2$ . Most of this  $CO_2$  derives from autotrophic and heterotrophic respiration  
 1613 within organic soils, and is transferred laterally from soils into drainage waters, where it is consistently present at

Final Draft

1614 concentrations well in excess of atmospheric CO<sub>2</sub> concentrations. This supersaturated CO<sub>2</sub> will be emitted  
1615 ('evaded' or 'degassed') to the atmosphere, typically within a few kilometres of its source (e.g., Hope *et al.*,  
1616 2001). Limited measurements of CO<sub>2</sub> evasion from natural peatlands suggest that this emission is a  
1617 quantitatively significant component of the overall carbon budget. For example, Dinsmore *et al.* (2010) recorded  
1618 a DIC flux of 0.12 to 0.16 t C ha<sup>-1</sup> yr<sup>-1</sup> at a Scottish peatland catchment, of which over 90% was evaded to the  
1619 atmosphere within the first 5 km of the stream length. Although this may be considered an 'on site' emission, in  
1620 practice it will not be measured as part of the terrestrial CO<sub>2</sub> emission using chamber-based methods, and is  
1621 unlikely to be captured by eddy covariance methods. Consequently, direct measurements of CO<sub>2</sub> emissions from  
1622 water bodies draining organic soils (e.g., using floating chambers or gas transfer coefficients linked to  
1623 measurements of dissolved CO<sub>2</sub> within the water column) are likely to be required in order to obtain reliable  
1624 estimates of this component of the carbon flux. Currently, only a few such measurements are available for  
1625 undrained organic soils (e.g., Hope *et al.*, 2001; Billett and Moore, 2008; Dinsmore *et al.*, 2009; Dinsmore *et al.*,  
1626 2010; Wallin *et al.*, 2012). For drained organic soils, insufficient data are currently available to permit default  
1627 emission factors to be developed. Further measurements of CO<sub>2</sub> evasion for a range of climate zones, soil types,  
1628 land-use classes and drainage systems are therefore required to support future methodological development in  
1629 this area. Care is required to avoid double-counting of CO<sub>2</sub> emissions associated with mineralisation of DOC  
1630 within downstream water bodies, as opposed to the direct degassing of CO<sub>2</sub> released from the organic soil into  
1631 the water body.

1632  
1633 As noted above, other components of the DIC flux can be considered minor for bogs, due to their low pH. This is  
1634 not the case for fens, which have a higher pH, so that HCO<sub>3</sub><sup>-</sup> and CO<sub>3</sub><sup>2-</sup> may form significant components of the  
1635 total DIC export. However, a high proportion of this flux may derive from weathering processes external to the  
1636 organic soil (i.e. in groundwater or river water inputs to the fen) and this geogenic flux cannot be considered a  
1637 part of the internal carbon budget of the organic soil. On the other hand, autotrophic and heterotrophic respiration  
1638 processes may also generate dissolved CO<sub>2</sub>, which can then dissociate to form HCO<sub>3</sub><sup>-</sup> and CO<sub>3</sub><sup>2-</sup> in alkaline  
1639 waters. This flux *does* form a component of the organic soil carbon balance, but further work is needed in order  
1640 to i) quantify this flux (particularly for drained organic soils); ii) differentiate this biogenic DIC from geogenic  
1641 DIC (for example using isotopic methods); and iii) determine the proportion of DIC exported from organic soils  
1642 which is ultimately returned to the atmosphere as CO<sub>2</sub>, rather than sequestered into sediments, such as marine  
1643 carbonate deposits.

1644  
1645 Finally, available data consistently suggest that, other than emissions from drainage ditches (see Section 2.2.2.1),  
1646 on- or off-site emissions of dissolved CH<sub>4</sub> from water bodies represent a negligible component of the total  
1647 carbon and greenhouse gas budget of organic soils (e.g., Hope *et al.*, 2001; Dinsmore *et al.*, 2010; Billett and  
1648 Harvey, 2013).  
1649

1650 **References**

- 1651 Ågren, A., Jansson, M., Ivarsson, H., Bishop, K., Seibert, J. 2007. Seasonal and runoff-related changes in total  
1652 organic carbon concentrations in the River Öre, Northern Sweden. *Aquatic Science* doi 10.1007/s00027-007-  
1653 0943-9.
- 1654 Ahlholm, U. and Silvola, J. 1990. Turvetuotannon ja turpeen käytön osuus maapallon ja Suomen hiilitaseessa,  
1655 Ministry of Trade and Industry, Ser. D 183, 1–57, [in Finnish].
- 1656 Akagi, S. K., R. J. Yokelson, C. Wiedinmyer, M. J. Alvarado, J. S. Reid, T. Karl, J. D. Crounse and P. O.  
1657 Wennberg (2011). Emission factors for open and domestic biomass burning for use in atmospheric models.  
1658 *Atmospheric Chemistry and Physics* 11: 4039-4072.
- 1659 Algesten, G., Sobek, S., Bergström, A-K., Ågren, A., Tranvik, L., Jansson, M. 2003. Role of lakes for organic  
1660 carbon cycling in the boreal zone. *Global Change Biol.*, 10, 141-147.
- 1661 Ali, M., Taylor, D., Inubushi, K. 2006. Effects of environmental variations on CO<sub>2</sub> efflux from a tropical  
1662 peatland in Eastern Sumatra. *Wetlands* 26, 612-618.
- 1663 Alkhatib, M., Jennerjahn, T.C., Samiaji, J. 2007. Biogeochemistry of the Dumai River estuary, Sumatra,  
1664 Indonesia, a tropical blackwater river. *Limnol. Oceanogr.*, 52: 2410–2417.
- 1665 Alm J, Schulman L, Walden J, Nykänen H, Martikainen PJ, Silvola J (1999a) Carbon balance of a boreal bog  
1666 during a year with an exceptionally dry summer. *Ecology*, 80(1), 161-174.
- 1667 Alm, J., Shurpali, N. J., Minkkinen, K., Aro, L., Hytönen, J., Laurila, T., Lohila, A., Maljanen, M., Mäkiranta, P.,  
1668 Penttilä, T., Saarnio, S., Silvan, N., Tuittila, E.-S. and Laine, J. 2007. Emission factors and their uncertainty  
1669 for the exchange of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O in Finnish managed peatlands, *Boreal Environ. Res.*, 12, 191–209.
- 1670 Amiro, B. D., J. B. Todd, B. M. Wotton, K. A. Logan, M. D. Flannigan, B. J. Stocks, J. A. Mason, D. J. Martell  
1671 and K. G. Hirsch (2001). Direct carbon emissions from Canadian forest fires, 1959-1999. *Canadian Journal*  
1672 *of Forest Research* 31: 512-525.
- 1673 Andreae, M.O., Merlet, P. 2001. Emission of trace gases and aerosols from biomass burning. *Global*  
1674 *Biogeochemical Cycles*, 15:955–966.
- 1675 Augustin, J., Merbach, W., Steffens, L., Snelinski, B. 1998. Nitrous oxide fluxes of disturbed minerotrophic  
1676 peatlands. *Agribiological Research*, 51, 47–57. Aulakh MS, Bijay-Singh 1997. Nitrogen losses and fertilizer  
1677 N use efficiency in irrigated porous soils. *Nutrient Cycling in Agroecosystems* 47, 197–212.
- 1678 Aulakh MS, Singh, B. 1997. Nitrogen losses and fertilizer N use efficiency in irrigated porous soils. *Nutr. cycl.*  
1679 *Agroecosyst.*, 47, 197–212.
- 1680 Ballhorn, U., F. Siegert, M. Mason and S. Limin (2009). Derivation of burn scar depths and estimation of carbon  
1681 emissions with LIDAR in Indonesian peatlands. *Proceedings of the National Academy of Sciences* 106:  
1682 21213–21218.
- 1683 Banaś, K., Gos, K. 2004. Effect of peat-bog reclamation on the physico-chemical characteristics of ground water  
1684 in peat. *Polish J. Ecol.* 52: 69-74.
- 1685 Basuki, S., Suwardi, Munoz, C.P., 2012. Emission of CO<sub>2</sub> and CH<sub>4</sub> from plantation forest of *Acacia crassiparpa*  
1686 on peatlands in Indonesia. 14th International peat congress, Stockholm, Sweden, 3-8 June 2012.
- 1687 Battin, T.J., Luyssaert, S., Kaplan L.A., Aufdenkampe, A.K., Richter, A., Tranvik, L.J. 2009. The boundless  
1688 carbon cycle. *Nature Geosci.* 2: 598-600.
- 1689 Baum, A., Rixen, T., Samiaji, J. 2007. Relevance of peat draining rivers in central Sumatra for the riverine input  
1690 of dissolved organic carbon into the ocean. *Estuarine Coastal Shelf Sci.* 73: 563-580.
- 1691 Benschoter, B. W., D. K. Thompson, J. M. Waddington, M. D. Flannigan, B. M. Wotton, W. J. de Groot and M. R.  
1692 Turetsky (2011). Interactive effects of vegetation, soil moisture, and bulk density on depth of burning of thick  
1693 organic soils. *International Journal of Wildland Fire* 20: 418-429.
- 1694 Best, E.P.H., Jacobs, F.H.H. 1997. The influence of raised water table levels on carbon dioxide and methane  
1695 production in ditch dissected peat Grasslands in the Netherlands. *Ecol. Eng.* 8: 129-144.
- 1696 Bianchi T.S. 2011. The role of terrestrially derived organic carbon in the coastal ocean: A changing paradigm  
1697 and the priming effect. *Proc. Nat. Acad. Sci.* 108: 19473–19481.

Final Draft

- 1698 Billett, M.F., & Harvey, F.H. (2013). Measurements of CO<sub>2</sub> and CH<sub>4</sub> evasion from UK Peatland headwater  
1699 streams. *Biogeochemistry*, DOI 10.1007/s10533-012-9798-9.
- 1700 Billett, M.F., Moore, T.R. (2008) Supersaturation and evasion of CO<sub>2</sub> and CH<sub>4</sub> in surface waters at Mer Bleue  
1701 peatland, Canada. *Hydrol. Process.* 22, 2044-2054.
- 1702 Billett M.F., Charman, D.J., Clark, J.M., Evans, C.D., Evans, M.G., Ostle, N.J., Worrall, F., Burden, A.,  
1703 Dinsmore, K.J., Jones, T., McNamara, N.P., Parry, L., Rowson, J.G., Rose, R. 2010. Carbon balance of UK  
1704 peatlands: current state of knowledge and future research challenges. *Climate Research*, 45: 13-29.
- 1705 Billett, M.F., Palmer, S.M., Hope, D., Deacon, C., Storeton-West, R., Hargreaves, K.J., Flechard, C., Fowler, D.  
1706 2004. Linking land-atmosphere-stream carbon fluxes in a lowland peatland system. *Global Biogeochem.*  
1707 *Cycl.* 18: GB1024.
- 1708 Blodau, C. (2002): Carbon cycling in peatlands: A review of processes and controls. *Environmental Reviews* 10:  
1709 111-134.
- 1710 Brady, M.A., 1997. Organic matter dynamics of coastal peat deposits in Sumatry, Indonesia, Department of  
1711 Forestry. The University of British Columbia, Vancouver, p. 258.
- 1712 Bremner JM 1997. Sources of nitrous oxide in soils. *Nutr. Cycl. Agroecosyst.*, 49, 7–16.
- 1713 Cahoon, D. J., B. J. Stocks, J. Levine, W. Cofer and J. Pierson (1994). Satellite analysis of the severe 1987 forest  
1714 fire in northern China and southeastern Siberia. *Journal of Geophysical Research* 99: 18 627 –  
1715 618 638.
- 1716 Charman, D. (2002). *Peatlands and Environmental Change*. Chichester, U.K., Wiley.
- 1717 Chimner, R.A., and K.C. Ewel (2005), A tropical freshwater wetland: II. Production, decomposition, and peat  
1718 formation, *Wetlands Ecology and Management*, 13, 671-684.
- 1719 Chimner, R.A., Ewel, K.C. (2004) Differences in carbon fluxes between forested and cultivated micronesia  
1720 tropical peatlands. *Wetlands Ecology and Management* 12, 419-427.
- 1721 Chimner, R.A., 2004. Soil respiration rates of tropical peatlands in Micronesia and Hawaii. *Wetlands* 24, 51–56.
- 1722 Chistotin M.V., Sirin A.A., Dulov L.E. 2006. Seasonal dynamics of carbon dioxide and methane emission from  
1723 a peatland in Moscow Region drained for peat extraction and agricultural use. *Agrokhimija* 6: 54–62.
- 1724 Christensen, N. L. (1977). Fire and soil-plant nutrient relations in a pine-wiregrass savanna on the coastal plain  
1725 of North Carolina. *Oecologia* 31: 27-44.
- 1726 Christian, T. J., B. Kleiss, R. J. Yokelson, R. Holzinger, P. J. Crutzen, W. M. Hao, B. H. Saharjo and D. E. Ward  
1727 (2003). Comprehensive laboratory measurements of biomass-burning emissions: 1. Emissions from  
1728 Indonesian, African and other fuels. *Journal of Geophysical Research* 108: No. D23, 4719,  
1729 doi:4710.1029/2003JD003704.
- 1730 Christian, T. J., R. J. Yokelson, J. A. Carvalho Jr., D. W. T. Griffith, E. C. Alvarado, J. C. Santos, T. G. S. Neto,  
1731 C. A. G. Veras and W. M. Hao (2007). The tropical forest and fire emissions experiment: Trace gases emitted  
1732 by smoldering logs and dung from deforestation and pasture fires in Brazil. *Journal of Geophysical Research*  
1733 112: D18308, doi:18310.11029/12006JD008147.
- 1734 Clair, T.A., Arp, P., Moore, T.R., Dalvac, M., Meng, F-R. 2002. Gaseous carbon dioxide and methane, as well as  
1735 dissolved organic carbon losses from a small temperate wetland under a changing climate. *Environ. Pollut.*  
1736 116: S143-S148.
- 1737 Clymo, R.S. and Reddaway, E.J.F. 1971. Productivity of Sphagnum (bog-moss) and peat accumulation.  
1738 *Hidrobiologia* 12: 181–192.
- 1739 Cofer, W. R., III, J. L. Levine, E. L. Winstead and B. J. Stocks. 1990. Gaseous emissions from Canadian boreal  
1740 forest fires. *Atmosphere and Environment, Part A* 24: 1653–1659.
- 1741 Cofer, W. R., III, J. S. Levine, D. I. Sebacher, E. L. Winstead, P. J. Riggan, B. J. Stocks, J. A. Brass, V. G.  
1742 Ambrosia and P. J. Bost 1989. Trace gas emissions from chaparral and boreal forest fuels. *Journal of*  
1743 *Geophysical Research* 94: 2255–2259
- 1744 Cofer III, W.R., Levine, J.S., Winstead, E.L. and Stocks, B.J. (1990) Gaseous emissions from Canadian boreal  
1745 forest fires. *Atmos. Environ. A-Gen.*, 24, 1653-1659..

Final Draft

- 1746 Cole, J.J., Prairie, Y.T., Caraco, N.F., McDowell, W.H., Tranvik, L.J., Striegl, R.G., Duarte, C.M., Kortelainen,  
1747 P., Downing, J.A., Middelburg, J.J., Melack, J. 2007. Plumbing the global carbon cycle: Integrating inland  
1748 waters into the terrestrial carbon budget. *Ecosystems* 10: 171-184
- 1749 Comeau, L.-P., Hergoualc'h, K., Smith, J. U. and Verchot, L. 2013. Conversion of intact peat swamp forest to oil  
1750 palm plantation: Effects on soil CO<sub>2</sub> fluxes in Jambi, Sumatra. Working Paper 110. CIFOR, Bogor, Indonesia.
- 1751 Comeau, L.-P., Hergoualc'h, K., Verchot, L.V., 2012. Soil respiration rates along a forest conversion to oil palm  
1752 plantation transition in Jambi, Sumatra, Indonesia. Technical report.
- 1753 Cooper M., Evans, C. 2013. CH<sub>4</sub> emissions from ditches in a drained upland blanket bog, North Wales, UK. In:  
1754 Emissions of greenhouse gases associated with peatland drainage waters: Report to Defra under project  
1755 SP1205: Greenhouse Gas Emissions Associated with Non Gaseous Losses of Carbon from Peatlands - fate of  
1756 Particulate and Dissolved Carbon. Report to the Department of Environment, Food and Rural Affairs,  
1757 UK. Couwenberg, J., Thiele, A., Tanneberger, F., Augustin, J., Bährisch, S., Dubovik, D., Liashchynskaya, N.,  
1758 Michaelis, D., Minke, M., Skuratovich, A. & Joosten, H. (2011) Assessing greenhouse gas emissions from  
1759 peatlands using vegetation as a proxy. *Hydrobiologia*, 674, 67–89.
- 1760 Couwenberg, J. and Hooijer, A. and, 2013. Towards robust subsidence-based soil carbon emission factors for peat  
1761 soils in south-east Asia, with special reference to oil palm plantations. *Mires and Peat*, 12:1–13.
- 1762 Czaplak I., Dembek W. 2000. Polish peatlands as a source of emission of greenhouse gases. *Zeszyty Edukacyjne*  
1763 wyd. IMUZ, 6: 61-71.
- 1764 Dariah, A., Marwanto, S., Agus, F., 2013. Peat CO<sub>2</sub> emissions from oil palm plantations, separating root-related  
1765 and heterotrophic respirations. Submitted to Mitigation and Adaptation Strategies for Global Change.
- 1766 Darung, U., Morishita, T., Takakai, F., Dohong, S., Limin, H.S., Hatano, R., (2005) The effects of forest fire and  
1767 agriculture on CO<sub>2</sub> emissions from tropical peatlands, Central Kalimantan, Indonesia. Proceedings of the  
1768 International Workshop on Human Dimension of Tropical Peatland under Global Environmental Changes,  
1769 Bogor, Indonesia, December 8 - 9, 2004, pp. 112-119
- 1770 Davidson EA 1991. Fluxes of nitrous oxide and nitric oxide from terrestrial ecosystems. In *Microbial Production*  
1771 *and Consumption of Greenhouse Gases: Methane, Nitrogen Oxides and Halomethanes*. Eds JE Roggers and  
1772 WB Whitman, pp. 219–235. American Society for Microbiology, Washington.
- 1773 Dawson, J.J.C., Bakewell, T., Billett, M.F. 2001. Is in-stream processing an important control on spatial changes  
1774 in carbon fluxes in headwater catchments? *Sci. Total Environ.* 265: 153-167.
- 1775 Dawson, J.J.C., Billett, M.F., Hope, D., Palmer, S.M., Deacon, C.M. 2004. Sources and sinks of aquatic carbon  
1776 in a peatland stream continuum. *Biogeochemistry* 70: 71–92.
- 1777 de Groot, W. J. and M. E. Alexander (1986). Wildfire behavior on the Canadian Shield; case study of the 1980  
1778 Chachukew Fire, east-central Saskatchewan. Proc. Third Central Region Fire Weather Committee Sci. and  
1779 Tech. Seminar, Winnipeg, Manitoba, Can. For. Serv., West. & North. Reg., North. For. Cent., Edmonton.
- 1780 Dechow R, Freibauer A (2011) Assessment of German nitrous oxide emissions using empirical modelling  
1781 approaches. *Nutr Cycl Agroecosystems* 91(3):235-254.
- 1782 Department of Irrigation and Drainage and Land and Water Research Group (DID & LAWOO). 1996. Western  
1783 Jalore Integrated Agricultural Development Project. Peat Soil Management Study. Final report, Wageningen,  
1784 The Netherlands. ISN 16849. Pp 171.
- 1785 Deverel, S.J. Leighton, D.A. 2010. Historic, recent, and future subsidence, Sacramento-San Joaquin Delta,  
1786 California. USA. *San Francisco Estuary and Watershed Science* 8.
- 1787 di Folco, M-B., Kirkpatrick, J.B. 2011. Topographic variation in burning-induced loss of carbon from organic  
1788 soils in Tasmanian moorlands. *Catena* 87: 216-255.
- 1789 Dinsmore, K.J., Billett, M.F., Moore, T.R. (2009) Transfer of carbon dioxide and methane through the soil-  
1790 water-atmosphere system at Mer Bleue peatland, Canada. *Hydrol Process* 23, 330-341. Dinsmore, K.J.,  
1791 Billett, M.F., Skiba, U.M., Rees, R.M., Drewer, J., Helfter, C. 2010. Role of the aquatic pathway in the  
1792 carbon and greenhouse gas budgets of a peatland catchment. *Global Change Biol.* 16: 2750-2762.
- 1793 Dobbie KE, McTaggart IP, Smith KA 1999. Nitrous oxide emissions from intensive agricultural systems:  
1794 variations between crops and seasons, key driving variables, and mean emission factors. *J. Geophys. Res.*,  
1795 104, 26891– 26899.

Final Draft

- 1796 Drösler, M., 2005. Trace gas exchange and climatic relevance of bog ecosystems, Southern Germany.  
1797 Technische Universität München, Freising. Online published at: [http://nbn-](http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:bvb:91-diss20050901-1249431017)  
1798 [resolving.de/urn/resolver.pl?urn:nbn:de:bvb:91-diss20050901-1249431017](http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:bvb:91-diss20050901-1249431017)
- 1799 Drösler, M., Adelman, W., Augustin, J., Bergman, L., Beyer, C., Chojnicki, B., Förster, Ch., Freibauer, A.,  
1800 Giebels, M., Görlitz, S., Höper, H., Kantelhardt, J., Liebersbach, H., Hahn-Schöfl, M., Minke, M., Petschow,  
1801 U., Pfadenhauer, J., Schaller, L., Schägner, Ph., Sommer, M., Thuille, A., Wehrhan, M. 2013. Klimaschutz  
1802 durch Moorschutz. Schlussbericht des BMBF-Vorhabens: Klimaschutz - Moornutzungsstrategien 2006-2010.  
1803 201 pp. published online at TIB/UB-Hannover: [http://edok01.tib.uni-](http://edok01.tib.uni-hannover.de/edoks/e01fb13/735500762.pdf)  
1804 [hannover.de/edoks/e01fb13/735500762.pdf](http://edok01.tib.uni-hannover.de/edoks/e01fb13/735500762.pdf)
- 1805 Drösler, M., Schaller, L., Kantelhardt, J., Schweiger, M., Fuchs, D., Tiemeyer, B., Augustin, J., Wehrhan, M.,  
1806 Förster, Ch., Bergmann, L., Kapfer A., Krüger G.-M. 2012. Beitrag von Moorschutz- und -  
1807 revitalisierungsmaßnahmen zum Klimaschutz am Beispiel von Naturschutzgroßprojekten. *Natur und*  
1808 *Landschaft*, 87, Heft 02, pp 70-76.
- 1809 Dutaur, L. and Verhot, L.V. 2007. A global inventory of the soil CH<sub>4</sub> sink, *Global Biogeochem. Cy.*, 21,  
1810 GB4013,doi:10.1029/2006GB002734, 2007.
- 1811 Eggelsmann R, Bartels R. 1975. Oxidativer Torfverzehr im Niedermoor in Abhängigkeit von Entwässerung,  
1812 Nutzung und Düngung. *Mitteilungen der Deutschen Bodenkundlichen Gesellschaft* 22: 215-221.
- 1813 Elsgaard, L., Gorres, C.-M., Hoffmann, C.C., Blicher-Mathiesen, G. Schelde, K., Petersen, S.O. 2012. Net  
1814 ecosystem exchange of CO<sub>2</sub> and carbon balance for eight temperate organic soils under agricultural  
1815 management. *Agriculture, Ecosystems and Environment* 162: 52-67.
- 1816 Epting, J., D. Verbyla and B. Sorbel (2005). Evaluation of remotely sensed indices for assessing burn severity in  
1817 interior Alaska using Landsat TM and ETM+. *Remote Sensing of Environment* 96: 328-339.
- 1818 Fiedler, S., Höll, B.S., Freibauer, A., Stahr, K., Drösler, M., Schloter, M., Jungkunst, H.F. 2008. Particulate  
1819 organic carbon (POC) in relation to other pore water carbon fractions in drained and rewetted fens in  
1820 Southern Germany. *Biogeosciences*, 5: 1615–1623.
- 1821 Finér L., Ohashi, M., Noguchi, K., Hirano, Y. 2011. Fine root production and turnover in forest ecosystems in  
1822 relation to stand and environmental characteristics. *For. Ecol. Manage.*, 262: 2008–2023
- 1823 Firestone MK, Davidson EA 1989. Microbiological basis of NO and N<sub>2</sub>O production and consumption in soil.  
1824 *In*, Exchange of Trace Gases between Terrestrial Ecosystems and the Atmosphere. Eds MO Andreae and DS  
1825 Schimel, 7–21. John Wiley, New York.
- 1826 Flessa, H., Wild, U., Klemisch, M. and Pfadenhauer, J. 1998. Nitrous oxide and methane fluxes from organic  
1827 soils under agriculture. *Eur. J. of Soil Scien.*, 49: 327-335.
- 1828 French, N. H. F., P. Goovaerts and E. S. Kasischke (2004). Uncertainty in estimating carbon emissions from  
1829 boreal forest fires. *Journal of Geophysical Research* 109, D14, 27, DOI: 10.1029/2003JD003635
- 1830 Furukawa Y, Inubushi K, Ali M, Itang AM, Tsuruta H 2005. Effect of changing groundwater levels caused by  
1831 land-use changes on greenhouse gas emissions from tropical peatlands. *Nutr. Cycl. Agroecosyst.*, 71, 81–91.
- 1832 Gaudinski, J., Torn, M., Riley, W., Dawson, T., Doslin, D. and Majdi, H.2010. Measuring and modelling the  
1833 spectrum of fine-root turnover times in three forests using isotopes, minirhizotrons, and the Radix model.  
1834 *Glob. Biogeochem. Cycles*, 24 DOI 10.1029/2009/GB003649.
- 1835 Gebhart, K. A., S. M. Kreidenweis and W. C. Malm (2001). Back-trajectory analyses of fine particulate matter  
1836 measured at Big Bend National Park in the historical database and the 1996 scoping study. *Science of the*  
1837 *Total Environment* 36: 185-204.
- 1838 Giglio, L., T. Loboda, D. P. Roy, B. Quayle and C. O. Justice (2009). An active-fire based burned area mapping  
1839 algorithm for the MODIS sensor. *Remote Sensing of Environment* 113: 408-420.
- 1840 Gill, R. A., and R. B. Jackson (2000), Global patterns of root turnover for terrestrial ecosystems, *New Phytol.*,  
1841 147, 13–31, doi:10.1046/j.1469-8137.2000.00681.x.
- 1842 Glagolev, M.V., Chistotin, M.V., Shnyrev, N.A., Sirin, A.A. 2008. The emission of carbon dioxide and methane  
1843 from drained peatlands changed by economic use and from natural mires during the summer-fall period (on  
1844 example of a region of Tomsk oblast). *Agrokhimija* 5: 46-58.
- 1845 Glatzel, S., Kalbitz, K., Dalva, M. Moore, T. 2003. Dissolved organic matter properties and their relationship to  
1846 carbon dioxide efflux from restored peat bogs. *Geoderma* 113: 397-411.

Final Draft

- 1847 Glenn S., Heyes A., Moore T. 1993. Carbon dioxide and methane fluxes from drained peat soils, southern  
1848 Quebec. *Global Biogeochem. Cycles* 7: 247-257.
- 1849 Gorham, E. (1991). Northern peatlands: role in the carbon cycle and probable responses to climatic warming.  
1850 *Ecological Applications* 1: 182-195.
- 1851 Gorham, E. 1991) Northern Peatlands: Role in the carbon cycle and probable responses to climatic warming,  
1852 *Ecol. Appl.*, 1: 182-195.
- 1853 Goulsbra, C. Evans, M., Allott, T. 2013. 5. Towards the estimation of CO<sub>2</sub> emissions associated with POC  
1854 fluxes from drained and eroding peatlands. In: Emissions of greenhouse gases associated with peatland  
1855 drainage waters: Report to Defra under project SP1205: Greenhouse Gas Emissions Associated with Non  
1856 Gaseous Losses of Carbon from Peatlands - fate of Particulate and Dissolved Carbon. Report to the  
1857 Department of Environment, Food and Rural Affairs, UK.
- 1858 Gregoire, J.-M., K. J. Tansey and J. M. N. Silva (2003). The GBA2000 initiative: Developing a global burnt area  
1859 database from SPOT-VEGETATION imagery. *International Journal of Remote Sensing* 24: 1369-1376.
- 1860 Grønlund, A., Hauge, A., Hovde, A., Rasse, D. P., 2008: Carbon loss estimates from cultivated peat soils in  
1861 Norway: a comparison of three methods. *Nutr. Cycl. Agroecosystems* 81: 157-167.
- 1862 Grønlund, A., Sveistrup, T. E., Søvik, A. K., Rasse, D. P. and Kløve, B. 2006. Degradation of cultivated peat  
1863 soils in Norway based on field scale CO<sub>2</sub>, N<sub>2</sub>O and CH<sub>4</sub> emission measurements, *Arch. Agron. Soil Sci.*, 52,  
1864 149–159.
- 1865 Guðmundsson, J. & Óskarsson, H. 2008. Summaries of GHG measurement studies. UNESCO/ IHA Greenhouse  
1866 Gas Research Project. Measurement Specification Workshop, London, UK, 12–14 Nov.
- 1867 Hadi, A., Inubushi, K., Furukawa, Y., Purnomo, E., Rasmadi, M., Tsuruta, H. (2005) Greenhouse gas emissions  
1868 from tropical peatlands of Kalimantan, Indonesia. *Nutrient Cycling in Agroecosystems* 71, 73-80.
- 1869 Hairiah, K., M. van Noordwijk, and G. Cadisch (1999), Roots as part of the carbon and nitrogen input and output  
1870 of three types of cropping systems on an Ultisol in North Lampung, in *Proceedings of the Seminar Toward  
1871 Sustainable Agriculture in Humid Tropics Facing 21st Century*, Bandar Lampung, Indonesia, 27–28  
1872 September 1999, edited by C. Ginting et al., pp. 86–95, Int. Cent. for Res. in Agrofor., Bogor, Indonesia.
- 1873 Hairiah, K., M. van Noordwijk, and G. Cadisch (2000), Crop yield, C and N balance of three types of cropping  
1874 systems on an Ultisol in Northern Lampung, *Neth. J. Agric. Sci.*, 48, 3–17.
- 1875 Hall, R. J., J. T. Freeburn, W. J. De Groot, J. M. Pritchard, T. J. Lynham and R. Landry (2008). Remote sensing  
1876 of burn severity: experience from western Canada boreal fires. *International Journal of Wildland Fire* 17:  
1877 476-489.
- 1878 Hamada, Y., Darung, U., Limin, S.H. and Hatano, R. 2013 Characteristics of fire-generated gas emission  
1879 observed during a large peatland fire in 2009 at Kalimantan, Indonesia. *Atmospheric Environment*, 74, 177-  
1880 181.
- 1881 Hargreaves, K.J., Milne, R., Cannell, M.G.R. 2003. Carbon balance of afforested peatland in Scotland, *Forestry*,  
1882 76, 299-317.
- 1883 Harrison, M.E., Cheyne, S.M., Sulistiyanto, Y., Rieley, J.O., 2007 Biological effects of smoke from dry-season  
1884 fires in non-burnt areas of the Sabangau peat swamp forest, Central Kalimantan, Indonesia. Paper presented  
1885 at International Symposium and Workshop Carbon-Climate-Human Interactions: Carbon Pools, Fire,  
1886 Mitigation, Restoration and Wise Use, Yogyakarta, Indonesia, 27–31 August. (Available at [http://](http://www.geog.le.ac.uk/carbopeat/media/pdf/yogyapapers/p9.pdf)  
1887 [www.geog.le.ac.uk/carbopeat/media/pdf/yogyapapers/p9.pdf](http://www.geog.le.ac.uk/carbopeat/media/pdf/yogyapapers/p9.pdf))
- 1888 Heikkinen, K., 1990. Transport of organic and inorganic matter in river, brook and peat mining water in the  
1889 drainage basin of the River Kiiminkijoki. *Aqua Fennica*, 20: 143-155.
- 1890 Heil, A., Langmann, B. and Aldrian, E. 2006. Indonesian peat and vegetation fire emissions: Study on factors  
1891 influencing large-scale smoke haze pollution using a regional atmospheric chemistry model. *Mitigation and  
1892 Adaptations Strategies for Global Change* 12: 113-133.
- 1893 Hendriks, D.M.D., Van Huissteden, J., Dolman, A.J. 2010. Multi-technique assessment of spatial and temporal  
1894 variability of methane fluxes in a peat meadow. *Agricultural and Forest Meteorology* 150: 757-774
- 1895 Hendriks, D.M.D., Van Huissteden, J., Dolman, A.J., Van der Molen, M.K. 2007 The full greenhouse gas  
1896 balance of an abandoned peat meadow. *Biogeosciences* 4:411-424.

Final Draft

- 1897 Henson, I. E., and M. T. Dolmat (2003), Physiological analysis of an oil palm density trial on a peat soil, *J. Oil*  
1898 *Palm Res.*, 15, 1–27.
- 1899 Hergoualc'h, K., Verchot, L.V., 2011. Stocks and fluxes of carbon associated with land-use change in Southeast  
1900 Asian tropical peatlands: a review. *Global Biochem. Cycles* 25, GB2001, doi:2010.1029/2009GB003718.
- 1901 Hergoualc'h, K., Verchot, L.V., 2012. Changes in soil CH<sub>4</sub> fluxes from the conversion of tropical peat swamp  
1902 forests: a meta-analysis. *Journal of Integrative Environmental Sciences* 9, 93–101.
- 1903 Hergoualc'h, K., Verchot, L.V., 2013. Greenhouse gas emission factors for land use and land-use change in  
1904 Southeast Asian peatlands. Submitted to *Mitigation and Adaptation Strategies for Global Change*.
- 1905 Hillebrand, K., 1993. The greenhouse effects of peat production and use compared with coal, oil, natural gas and  
1906 wood. VTT Tiedotteita - Meddelanden - Research Notes 1494, Technical Research Centre of Finland, Espoo.
- 1907 Hirano, T, Segah, H., Harada, T., Limin, S., June, T., Hirata, R., Osaki, M. 2007. Carbon dioxide balance of a  
1908 tropical peat swamp forest in Kalimantan, Indonesia. *Global Change Biol.* 13: 412–425.
- 1909 Hirano, T., Jauhiainen, J., Inoue, T., and Takahashi, H. 2009. Controls on the carbon balance of tropical  
1910 peatlands, *Ecosystems*, 12, 873–887
- 1911 Hirano, T., Segah, H., Kusin, K., Limin, S., Takahashi, H., Osaki, M. (2012) Effects of disturbances on the  
1912 carbon balance of tropical peat swamp forests. *Global Change Biology* 18, 3410-3422.
- 1913 Holden, J., Chapman, P.J., Palmer, S.M., Kay, P., Grayson, R., 2012. The impacts of prescribed moorland  
1914 burning on water colour and dissolved organic carbon: A critical synthesis. *Journal of Environmental*  
1915 *Management* 101, 92-103.
- 1916 Honrath, R. E., R. C. Owen, M. V. Martin, J. S. Reid, K. Lapina, P. Fialho, M. P. Dziobak, J. Kleissl and D. L.  
1917 Westphal (2004). Regional and hemispheric impacts of anthropogenic and biomass burning emissions on  
1918 summertime CO and O<sub>3</sub> in the North Atlantic lower free troposphere. *Journal of Geophysical Research* 109:  
1919 D24310, doi:24310.21029/22004JD005147.
- 1920 Hooijer, A., Page, S., Jauhiainen, J., Lee, W.A., Lu, X.X., Idris, A., Anshari, G., 2012. Subsidence and carbon  
1921 loss in drained tropical peatlands. *Biogeosciences* 9, 1053-1071.
- 1922 Hope, D., Palmer, S.M., Billett, M.F., Dawson, J.J.C. (2001) Carbon dioxide and methane evasion from a  
1923 temperate Peatland stream. *Limnol. Oceanogr.* 46, 847-857.
- 1924 Höper, H. 2002. Carbon and nitrogen mineralization rates in German agriculturally used fenlands. 149-164. In:  
1925 Broll, G. Merbach, W. and E.-M. Pfeiffer (Eds.). *Wetlands in Central Europe. Soil organisms, soil ecological*  
1926 *processes, and trace gas emissions.* Springer, Berlin. 244 p.
- 1927 Hoscilo, A., K. J. Tansey and S. E. Page (2013). Post-fire vegetation response as a proxy to quantify the  
1928 magnitude of burn severity in tropical peatland. *International Journal of Remote Sensing* 34: 412-433.
- 1929 Huttunen J.T., Nykänen H., Turunen J. & Martikainen P.J. 2003a. Methane emissions from natural peatlands in  
1930 the northern boreal zone in Finland, Fennoscandia. *Atmos. Environ.* 37: 147–151.
- 1931 Hyvönen, N. P., Huttunen, J. T., Shurpali, N. J., Lind, S.E., Marushchak, M.E., Heitto, L., and Martikainen, P. J.  
1932 20132012. The role of drainage ditches in greenhouse gas emissions and surfave leaching losses from a  
1933 cutaway peatland cultivated with a perennial bioenergy crop, *Boreal Env. Res.* 18., 109-126.
- 1934 Hyvönen, N. P., Huttunen, J. T., Shurpali, N. J., Tavi, N. M., Repo, M. E. and Martikainen, P. J. 2009. Fluxes of  
1935 N<sub>2</sub>O and CH<sub>4</sub> on an organic soil: Effect of bioenergy crop cultivation, *Biores. Techn.*,  
1936 doi:10.1016/j.biortech.2009.04.043.
- 1937 Immirzi, C. P., E. Maltby and R. S. Clymo (1992). The global status of peatlands and their role in carbon cycling.  
1938 London, Wetland Ecosystems Research Group, Dept. Geography, University of Exeter: 1-145.
- 1939 Inubushi K, Furukawa Y, Hadi A, Purnomo E, Tsuruta H 2003. Seasonal changes of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O fluxes  
1940 in relation to land-use change in tropical peatlands located in coastal area of South Kalimantan. *Chemosphere*,  
1941 52, 603–608.
- 1942 Inubushi, K., Hadi, A., Okazaki, M., Yonebayashi, K. 1998. Effect of converting wetland forest to sago palm  
1943 plantations on methane gas flux and organic carbon dynamics in tropical peat soil. *Hydrol. Process.* 12: 2073-  
1944 2080.
- 1945 IPCC 2010, Use of Models and Facility-Level Data in Greenhouse Gas Inventories (Report of IPCC Expert  
1946 Meeting on Use of Models and Measurements in Greenhouse Gas Inventories 9-11 August 2010, Sydney,

Final Draft

- 1947 Australia) eds: Eggleston H.S., Srivastava N., Tanabe K., Baasansuren J., Fukuda M., GHG Pub. IGES, Japan  
1948 2010.
- 1949 Ishida, T., Suzuki, S., Nagano, T., Osawa, K., Yoshino, K., Fukumura, K., Nuyim, T. 2001. CO<sub>2</sub> emission rate  
1950 from a primary peat swamp forest ecosystem in Thailand. *Environ Control Biol* 39(4): 305–12.
- 1951 Ishizuka, S., Iswandi, A., Nakajima, Y., Yonemura, S., Sudo, S., Tsuruta H., and Murdiyarso, D., 2005. The  
1952 variation of greenhouse gas emissions from soils of various land-use/cover types in Jambi province,  
1953 Indonesia. *Nutrient Cycling in Agroecosystems* 71: 17–32, DOI 10.1007/s10705-004-0382.
- 1954 Itkonen, A. and M. J. Jantunen (1986). Emissions and particle-size distribution of some metallic elements of two  
1955 peat/oil-fired boilers. *Environmental Science and Technology* 20: 335-341.
- 1956 Jaakkola, A. 1985. Lannoite ja kasviainestysten hyväksikäyttö ja häviö. Biologisen typensidonnan ja  
1957 ravinnetyypen hyväksikäytön projekti. Suomen itsenäisyyden juhluvuoden 1967 rahasto. Julkaisu 13. Helsinki.  
1958 107 pp. (in Finnish).
- 1959 Jager, D.F., Wilmking, M., Kukkonen, J.V.K. 2009. The influence of summer seasonal extremes on dissolved  
1960 organic carbon export from a boreal peatland catchment: Evidence from one dry and one wet growing season.  
1961 *Sci. Total Environ.* 407: 1373-1382.
- 1962 Jauhiainen, J., Hooijer, A., Page, S.E., 2012a. Carbon dioxide emissions from an Acacia plantation on peatland  
1963 in Sumatra, Indonesia. *Biogeosciences* 9, 617-630.
- 1964 Jauhiainen, J., Silvennoinen, H., Hämäläinen, R., Kusin, K., Limin, S., Raison, R.J., Vasander, H., 2012b.  
1965 Nitrous oxide fluxes from tropical peat with different disturbance history and management. *Biogeosciences* 9,  
1966 1337-1350.
- 1967 Jauhiainen, J., Hooijer, A., Page, S.E., 2012c. Greenhouse gas emissions from a plantation on thick tropical peat.  
1968 14th International peat congress, Stockholm, Sweden, 3-8 June 2012. Jauhiainen, J., Limin, S., Silvennoinen,  
1969 H., Vasander, H. (2008) Carbon dioxide and methane fluxes in drained tropical peat before and after  
1970 hydrological restoration. *Ecology* 89, 3503-3514.
- 1971 Jauhiainen, J., Silvennoinen, H. 2012. Diffusion GHG fluxes at tropical peatland drainage canal water surfaces.  
1972 *Suo* 63, 93-105.
- 1973 Johnson, E. A. (1992). *Fire and vegetation dynamics: studies from the North American boreal forest*. Cambridge,  
1974 UK, Cambridge University Press.
- 1975 Jones, T., Jones, D., Evans, C. 2013. Conversion of waterborne DOC to CO<sub>2</sub> – results of laboratory experiments.  
1976 In: *Emissions of greenhouse gases associated with peatland drainage waters: Report to Defra under project*  
1977 *SP1205: Greenhouse Gas Emissions Associated with Non Gaseous Losses of Carbon from Peatlands - fate of*  
1978 *Particulate and Dissolved Carbon*. Report to the Department of Environment, Food and Rural Affairs, UK.
- 1979 Jonsson, A. Algesten, G., Bergström, A-K, Bishop, K., Sobek, S., Tranvik, L.J., Jansson, M. (2009). Integrating  
1980 aquatic carbon fluxes in a boreal catchment carbon budget. *J. Hydrol.* 334: 141-150.
- 1981 Juutinen, S., Välijanta, M., Kuutti, V., Laine, A.M., Virtanen, T., Seppä, H., Weckström, J., Tuittila, E-S. (2013).  
1982 Short-term and long-term carbon dynamics in a northern peatland-stream-lake continuum: A catchment  
1983 approach. *Journal of Geophysical Research: Biogeosciences*, 118, 171-183.
- 1984 Kajii, Y., S. Kato, D. Streets, N. Tsai, A. Shvidenko, S. Nilsson, I. McCallum, N. Minko, N. Abushenko, D.  
1985 Altyntsev and T. Khodzer (2002). Boreal forest fires in Siberia in 1998: estimation of area burned and  
1986 emissions of pollutants by advanced very high resolution radiometer satellite data. *Journal of Geophysical*  
1987 *Research* 107. D24, ACH 4-1–ACH 4-8, 27, DOI: 10.1029/2001JD001078.
- 1988 Kakuda, K-I; Watanabe, A; Ando, H; Jong, FS (2005) Effects of Fertilizer Application on the Root and  
1989 Aboveground Biomass of Sago Palm (*Metroxylon sago* Rottb.) Cultivated in Peat Soil. *Jpn. J. Trop. Agr.*  
1990 49(4) : 264 – 269.
- 1991 Kane, E.S., Turetsky, M.R., Harden, J.W., McGuire, A.D., Waddington, J.M. Seasonal ice and hydrologic  
1992 controls on dissolved organic carbon and nitrogen concentrations in a boreal-rich fen. *J. Geophys. Res.* 115,  
1993 G04012. doi:10.1029/2010JG001366.
- 1994 Kasimir-Klemedtsson Å., Klemedtsson L., Berglund K., Martikainen P., Silvola J., Oenema O. 1997.  
1995 Greenhouse gas emissions from farmed organic soils: a review. *Soil Use and Manag.*, 13: 245-250.
- 1996 Kasimir-Klemedtsson, Å., Weslien, P. and Klemedtsson, L. (2009). Methane and nitrous oxide fluxes from a  
1997 farmed Swedish Histosol, *Eur. J. Soil Sci.*, 60, 321–331, doi:10.1111/j.1365- 2389.2009.01124.x.

Final Draft

- 1998 Kasischke, E. S. 2000. Boreal ecosystems in the carbon cycle. *Fire, Climate Change and Carbon Cycling in the*  
1999 *North American Boreal Forest*. E. S. Kasischke and B. J. Stocks. New York, Springer-Verlag: 19-30.
- 2000 Kasischke, E. S. and Bruhwiler, L. P. (2003). Emissions of carbon dioxide, carbon monoxide and methane from  
2001 boreal forest fires in 1998. *Journal of Geophysical Research* 108: 8146, doi:8110.1029/2001JD000461.
- 2002 Kasischke, E. S., Hyer, E. J. Novelli, P. C. Bruhwiler, L. P. French, N. H. F. Sukhinin, A. I. Hewson J. H. and  
2003 Stocks, B. J. (2005). Influences of boreal fire emissions on Northern Hemisphere atmospheric carbon and  
2004 carbon monoxide. *Global Biogeochemical Cycles* **19**: GB1012, doi:10.1029/2004GB002300.
- 2005 Kasischke, E. S., Bourgeau-Chavez, L. L. Rober, A. R. Wyatt, K. H. Waddington J. M. and Turetsky M. R.  
2006 (2009). Effects of soil moisture and water depth on ERS SAR backscatter measurements from an  
2007 Alaskan wetland complex. *Remote Sensing of Environment* **113**: 1868-1873.
- 2008 Kasischke, E. S., Christensen, N. L. and Stocks, B. J. (1995). Fire, global warming, and the carbon balance of  
2009 boreal forests. *Ecological Applications* 5: 437-451.
- 2010 Kasischke, E. S., Loboda, T., Giglio, L., French, N. H. F. Hoy, E. E. de Jong, B. and Riaño, D. (2011).  
2011 Quantifying burned area from fires in North American forests: Implications for direct reduction of carbon  
2012 stocks. *Journal of Geophysical Research* 116: doi:10.1029/2011JG001707.
- 2013 Keeney D.R., Fillery I.R., Marx G.P. 1979. Effect of temperature on the gaseous nitrogen products of  
2014 denitrification in a silty loam soil. *Soil Sci. Soc. Am. J* 43, 1124–1128.
- 2015 Klemetsson, L., von Arnold, K., Weslien, P. and Gundersen, P. (2005). Soils CN ratio as scalar parameter to  
2016 predict nitrous oxide emissions. *Global Change Biology* 11:1142–1147.
- 2017 Koehler, A-K., Murphy, K., Kiely, G., Sottocornola, M. (2009). Seasonal variation of DOC concentration and  
2018 annual loss of DOC from an Atlantic blanket bog in South Western Ireland. *Biogeochemistry* 95: 231–242.
- 2019 Koehler, A-K., Sottocornola, M., Kiely, G. (2011). How strong is the current carbon sequestration of an Atlantic  
2020 blanket bog? *Global Change Biol.* 17: 309–319.
- 2021 Köhler, S., Buffam, I., Jonsson, A., Bishop, K. (2002). Photochemical and microbial processing of stream and  
2022 soil water dissolved organic matter in a boreal forested catchment in northern Sweden. *Aquat. Sci.* 64, 1-13.
- 2023 Kolka, R.K., Grigal, D.F., Verry, E.S., Nater, E.A. (1999). Mercury and organic carbon relationships in streams  
2024 draining forested upland peatland watersheds. *J. Environmental Quality* 28: 766-775.
- 2025 Komulainen, V.-M., Nykänen, H., Martikainen, P. J. and Laine, J. (1998). Short-term effect of restoration on  
2026 vegetation change and methane emissions from peatlands drained for forestry in southern Finland, *Can. J.*  
2027 *For. Res.*, 28, 402–411.
- 2028 Komulainen, V.-M., Tuittila, E. S., Vasander, H. and Laine, J. (1999). Restoration of drained peatlands in  
2029 southern Finland: initial effects on vegetation change and CO<sub>2</sub> balance, *J. Appl. Ecol.*, 36, 634–648.
- 2030 Koprivnjak, J-F, Moore, T.R. 1992. Sources, sinks and fluxes of dissolved organic carbon in subarctic fen  
2031 catchments. *Arctic and Alpine Research*, 24: 204-210.
- 2032 Kortelainen, P., Mattsson, T., Finér, L., Ahtiainen, M., Saukkonen, S., Sallantausta, T. (2006). Controls on the  
2033 export of C, N, P and Fe from undisturbed boreal catchments, Finland. *Aquat. Sci.* 68: 453-468.
- 2034 Kreshtapova V.N., Maslov B.S. 2004. Contents of carbon compounds in reclaimed peat soils as a function of the  
2035 properties of peat organic matter. *Proc of 12<sup>th</sup> Peat Cong., Tampere*, volume 2: 988-992.
- 2036 Kroon, P. S., Schrier-Uijl, A. P., Hensen, A., Veenendaal, E. M., Jonker, H. J. J. (2010). Annual balances of  
2037 CH<sub>4</sub> and N<sub>2</sub>O from a managed fen meadow using eddy covariance flux measurements. *European Journal of*  
2038 *Soil Science* 61:773-784, 10.1111/j.1365-2389.2010.01273.x.
- 2039 Kuhry, P. (1994). The role of fire in the development of Sphagnum-dominated peatlands in western boreal  
2040 Canada. *Journal of Ecology* 82: 899-910.
- 2041 Kuntze, H., (1992). Peat losses by liming and fertilization of peatlands used as Grassland. *Proc 9th Int Peat*  
2042 *Congress*, vol. 2: 306–314.
- 2043 Laine, J., Minkinen, K., Sinisalo, J., Savolainen, I. and Martikainen, P. J. 1996. Greenhouse Impact of a mire  
2044 after drainage for forestry, in: *Northern Forested Wetlands, Ecology and Management*, edited by: Trettin, C.  
2045 C., Jurgensen, M. F., Grigal, D. F., Gale, M. R. and Jeglum, J. K., CRC Lewis Publishers Boca Raton, USA,  
2046 437–447.

Final Draft

- 2047 Lamade, E., and J-P. Bouillet (2005), Carbon storage and global change: the role of oil palm, *OCL - Oléagineux,*  
2048 *Corps Gras, Lipides*, 12, 154-160.
- 2049 Langeveld C.A., Segers R., Dirks B.O.M., van den Pol-van Dasselaar A., Velthof G.L., Hensen A. 1997.  
2050 Emissions of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O from pasture on drained peat soils in the Netherlands. *Europ. J. Agr.* 7: 35-  
2051 42.
- 2052 Lapina, K., R. E. Honrath, R. C. Owen, M. Val Martin, E. J. Hyer and P. Fialho (2008). Late-summer changes in  
2053 burning conditions in the boreal regions and thier implications for NO<sub>x</sub> and CO emissions from boreal fires.  
2054 *Journal of Geophysical Research* 113: D11304, doi:11310.11029/12007JD009421.
- 2055 Laurila, T., Lohila, A., Aurela, M., Tuovinen, J.-P., Thum, T., Aro, L., Laine, J., Penttilä, T., Minkkinen, K.,  
2056 Riutta, T., Rinne, J., Pihlatie, M. and Vesala, T. 2007. Ecosystem-level carbon sink measurements on forested  
2057 peatlands, in: *Greenhouse Impacts of the Use of Peat and Peatlands in Finland*, edited by: Sarkkola, S.,  
2058 Ministry of Agriculture and Forestry 11a/2007, 38–40.
- 2059 Leifeld J., Müller M. & Fuhrer J., 2011. Peatland subsidence and carbon loss from drained temperate fens. *Soil*  
2060 *Use and Management*, June 2011, 27, 170-176.
- 2061 Lindroth, A., Klemetsson, L., Grelle, A., Weslien, P. and Langvall O. 2007. Measurement of net ecosystem  
2062 exchange, productivity and respiration in three spruce forests in Sweden shows unexpectedly large soil  
2063 carbon losses. *Biogeochemistry* 89(1): 43–60. DOI 10.1007/s10533-007-9137-8.
- 2064 Lloyd, C. R., (2006). Annual carbon balance of a managed wetland meadow in the Somerset Levels, UK.  
2065 *Agricultural and Forest Meteorology* 138 :168–179. Lohila, A., Aurela, M., Tuovinen, J.-P. and Laurila, T.  
2066 2004. Annual CO<sub>2</sub> exchange of a peat field growing spring barley or perennial forage, *J. Geophys. Res.*, 109,  
2067 D18116, doi:10.1029/2004JD004715.
- 2068 Lohila, A., Laurila, T., Aro, L., Aurela, M., Tuovinen, J.-P., Laine, J. and Minkkinen, K. 2007. Carbon dioxide  
2069 exchange above a 30-year-old Scots pine plantation established on organic-soil Cropland, *Boreal Environ.*  
2070 *Res.*, 12, 141–157.
- 2071 Lohila, A, Minkkinen, K, Aurela, M, Tuovinen, J-P, Penttilä, T and Laurila, T. Lohila, A, Minkkinen, K, Aurela,  
2072 M, Tuovinen, J-P, Penttilä, T and Laurila, T. 2011. Greenhouse gas flux measurements in a forestry-drained  
2073 peatland indicate a large carbon sink. *Biogeosciences Discuss*, 8: 5787–5825.
- 2074 Lorenz, W. D., Sauerbrey, R., Eschner, D., Lehrkamp, H. Zeitz, J. 1992. Zustand der landwirtschaftlich  
2075 genutzten Niedermoore in der ehemaligen DDR, *Wasser und Boden*, 44, 58-61
- 2076 Lucas, R. E. (1982). *Organic soils (Histosols). Formation, distribution, physical and chemical properties and*  
2077 *management for crop production.* Michigan State University.
- 2078 Mäkiranta, P., Hytönen, J., Aro, L., Maljanen, M., Pihlatie, M., Potila, H., Shurpali, N. J., Laine, J., Lohila, A.,  
2079 Martikainen, P. J. and Minkkinen, K. 2007. Soil greenhouse gas emissions from afforested organic soil  
2080 Croplands and peat extraction peatlands, *Boreal Environ. Res.*, 12, 159–175.
- 2081 Malhi, Y., *et al.* 2009. Comprehensive assessment of carbon productivity, allocation and storage in three  
2082 Amazonian forests, *Global Change Biol.*, doi:10.1111/j.1365-2486.2008.01780.x.
- 2083 Maljanen, M. Hytönen, J. Martikainen, P.J. 2010. Cold-season nitrous oxide dynamics in a drained boreal  
2084 peatland differ depending on land-use practice. *Canadian Journal of Forest Research*, 40 (3): 565-572.
- 2085 Maljanen, M. Sigurdsson, B.D., Guðmundsson, J., Óskarsson, H., Huttunen, J.T., and Martikainen, P.J. 2010.  
2086 Greenhouse gas balances of managed peatlands in the Nordic countries – present knowledge and gaps.  
2087 *Biogeosciences*, 7, 2711–2738.
- 2088 Maljanen, M., Alm, J., Martikainen, P. J. and Repo, T. 2009a. Prolongation of soil frost resulting from reduced  
2089 snow cover increases nitrous oxide emissions from boreal forest soil, *Boreal Environ. Res.*, in press.
- 2090 Maljanen, M., Hytönen, J. and Martikainen, P. J. 2001b. Fluxes of N<sub>2</sub>O, CH<sub>4</sub> and CO<sub>2</sub> on afforested boreal  
2091 agricultural soils, *Plant Soil*, 231, 113–121.
- 2092 Maljanen, M., Hytönen, J., Mäkiranta, P., Alm, J., Minkkinen, K., Laine, J. and Martikainen, P. J. 2007a.  
2093 Greenhouse gas emissions from cultivated and abandoned organic Croplands in Finland, *Boreal Environ.*  
2094 *Res.*, 12, 133–140.
- 2095 Maljanen, M., Komulainen, V.-M., Hytönen, J., Martikainen, P. J. and Laine, J. 2004. Carbon dioxide, nitrous  
2096 oxide and methane dynamics in boreal organic agricultural soils with different soil management, *Soil Biol.*  
2097 *Biochem.*, 36, 1801–1808.

Final Draft

- 2098 Maljanen, M., Liikanen, A., Silvola, J. and Martikainen, P. J. 2003a. Methane fluxes on agricultural and forested  
2099 boreal organic soils, *Soil Use Manage.*, 19, 73–79.
- 2100 Maljanen, M., Liikanen, A., Silvola, J. and Martikainen, P. J. 2003b. Nitrous oxide emissions from boreal  
2101 organic soil under different land-use, *Soil Biol. Biochem.*, 35, 689–700.
- 2102 Maljanen, M., Martikainen, P. J., Walden, J. and Silvola, J. 2001a. CO<sub>2</sub> exchange in an organic field growing  
2103 barley or grass in eastern Finland, *Glob. Change Biol.*, 7, 679–692.
- 2104 Maljanen, M., Nykänen, H., Moilanen, M. and Martikainen, P. J. 2006a. Greenhouse gas fluxes of coniferous  
2105 forest floors affected by wood ash fertilization, *For. Ecol. Man.*, 237, 143–149.
- 2106 Maljanen, M., Virkajärvi, P., Hytönen, J., Öquist, M., Sparrman, T. and Martikainen, P. J. 2009b. Nitrous oxide  
2107 production in boreal soils with variable organic matter content at low temperature snow manipulation  
2108 experiment, *Biogeosciences Discuss.*, 6, 5305–5337, 2009, [http://www.biogeosciences-](http://www.biogeosciences-discuss.net/6/5305/2009/)  
2109 [discuss.net/6/5305/2009/](http://www.biogeosciences-discuss.net/6/5305/2009/).
- 2110 Marsden, K. and S. Ebmeier (2012). Peatlands and climate change. SPICe briefing: 35.
- 2111 Martikainen, P. J., Nykänen, H., Alm, J. and Silvola, J. 1995a. Change in fluxes of carbon dioxide, methane and  
2112 nitrous oxide due to forest drainage of mire sites of different trophy, *Plant Soil*, 168–169, 571–577.
- 2113 Martikainen, P. J., Nykänen, H., Crill, P. and Silvola, J. 1993. Effect of a lowered water table on nitrous oxide  
2114 fluxes from northern peatlands, *Nature*, 366, 51–53.
- 2115 Martikainen, P. J., Nykänen, H., Crill, P. and Silvola, J. 1992. The effect of changing water table on methane  
2116 fluxes at two Finnish mire sites, *Suo*, 43, 237–240.
- 2117 Martikainen, P. J., Nykänen, H., Regina, K., Lehtonen, M. and Silvola, J. 1995b. Methane fluxes in a drained and  
2118 forested peatland treated with different nitrogen compounds, in: *Northern Peatlands in Global Climatic*  
2119 *Change*, edited by: Laiho, R., Laine, J. and Vasander, H. Proceedings of the International Workshop Held in  
2120 Hyytiälä, Finland, Helsinki, 105–109.
- 2121 Marwanto, S. and Agus, F. 2013. Is CO<sub>2</sub> flux from oil palm plantations on peatland controlled by water table,  
2122 soil moisture, day/night rhythm and/or temperature? *Mitigation and Adaptation Strategies for Global Change*.  
2123 Accepted.
- 2124 Matthews, R. B., R. Wassmann, L. V. Buendia, and J. W. Knox (2000), Using a crop/soil simulation model and  
2125 GIS techniques to assess methane emissions from rice fields in Asia: II. Model validation and sensitivity  
2126 analysis, *Nutr. Cycl. Agroecosyst.*, 58, 161–177, doi:10.1023/A:1009846703516.
- 2127 McNamara, N. 2013. CH<sub>4</sub> emissions from ditches in a drained lowland peat Grassland, Somerset, UK. In:  
2128 Emissions of greenhouse gases associated with peatland drainage waters: Report to Defra under project  
2129 SP1205: Greenhouse Gas Emissions Associated with Non Gaseous Losses of Carbon from Peatlands - fate of  
2130 Particulate and Dissolved Carbon. Report to the Department of Environment, Food and Rural Affairs, UK.
- 2131 McNeil, P. and Waddington, J.M. 2003. Moisture controls on Sphagnum growth and CO<sub>2</sub> exchange on a cutover  
2132 bog. *Journal of Applied Ecology*, 40:354–367.
- 2133 Melling, L., Chaddy, A., Goh, K.J. and Hatano, R. 2013. Soil CO<sub>2</sub> Fluxes from Different Ages of Oil Palm in  
2134 Tropical Peatland of Sarawak, Malaysia as Influenced by Environmental and Soil Properties. *Acta Hort.*  
2135 982:25-35.
- 2136 Melling L, Hatano R, and Goh KJ 2007b. Nitrous oxide emissions from three ecosystems in tropical peatland of  
2137 Sarawak, Malaysia. *Soil Sci. Plant Nutr.* 53, 792–805
- 2138 Melling, L., Goh, K.J., Beauvais, C., Hatano, R., 2007a. Carbon flow and budget in a young mature oil palm  
2139 agroecosystem on deep tropical peat. International symposium and workshop on tropical peatland,  
2140 Yogyakarta, Indonesia, 27-31 August 2007.
- 2141 Melling, L., Hatano, R., Goh, K.J. (2005a) Soil CO<sub>2</sub> flux from three ecosystems in tropical peatland of Sarawak,  
2142 Malaysia. *Tellus* 57B, 1-11.
- 2143 Melling, L., Hatano, R., Goh, K.J., 2005b. Methane fluxes from three ecosystems in tropical peatland of Sarawak,  
2144 Malaysia. *Soil Biology & Biochemistry* 37, 1445–1453.
- 2145 Meyer, K. Höper H., Blankenburg J. 2001. Spurengashaushalt und Klimabilanz von Niedermooren unter dem  
2146 Einfluß des Vernässungsmanagements. In *Ökosystemmanagement für Niedermoore. Strategien und*  
2147 *Verfahren zur Renaturierung*. (Kratz R., Pfadenhauer J., eds) Ulmer, Stuttgart, 104-111.

Final Draft

- 2148 Milner, L., A. Boom, S. E. Page, S. Moore and R. Matthews (2013). Effects of fire on the organic matter  
2149 composition of a tropical peatland in Central Kalimantan, Indonesia. *Journal of Organic Geochemistry*.  
2150 Accepted.
- 2151 Minkkinen, K. & Laine, J. 2006. Vegetation heterogeneity and ditches create spatial variability in methane fluxes  
2152 from peatlands drained for forestry. *Plant and Soil*: 289–304.
- 2153 Minkkinen, K. and Laine, J. 1998. Long term effect of forest drainage on the peat carbon stores of pine mires in  
2154 Finland. *Can. J. For. Res*: 28: 1267–1275.
- 2155 Minkkinen, K., Penttilä, T. & Laine, J. 2007a. Tree stand volume as a scalar for methane fluxes in forestry-  
2156 drained peatlands in Finland. *Boreal Environment Research* 12: 127-132.
- 2157 Minkkinen, K., Laine, J., Shurpali, N., Mäkiranta, P., Alm, J. and Penttilä, T. 2007b. Heterotrophic soil  
2158 respiration in forestry drained peatlands. *Boreal Environment Reserach* 12(2): 115-126.
- 2159 Minkkinen, K., Vasander, H., Jauhiainen, S., Karsisto, M. and Laine, J. 1999. Post-drainage changes in  
2160 vegetation composition and carbon balance in Lakkasuo mire, Central Finland. *Plant and Soil* 207:107–120.
- 2161 Moore, S., Evans, C.D., Page, S.E., Garnett, M.G., Jones, T.G., Freeman, C., Hooijer A., Wiltshire, A. Limin, S.  
2162 Gauci, V. 2013. Deep instability of deforested tropical peatlands revealed by fluvial organic carbon fluxes.  
2163 *Nature*, 493, 660-664..
- 2164 Moore, T.R. 2003. Dissolved organic carbon in a northern boreal landscape. *Global Biogeochem. Cycles*, 17,  
2165 1109, doi: 10.1029/2003GB002050.
- 2166 Moore, T.R. and Clarkson, B.R. 2007. Dissolved organic carbon in New Zealand peatlands. *New Zealand J.*  
2167 *Marine Freshwater Res.* 41: 137-141. Moore, T.R., Matos, L., Roulet, N.T. 2003. Dynamics and chemistry of  
2168 dissolved organic carbon in Precambrian Shield catchments and an impounded wetland. *Can. J. Fish. Aquat.*  
2169 *Sci.* 60: 612-623.
- 2170 Moore, T.R. and Knowles, R. 1990. Methane emissions from fen, bog and swamp peatlands in Quebec.  
2171 *Biogeochemistry* 11, 45–61.
- 2172 Morrison R., Cumming A., Taft H., Page S., Kaduk, J., Harding, R., Jones, D. & Balzter, H. (2013). Carbon  
2173 dioxide budget of a drained and intensively cultivated lowland fen in the East Anglian Fens. In: Emissions of  
2174 greenhouse gases from UK managed lowland peatlands: Report to Defra under project SP1210:  
2175 Lowland peatland systems in England and Wales – evaluating greenhouse gas fluxes and carbon balances.
- 2176 Morrison R., Page, S., Kaduk, J., Acreman, M., Harding, R. & Balzter, H. (2013). Annual CO<sub>2</sub> budget of a  
2177 regenerating ex-arable peatland in the East Anglian Fens. In: Emissions of greenhouse gases from UK  
2178 managed lowland peatlands: Report to Defra under project SP1210: Lowland peatland systems in England  
2179 and Wales – evaluating greenhouse gas fluxes and carbon balances.
- 2180 Mundel, G. 1976. Untersuchungen zur Torfmineralisation in Niedermooren. *Arch. Acker Pflanzenbau Bodenk.*  
2181 20: 669-679.
- 2182 Murdiyarso, D., K. Hergoualc'h, and L.V. Verchot. 2010. Opportunities for reducing GHG emissions in  
2183 Tropical Peatlands. *Proceedings of the American National Academy of Sciences* 107: 19655–19660,  
2184 doi10.1073/pnas.0911966107.
- 2185 NCDENR (1998) Smoke from Peat Fire Could Pose Health Concerns in Craven County.  
2186 <http://www.ehnr.state.nc.us/newsrels/presrels.htm>.
- 2187 Nieveen, J.P., Campbell, D.I., Schipper, L.A., Blair, I.J. 2005. Carbon exchange of grazed pasture on a drained  
2188 peat soil. *Global Change Biology* 11: 607-618.
- 2189 Nilsson, M., Sagerfors, J., Buffam, I., Laudon, H., Eriksson, T., Grelle, A., Klemetsson, L., Weslien, P. &  
2190 Lindroth, A. 2008. Contemporary carbon accumulation in a boreal oligotrophic minerogenic mire - a  
2191 significant sink after accounting for all C-fluxes. *Global Change Biology* 14: 2317–2332.
- 2192 Nouvellon, Y., Laclau, J.-P., Epron, D., Le Maire, G., Bonnefond, J.-M., Gonçalves, J.L.M., Bouillet, J.-P., 2012.  
2193 Production and carbon allocation in monocultures and mixed-species plantations of *Eucalyptus grandis* and  
2194 *Acacia mangium* in Brazil. *Tree Physiology* 32, 680-695.
- 2195 Nykänen, H., Alm, J., La<sup>o</sup>ng, K., Silvola, J. and Martikainen, P. J. 1995. Emissions of CH<sub>4</sub>, N<sub>2</sub>O and CO<sub>2</sub> from a  
2196 virgin fen and a fen drained for Grassland in Finland, *J. Biogeogr.*, 22, 351–357.

Final Draft

- 2197 Nykänen, H., Alm, J., Silvola, J., Tolonen, K. and Martikainen, P. J. 1998. Methane fluxes on boreal peatlands of  
2198 different fertility and the effect of long term experimental lowering of the water table on flux rates, *Glob.*  
2199 *Biogeochemical Cycles*, 12, 53–69.
- 2200 Nykänen, H., Silvola, J., Alm, J. and Martikainen, P. J. 1996. Fluxes of greenhouse gases CH<sub>4</sub>, CO<sub>2</sub> and N<sub>2</sub>O on  
2201 some peat mining areas in Finland, in: *Northern Peatlands in Global Climatic Change*, edited by: Laiho, R.,  
2202 Laine, J. and Vasander, H., *Proceedings of the International Workshop Held in Hyytiälä, Finland*. Publication  
2203 of the Academy of Finland, Helsinki 1/96, 141–147.
- 2204 Ojanen, P., Minkkinen, K. Alm, J. and Penttilä, T. 2010. Soil-atmosphere CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O fluxes in boreal  
2205 forestry-drained peatlands. *Forest Ecology and Management* 260:411-421.
- 2206 Ojanen, P., Minkkinen, K. and Penttilä, T. 2013. The current greenhouse impact of forestry-drained peatlands.  
2207 *Forest Ecology and Management* 289:201-208.
- 2208 Ojanen, P., Minkkinen, K., Lohila, A., Badorek, T. and Penttilä, T. 2012. Chamber measured soil respiration: a  
2209 useful tool for estimating the carbon balance of peatland forest soils? *Forest Ecology and Management*  
2210 277:132-140.
- 2211 Okruszko, H., 1989. Wirkung der Bodennutzung auf die Niedermoorentwicklung. Ergebnisse eines längjährigen  
2212 Feldversuches. *Z f Kulturtechnik und Landentwicklung* 30: 167–176.
- 2213 Opsahl S., Benner, R. 1997. Distribution and cycling of terrigenous dissolved organic matter in the ocean. *Nature*  
2214 386: 480-482.
- 2215 Opsahl S., Benner, R. 1998. Photochemical reactivity of dissolved lignin in river and ocean waters. *Limnol.*  
2216 *Oceanogr.* 43: 1297-1304.
- 2217 Page, S. E., F. Siegert, J. O. Rieley, H.-D. V. Bohm, A. Jaya and S. Limin (2002). The amount of carbon released  
2218 from peat and forest fires in Indonesia during 1997. *Nature* 420: 61-65.
- 2219 Page, S. E., J. O. Rieley and C. J. Banks (2011). Global and regional importance of the tropical peatland carbon  
2220 pool. *Global Change Biology* 17: 798-818.
- 2221 Pawson, R. R., Lord, D. R., Evans, M. G., Allott, T.E.H. (2008). Fluvial organic carbon flux from an eroding  
2222 peatland catchment, southern Pennines, UK. *Hydrol. Earth Syst. Sci.* 12: 625–634.
- 2223 Petersen, S.O., Hoffmann, C.C., Schafer, C.-M, Blicher-Mathiesen, G., Elsgaard, L., Kristensen, K. Larsen, S.E.,  
2224 Torp, S.B., Greve, M.H. 2012. Annual emissions of CH<sub>4</sub>, and N<sub>2</sub>O, and ecosystem respiration, from eight  
2225 organic soils in Western Denmark managed by agriculture. *Biogeosciences* 9: 403-422.
- 2226 Pihlatie, M., Rinne, J., Lohila, A., Laurila, T., Aro, L. and Vesala, T. 2004. Nitrous oxide emissions from an  
2227 afforested peat field using eddy covariance and enclosure techniques. In: Päivänen *et al.* (Eds.), *Proceedings*  
2228 *of 12th International Peat Congress, Tampere, Finland 6–11 Jun 2004, Vol 2*, 1010–1014.
- 2229 Pitkänen, A., J. Turunen and K. Tolonen (1999). The role of fire in the carbon dynamics of a mire, eastern  
2230 Finland. *Holocene* 9: 453-462.
- 2231 Policy, U. S. F. F. M. (2001). Review and Update of the 1995 Federal Wildland Fire Management Policy.
- 2232 Poulter, B., N. L. Christensen and P. N. Halpin (2006). Carbon emissions from a temperate peat fire and its  
2233 relevance to interannual variability of trace atmospheric greenhouse gases. *Journal of Geophysical Research*  
2234 111: D06301, doi:06310.01029/02005JD006455.
- 2235 Rahajoe, J. S., T. Kohyama, and S. H. Limin (2000), Litter decomposition process in two contrastive nutrient  
2236 limited forest types in central Kalimantan, in *Proceedings of the International Symposium on Tropical*  
2237 *Peatlands, Bogor, Indonesia, 22–23 November 1999*, edited by T. Iwakuma et al., pp. 223–231, Hokkaido  
2238 Univ. and Indonesian Inst. of Sci., Bogor, Indonesia.
- 2239 Ramadan, Z., X. H. Song and P. K. Hopke (2000). Identification of sources of Phoenix aerosol by positive  
2240 matrix factorization. *Journal of Air Waste Management* 50: 1308-1320.
- 2241 Rantakari, M., Mattsson, T., Kortelainen, P., Piirainen, S., Finér, L., Ahtiainen, M. 2010. Organic and inorganic  
2242 carbon concentrations and fluxes from managed and unmanaged boreal first-order catchments. *Sci. Total*  
2243 *Environ.*, 408: 1649-1658.
- 2244 Raymond, P.A., Bauer, J.E. 2001. Riverine export of aged terrestrial organicmatter to the North Atlantic Ocean.  
2245 *Nature*, 497, 497-500.

Final Draft

- 2246 Regina, K., Nykänen, H., Maljanen, M., Silvola, J. and Martikainen, P. J. 1998. Emissions of N<sub>2</sub>O and NO and  
2247 net nitrogen mineralization in a boreal forested peatland treated with different nitrogen compounds, *Can. J.*  
2248 *For. Res.*, 28, 132–140.
- 2249 Regina, K., Nykänen, H., Silvola, J. and Martikainen, P. J. 1996. Fluxes of nitrous oxide from boreal peatlands  
2250 as affected by peatland type, water table level and nitrification capacity, *Biogeochemistry*, 35, 401–418.
- 2251 Regina, K., Pihlatie, M., Esala, M. and Alakukku, L. 2007. Methane fluxes on boreal arable soils, *Agr. Ecosyst.*  
2252 *Environ.*, 119, 346–352.
- 2253 Regina, K., Syväsalö, E., Hannukkala, A. and Esala, M. 2004. Fluxes of N<sub>2</sub>O from farmed peat soils in Finland,  
2254 *Eur. J. Soil Sci.*, 55, 591–599.
- 2255 Rein, G., Cleaver, N., Ashton, C., Pironi, P., and Torero, J.L. 2008. The severity of smouldering peat fires and  
2256 damage to the forest soil. *Catena*, 74: 304–309.
- 2257 Roulet, N.T. and Moore, T.R. (1995). The effect of forestry drainage practices on the emission of methane from  
2258 northern peatlands. *Canadian Journal of Forest Research* 25: 491-499.
- 2259 Roulet, N.T., LaFleur, P.M., Richards, P.J., Moore, T.R., Humphreys, E.R., Bubier, J. (2007). Contemporary  
2260 carbon balance and late Holocene carbon accumulation in a northern peatland. *Global Change Biol.* 13, 397-  
2261 411.
- 2262 Rowson, J.G., Gibson, H.S., Worrall, F., Ostle, N., Burt, T.P., Adamson, J.K. (2010). The complete carbon  
2263 budget of a drained peat catchment. *Soil Use and Management* 26: 261-273.
- 2264 Roy, D. P., L. Boschetti, C. O. Justice and J. Ju (2008). The collection 5 MODIS burned area product—Global  
2265 evaluation by comparison with the MODIS activefire product. *Remote Sensing of Environment* 112: 3690-  
2266 3707.
- 2267 Ruser R., Flessa H., Schilling R., Beese F., Munch J.C. 2001. Effect of crop-specific field management and N  
2268 fertilization on N<sub>2</sub>O emissions from a fine-loamy soil. *Nutr. Cycl. Agroecosyst.*, 59, 177–191.
- 2269 Ryden JC, Lund LJ 1980. Nitrous oxide evolution from irrigated land. *J. Environ. Qual.*, 9, 387–393.
- 2270 Saari, P., Saarnio, S., Kukkonen, J. V. K., Akkanen, J., Heinonen, J., Saari, V. and Alm, J. (2009). DOC and  
2271 N<sub>2</sub>O dynamics in upland and peatland forest soils after clear-cutting and soil preparation, *Biogeochemistry*,  
2272 94, 217–231, doi:10.1007/s10533-009-9320-1.
- 2273 Sah, S.P., Jungner, H., Oinonen, M., Kukkola, M., Helmisaari, H-S. (2010). Does the age of fine root carbon  
2274 indicate the age of fine roots in boreal forests? *Biogeochem.* DOI 10.1007/s10533-010-9485-7
- 2275 Saharjo, B. H. and Nurhayati, A. D. (2005). Changes in chemical and physical properties of hemic peat under  
2276 fire-based shifting cultivation. *Tropics* 14: 263-269.
- 2277 Saharjo, B. H. and Munoz, C. P. (2005). Controlled burning in peat lands owned by small farmers: a case study  
2278 in land preparation. *Wetlands Ecology and Management* 13: 105-110.
- 2279 Schothorst C.J. (1977). Subsidence of low moor peat soils in the Western Netherlands. *Proc of 5<sup>th</sup> Int Peat*  
2280 *Congress, Poznan*, 1: 206–217.
- 2281 Schrier-Uijl, A.P., Kroon, P.S., Hensen, A., Leffelaar, P.A., Berendse, F. & Veenendaal, E.M. (2010a).  
2282 Comparison of chamber and eddy covariance based CO<sub>2</sub> and CH<sub>4</sub> emission estimates in a heterogeneous grass  
2283 ecosystem on peat, *Agric. For. Meteorol.*, doi:10.1016/j.agrformet.2009.11.007.
- 2284 Schrier-Uijl, A.P., Hendriks, D.M.D., Kroon, P.S., Hensen, A., van Huissteden, J., Leffelaar, P. A., Nol, L.,  
2285 Veenendaal, E.M. and Berendse, F. (2010b). Agricultural peat lands; towards a greenhouse gas sink – a  
2286 synthesis.
- 2287 Schrier-Uijl, A.P., Hendriks, D.M.D., Kroon, P.S., Hensen, A., van Huissteden, J., Leffelaar, P. A., Nol, L.,  
2288 Veenendaal, E.M. and Berendse, F. (2010c). Flushing meadows - The influence of management alternatives  
2289 on the greenhouse gas balance of fen meadow areas, *Academic Thesis, Wageningen University*
- 2290 Schrier-Uijl, A.P., Kroon, P.S., Leffelaar, P.A., van Huissteden, J.C., Berendse, F. and Veenendaal, E.M. 2010.  
2291 Methane emissions in two drained peat agro-ecosystems with high and low agricultural intensity. *Plant and*  
2292 *Soil* (2010) 329: 509-520. DOI 10.1007/s11104-009-0180-1.
- 2293 Schrier-Uijl, A.P., Veraart, A.J., Leffelaar, P.J., Berendse, F., Veenendaal, E.M. 2011. Release of CO<sub>2</sub> and CH<sub>4</sub>  
2294 from lakes and drainage ditches in temperate wetlands. *Biogeochemistry*, 102: 265–279.

Final Draft

- 2295 Shimamura, T., Momose, K. (2005) Organic matter dynamics control plant species coexistence in a tropical peat  
2296 swamp forest. *Proceedings of the Royal Society B* 272, 1503-1510.
- 2297 Shurpali, N. J., Hyvönen, N. P., Huttunen, J. T., Biasi, C., Nykänen, H., Pekkarinen, N. and Martikainen, P. J.  
2298 2008. Bare soil and reed canary grass ecosystem respiration in peat extraction sites in Eastern Finland, *Tellus*,  
2299 60B, 200–209.
- 2300 Shurpali, N. J., Hyvönen, N., Huttunen, J. T., Clement, R., Reichstein, M., Nykänen, H., Biasi, C. and  
2301 Martikainen, P. J. 2009. Cultivation of perennial grass for bioenergy use on a boreal organic soil – carbon  
2302 sink or source?, *Glob. Change Biol. Bioenerg.*, 1, 35–50, doi:10.1111/j.1757.2009.01003.x.
- 2303 Sikström, U., Björk, R. G., Ring, E., Ernfors, M., Jacobson, S., Nilsson, M. and Klemetsson, L. 2009. Tillförsel  
2304 av aska i skog på dikad torvmark i södra Sverige. Effekter på skogsproduktion, flöden av växthusgaser,  
2305 torvegenskaper, markvegetation och grundvattenkemi. VÄRMEFORSK Service AB, Stockholm, 75 pp., (in  
2306 Swedish).
- 2307 Simola, H., Pitkänen, A. & Turunen, J., 2012. Carbon loss in drained forestry peatlands in Finland, estimated by  
2308 re-sampling peatlands surveyed in the 1980s. *European Journal of Soil Science*, Dezember 2012, 63, 798-807.  
2309 DOI: 10.1111/j.1365-2389.2012.01499.x.
- 2310 Simon, M., S. Plummer, F. Fierens, J. J. Hoelzemann and O. Arino (2004). Burnt area detection at global scale  
2311 using ATSR-2: The GLOBSCAR products and their qualification. *Journal of Geophysical Research* **109**:  
2312 D14S02, doi:10.1029/2003JD003622.
- 2313 Sinsabaugh, R.L., Findlay, S. 1995. Microbial production, enzyme activity, and carbon turnover in surface  
2314 sediments of the Hudson River estuary. *Microb. Ecol.* 30: 127-141.
- 2315 Sobek, S., Algesten, G., Bergström, A-K, Jansson, M., Tranvik, J. (2003). The catchment and climate regulation  
2316 of pCO<sub>2</sub> in boreal lakes. *Global Change Biol.* 9, 630-641.
- 2317 Soja, A. J., W. R. Cofer, H. H. Shugart, A. I. Sukhinin, P. W. Stackhouse, D. J. McRae and S. G. Conard (2004).  
2318 Estimating fire emissions and disparities in boreal Siberia (1998-2002). *Journal of Geophysical Research-  
2319 Atmospheres* 109: D14S06, doi:10.1029/2004JD004570.
- 2320 Stehfest E., Bouwman L. (2006). N<sub>2</sub>O and NO emission from agricultural fields and soils under natural  
2321 vegetation: summarizing available measurement data and modelling of global annual emissions *Nutrient  
2322 Cycling in Agroecosystems* (2006) 74, 207-228.
- 2323 Strack, M., Waddington, J. M. and Tuittila, E.-S. 2004. Effect of water table drawdown on northern peatland  
2324 methane dynamics: Implications for climate change. *Global Biogeochem. Cycles*, 18, GB4003,  
2325 doi:10.1029/2003GB002209.
- 2326 Strack, M., Waddington, J.M., Bourbonniere, R.A., Buckton, L., Shaw, K. Whittington, P., Price, J.S. (2008).  
2327 Effect of water table drawdown on peatland dissolved organic carbon export and dynamics. *Hydrol. Process.*  
2328 22: 3373-3385
- 2329 Strack, M. & Zuback, Y.C.A. (2013). Annual carbon balance of a peatland 10 yr following restoration.  
2330 *Biogeosciences* 10, 2885–2896
- 2331 Stutter, M.I., Baggaley, N., Barry, C., Chapman, S., Dawson, J.J.C., Helliwell, R.C., Higgins, A., Howden, L.,  
2332 Jackson-Blake, L., Lumsdon, D.G., Malcolm, I., Sample, J., Potts, J., Worrall, F. 2011. Assessment of the  
2333 contribution of aquatic carbon fluxes to carbon losses from UK peatlands. SNIFFER Report, Project ER18,  
2334 James Hutton Institute, Aberdeen, 194 pp.
- 2335 Sulistiyanto, Y. (2004), Nutrient dynamics in different sub-types of peat swamp forest in central Kalimantan,  
2336 Indonesia, Ph.D. thesis, 351 pp., Univ. of Nottingham, Nottingham, U. K.
- 2337 Sundari, S., Hirano, T., Yamada, H., Kusin, K., Limin, S. (2012) Effects of groundwater level on soil respiration  
2338 in tropical peat swamp forests. *J. Agric. Meteorol.* 68, 121-134.
- 2339 Sundh, I., Nilsson, M., Mikkelä, C., Granberg, G. and Svensson, B. H. 2000. Fluxes of methane and carbon  
2340 dioxide on peat-mining areas in Sweden, *Ambio*, 29, 499–503.
- 2341 Swetnam, T. W. (1993). Fire history and climate change in giant sequoia groves. *Science* 262: 885-889.
- 2342 Taft, H., Cross, P., Jones, D. 2013. Annual emission cycle of greenhouse gases from peat soils managed for  
2343 horticultural production. In: *Emissions of greenhouse gases from UK managed lowland peatlands: Report to  
2344 Defra under project SP1210: Lowland peatland systems in England and Wales – evaluating greenhouse gas  
2345 fluxes and carbon balances.*

Final Draft

- 2346 Takakai F, Morishita T, Hashidoko Y, Darung U, Kuramochi K, Dohong S, Limin SH, and Hatano R 2006.  
2347 Effects of agricultural landuse change and forest fire on N<sub>2</sub>O emission from tropical peatlands, Central  
2348 Kalimantan, Indonesia. *Soil Sc. Plant Nutr.* 53:662-674.
- 2349 Tansey, K., Grégoire, J.-M., Defourny, P., Leigh, R., Pekel, J.-F., van Bogaert, E., and Bartholomé, E. (2008). A  
2350 new, global, multi-annual (2000–2007) burnt area product at 1 km resolution. *Geophysical Research Letters*  
2351 35: L01401, doi:01410.01029/02007GL031567.
- 2352 Teh, Y.A., Silver, W.L., Sonnentag, O., Detto, M., Kelly, M., Baldocchi, D.D. 2011. Large greenhouse gas  
2353 emissions from a temperate peatland pasture.
- 2354 Treat CC, Bubier JL, Varner RK, Crill PM. 2007. Timescale dependence of environmental and plant-mediated  
2355 controls on CH<sub>4</sub> flux in a temperate fen. *Journal of Geophysical Research. G, Biogeosciences* 112, G01014.
- 2356 Tuittila, E.-S. and Komulainen, V.-M. 1995. Vegetation and CO<sub>2</sub> balance in an abandoned harvested peatland in  
2357 Aitoneva, southern Finland, *Suo*, 46, 69–80.
- 2358 Tuittila, E. S., Komulainen, V. M. Vasander, H. Nykänen, H. Martikainen, P. J., Laine, J. 2000. Methane  
2359 dynamics of a restored cut-away peatland, *Global Change Biol.*, 6, 569-581.
- 2360 Tuittila E-S. Vasander, H. and Laine J. 2004. Sensitivity of C Sequestration in Reintroduced Sphagnum to  
2361 Water-Level Variation in a Cutaway Peatland. *Restoratipon Ecology Vol 12 No 4 pp.* 483-493.
- 2362 Turetsky, M. R. and R. K. Wieder (2001). A direct approach to quantifying organic matter lost as a result of  
2363 peatland fire. *Canadian Journal of Forest Research* 31: 363-366.
- 2364 Turetsky, M. R., Kane, E. S., Harden, J. W., Ottmar, R. D., Manies, K. L. Hoy, E., and Kasischke, E. S. (2011 a).  
2365 Recent acceleration of biomass burning and carbon losses in Alaskan forests and peatlands. *Nature*  
2366 *Geoscience* 4: 27–31.
- 2367 Turetsky, M. R., Donahue, W. F., and Benscoter, B. W. (2011b). Experimental drying intensifies burning and  
2368 carbon losses in a northern peatland. *Nature Communications* 2:514: DoI: 10.1038/ncomms1523.
- 2369 Turunen, J., Roulet, N.T., Moore, T.R. (2004). Nitrogen deposition and increased carbon accumulation in  
2370 ombrotrophic peatlands in eastern Canada. *Global Biogeochem. Cycl.* 18, GB3002.
- 2371 Urban, N.R., Bayley, S.E., Eisenreich, S.J., 1989. Export of dissolved organic carbon and acidity from peatlands.  
2372 *Water Resour. Res.* 25: 1619-1628.
- 2373 Urbanová, Z., Pícek, T, Bárta, J. 2011. Effect of peat re-wetting on carbon and nutrient fluxes, greenhouse gas  
2374 production and diversity of methanogenic archaeal community. *Ecol. Engineering* 37: 1017-1026.
- 2375 Usop, A., Y. Hashimoto, H. Takahashi and H. Hayasaka (2004). Combustion and thermal characteristics of peat  
2376 fire in tropical peatland in Central Kalimantan, Indonesia. *Tropics* 14: 1-19.
- 2377 Val Martin, M., Honrath, R.E., Owen, R.C., Pfister, G., Fialho, P., and Barata, F. (2006). Significant  
2378 enhancements of nitrogen oxides, black carbon, and ozone in the North Atlantic lower free troposphere  
2379 resulting from North American boreal wildfires. *Journal of Geophysical Research* 111: D23S60,  
2380 doi:10.1029/2006JD007530.
- 2381 van Beek CL, Pleijter M, Jacobs CMJ, Velthof GL, van Groenigen JW and Kuikman J. 2010. Emissions of N<sub>2</sub>O  
2382 from fertilized and grazed Grassland on organic soil in relation to groundwater level. *Nutr Cycl Agroecosyst*  
2383 (2010) 86:331-340. DOI 10.1007/s10705-009-9295-2.van den Pol-van Dasselaar A., van Beusichem M.L and  
2384 Oenema O. 1999c. Effects of nitrogen input and grazing on methane fluxes of extensively and intensively  
2385 managed Grasslands in the Netherlands. *Biol. Fertil. Soils* 29: 24-30.
- 2386 van den Pol-van Dasselaar, A. 1998. Methane emissions from Grassland. PhD thesis. Wageningen Agricultural  
2387 University, Wageningen, The Netherlands. 179 pp.
- 2388 van den Pol-van Dasselaar, A., van Beusichem, M.L. and Oenema, O. 1999b. Determinants of spatial variability  
2389 of methane emissions from wet Grasslands on peat soil. *Biogeochemistry* 44: 221–237.
- 2390 van den Pol-van Dasselaar, A., van Beusichem, M.L. and Oenema, O. 1999a. Methane emissions from wet  
2391 Grasslands on peat soil in a nature preserve. *Biogeochemistry* 44, 205–220.
- 2392 van Huissteden J., van den Bos R. & Martikonvena Alvarez. 2006. Modelling the effect of water-table  
2393 management on CO<sub>2</sub> and CH<sub>4</sub> fluxes from peat soils. *Netherlands Journal of Geosciences* 85 – 1; 3 – 18.
- 2394 van Wagtendonk, J. W., R. R. Root and C. H. Key (2004). Comparison of AVIRIS and Landsat ETM+ detection  
2395 capabilities for burn severity. *Remote Sensing of Environment* 92: 397-408.

Final Draft

- 2396 Veenendaal, E.M., Kolle, O., Leffelaar, P.A., Schrier-Uijl, A.P., Van Huissteden, J., Van Walsem, J., Moller, F.,  
2397 and Berendse, F. 2007. CO<sub>2</sub> exchange and carbon balance in two grassland sites on eutrophic drained peat  
2398 soils. *Biogeosciences* 4:1027-1040.
- 2399 Velthof G. L., Brader, A. B., Oenema O.1996. Seasonal variations in nitrous oxide losses from managed  
2400 grasslands in The Netherlands. *Plant Soil*, 181(2): 263-274.
- 2401 Verchot, L.V., Davidson, E.A. Cattânio, J.H. Ackerman, I.L. Erickson, H.E. and Keller, M. 1999. Land-use  
2402 change and biogeochemical controls of nitrogen oxide emissions from soils in eastern Amazonia. *Global*  
2403 *Biogeochemical Cycles*. 13: 31-46.
- 2404 Verchot, L.V., Davidson, E.A. Cattânio, J.H. and Ackerman, I.L.. 2000. Land-use change and biogeochemical  
2405 controls of methane fluxes in soils of eastern Amazonia. *Ecosystems*. 3: 41-56  
2406 Verchot, L.V., L. Hutabarat,  
2407 K. Hairiah and M. van Noordwijk. 2006. Nitrogen Availability and Soil N<sub>2</sub>O Emissions Following  
2408 Conversion of Forests to Coffee in Southern Sumatra. *Global Biogeochemical Cycles*. 20: GB4008,  
doi10.1029/2005GB002469.
- 2409 Vermaat, J.E., Hellmann, F., Dias, A.T.C., Hoorens, B., van Logtestijn, R.S.P., Aerts, R. 2011. Greenhouse gas  
2410 fluxes from Dutch peatland water bodies: Importance of the surrounding landscape. *Wetlands*, 31: 493-498.
- 2411 von Arnold, K., Nilsson, M., Hånell, B., Weslien, P. and Klemedtsson, L. 2005a. Fluxes of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O  
2412 from drained organic soils in deciduous forests. *Soil Biology and Biochemistry*, 37:1059-1071.
- 2413 von Arnold, K., Hånell, B., Stendahl, J. and Klemedtsson, L. 2005c. Greenhouse gas fluxes from drained organic  
2414 Forest Land in Sweden. *Scandinavian Journal of Forest Research*, 20:5, 400 - 411. DOI:  
2415 10.1080/02827580500281975.
- 2416 von Arnold, K., Weslien, P., Nilsson, M., Svensson, B. H. and Klemedtsson, L. 2005b. Fluxes of CO<sub>2</sub>, CH<sub>4</sub> and  
2417 N<sub>2</sub>O from drained coniferous forests on organic soils, *Forest Ecol. Manage.*, 210, 239–254.
- 2418 Waddington, J.M., and Day, S.M. 2007. Methane emissions from a peatland following restoration. *J. Geophys.*  
2419 *Res.* 112, doi:10.1029/2007JG000400.
- 2420 Waddington, J.M., Strack, M. and Greenwood, M.J. 2010. Toward restoring the net carbon sink function of  
2421 degraded peatlands: Short-term response in CO<sub>2</sub> exchange to ecosystem-scale restoration. *J. Geophys. Res.*  
2422 115, doi:10.1029/2009JG001090.
- 2423 Wallage, Z.E., Holden, J., McDonald, A.T. 2006. Drain blocking: An effective treatment for reducing dissolved  
2424 organic carbon loss and water discoloration in a drained peatland. *Sci. Total Environ.* 367: 811-821.
- 2425 Wallin, M.B., Öquist, M.G., Buffam, I., Billett, M.F., Nisell, J., Bishop, K.H. (2011) Spatiotemporal variability  
2426 of the gas transfer coefficient (KCO<sub>2</sub>) in boreal streams; implications for large scale estimates of CO<sub>2</sub> evasion.  
2427 *Global Biogeochem. Cycles* GB3025. doi:10.1029/2010GB003975
- 2428 Ward, D. E. and C. C. Hardy (1984). Advances in the characterisation and control of emissions from prescribed  
2429 fires. 77th Annual Meeting of the Air Pollution Control Association. San Francisco, California.
- 2430 Ward, S.E., Bardgett, R.D., McNamara, N.P., Adamson, J.K., Ostle, N.J. 2007. Long-term consequences of  
2431 grazing and burning on northern peatland carbon dynamics. *Ecosystems* 10: 1069–1083.
- 2432 Warren, M.W., Kauffmann, J.B., Murdiyarto, D., Anshari, G., Hergoualc'h, K., Kurnianto, S., Purbopuspito, J.,  
2433 Gusmayanti, E., Afifudin, M., Rahajoie, J., Alhamd, L., Limin, S., Iswandi, A. (2012) A cost-efficient  
2434 method to assess carbon stocks in tropical peat soil. *Biogeosciences* 9, 4477-4485.
- 2435 Watanabe, A., Purwanto, B.H., Ando, H., Kakuda, K.-i., Jong, F.-S., (2009). Methane and CO<sub>2</sub> fluxes from an  
2436 Indonesian peatland used for sago palm (*Metroxylon sagu* Rottb.) cultivation: Effects of fertilizer and  
2437 groundwater level management. *Agriculture, Ecosystems and Environment* 134, 14-18.
- 2438 Weinzierl W. (1997). Niedermoore in Baden-Württemberg – Bilanzierung der CO<sub>2</sub>-Emission am Beispiel des  
2439 Donaurieds. *Mitteilungen der Deutschen Bodenkundlichen Gesellschaft* 85: 1059–1062.
- 2440 Weslien, P., Kasimir Klemedtsson, Å , Börjesson, G. and Klemedtsson, L. 2009. Strong pH influence on N<sub>2</sub>O and  
2441 CH<sub>4</sub> fluxes from forested organic soils, *Eur. J. Soil Sci.*, 60, 311–320, doi:10.1111/j.1365-  
2442 2389.2009.01123.x.
- 2443 Wickland, K.P., Neff, J.C., Aiken, G.R. 2007. Dissolved organic carbon in Alaskan boreal forest: Sources,  
2444 chemical characteristics, and biodegradability. *Ecosystems* 10: 1323-1340.
- 2445 Wild, U., Kampp, T., Lenz, A., Heinz, S. And Pfadenhauer, J. 2001. Cultivation of *Thpha* spp. In constructed  
2446 wetlands for peatland restoration. *Ecological Engineering* 17:49-54.

Final Draft

- 2447 Worrall F., Armstrong, A. Adamson, J.K. 2007. The effects of burning and sheep-grazing on water table depth  
2448 and soil water quality in a upland peat. *J. Hydrol.* 339: 1-14.
- 2449 Worrall, F., G. Clay, R. Marrs and M. S. Reed (2010). Impacts of burning management on peatlands. Report to  
2450 IUCN UK Peatland Programme, Edinburgh.
- 2451 Worrall, F., Moody, C., Jones, T., Evans, C., (2013). Conversion of waterborne DOC to CO<sub>2</sub> – results of field  
2452 experiments. In: Emissions of greenhouse gases associated with peatland drainage waters: Report to Defra  
2453 under project SP1205: Greenhouse Gas Emissions Associated with Non Gaseous Losses of Carbon from  
2454 Peatlands - fate of Particulate and Dissolved Carbon. Report to the Department of Environment, Food and  
2455 Rural Affairs, UK.
- 2456 Worrall, F., Rowson, J.G., Evans, M.G., Pawson, R., Daniels, S., Bonn, A. (2011). Carbon fluxes from eroding  
2457 peatlands – the carbon benefit of revegetation following wildfire. *Earth Surface Process. Landforms* 36:  
2458 1487–1498.
- 2459 Wösten, J.M.H., Ismail, A.B., van Wijk, A.L.M., (1997). Peat subsidence and its practical implications: a case  
2460 study in Malaysia. *Geoderma* 78, 25-36. Yallop, A.R., Clutterbuck, B., Thacker, J. 2010. Increases in humic  
2461 dissolved organic carbon export from upland peat catchments: the role of temperature, declining sulfate  
2462 deposition and changes in land management. *Climate Res.* 45: 43-56. Yallop, A.R., Clutterbuck, B., Thacker,  
2463 J., (2010). Increases in humic dissolved organic carbon export from upland peat catchments: the role of  
2464 temperature, declining sulphur deposition and changes in land management. *Climate Research* 45, 43-56.
- 2465 Yamulki, S., Anderson, R. Peace, A., and Morison, J.I.L. 2013. Soil CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O fluxes from an afforested  
2466 lowland raised peatbog in Scotland: implications for drainage and restoration. *Biogeosciences* 10:1051-1065.
- 2467 Yefremova, T. T. and S. P. Yefremov (1996). Ecological effects of peat fire on forested bog ecosystems. *Fire in*  
2468 *ecosystems of boreal Eurasia.* J. G. Goldammer and V. V. Furyaev. Netherlands, Kluwer Academic  
2469 Publishers: 350-357.
- 2470 Yokelson, R. J., Burling, I. R., Gilman, J. B., Warneke, C., Stockwell, C. E., de Gouw, J., Akagi, S. J., Urbanski,  
2471 S. P., Veres, P., Roberts, J. M., Kuster, W. C., Reardon, J., Griffith, D. W. T., Johnson, D. T., Hosseini, S.,  
2472 Miller, J. W., Cocker III, D. R., Jung, D. and Weise, D. R. 2013. Coupling field and laboratory  
2473 measurements to estimate the emission factors of identified and unidentified trace gases for prescribed fires.  
2474 *Atmospheric Chemistry and Physics* **13**: 89-116.
- 2475 Yokelson, R.J., Karl, T., Artaxo, P., Blake, D. R., Christian, T.J., Griffith, D.W.T., Guenther, A., and Hao, W.M.  
2476 (2007). The tropical forest and fire emissions experiment: overview and airborne fire emission factor  
2477 measurements. *Atmos. Chem. Phys.*, 7, 5175–5196.
- 2478 Yokelson, R. J., R. Susott, D. E. Ward, J. Reardon and D. W. T. Griffith (1997). Emissions from smoldering  
2479 combustion of biomass measured by open-path Fourier transform infrared spectroscopy. *Journal of*  
2480 *Geophysical Research* 102: 18865-18877.
- 2481 Yule, C.M., Gomez, L.N. 2009. Leaf litter decomposition in a tropical peat swamp forest in Peninsular Malaysia.  
2482 *Wetlands Ecol. Manage.* 17: 231–241
- 2483 Zoltai, S. C., Morrissey, L. A., Livingston, G. P. and de Groot, W. J. (1998). Effects of fires on carbon cycling  
2484 in North American boreal peatlands. *Environmental Reviews* 6: 13-24.
- 2485 Zulkifli, Y. 2002. Hydrological attributes of a disturbed peat swamp forest. In: Parish F, Padmanabhan E, Lee  
2486 CL, Thang HC (eds) Prevention and control of fire in peatlands. Proceedings of workshop on prevention and  
2487 control of fire in peatlands, 19–21 March 2002, Kuala Lumpur. Global Environment Centre & Forestry  
2488 Department Peninsular Malaysia. Cetaktama, Kuala Lumpur, pp 51–5.
- 2489